

Nontrivial edge coupling from a Dirichlet network squeezing: the case of a bent waveguide

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2007 J. Phys. A: Math. Theor. 40 F511

(<http://iopscience.iop.org/1751-8121/40/26/F02>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.109

The article was downloaded on 03/06/2010 at 05:17

Please note that [terms and conditions apply](#).

FAST TRACK COMMUNICATION

Nontrivial edge coupling from a Dirichlet network squeezing: the case of a bent waveguide

Claudio Cacciapuoti¹ and Pavel Exner^{2,3}¹ Institut für Angewandte Mathematik, Universität Bonn, Wegelerstr. 6, 53115 Bonn, Germany² Nuclear Physics Institute, Czech Academy of Sciences, 25068 Řež near Prague, Czech Republic³ Doppler Institute, Czech Technical University, Břehová 7, 11519 Prague, CzechiaE-mail: caccia@na.infn.it and exner@ujf.cas.cz

Received 23 April 2007, in final form 16 May 2007

Published 12 June 2007

Online at stacks.iop.org/JPhysA/40/F511**Abstract**

In distinction to the Neumann case, the squeezing limit of a Dirichlet network leads in the threshold region generically to a quantum graph with disconnected edges, exceptions may come from threshold resonances. Our main point in this paper is to show that modifying locally the geometry we can achieve in the limit a nontrivial coupling between the edges including, in particular, the class of δ -type boundary conditions. We work out an illustration of this claim in the simplest case when a bent waveguide is squeezed.

PACS numbers: 03.65.Db, 73.21.Hb

Mathematics Subject Classification: 81V99, 35J10, 34L40

1. Introduction

Quantum mechanics on graphs attracted a lot of attention recently—let us just mention the preceding volume [BCFK06] as a guide to the abundant bibliography in the field. The interest has different sources, important among them are numerous existing and potential applications. While simple and versatile, however, quantum graph models have a problem: in a sense they offer too much freedom. The requirement of self-adjointness, or probability current conservation, determines a class of boundary conditions which connect the column vectors Ψ and Ψ' of boundary values of the wavefunctions and their derivatives at a given graph vertex. Following [Ha00, KS00] these conditions can be cast into the unique form

$$(U - I)\Psi + i(U + I)\Psi' = 0, \quad (1.1)$$

where U is an $n \times n$ unitary matrix, with n being the number of edges sprouting of the vertex.

Hence, a vertex coupling contains n^2 free parameters to be fixed. Asking about the meaning of various couplings one can naturally get useful insights by obtaining boundary conditions through limit of families of (regular or singular) interactions on the graph

[E96, CE04, ET07]. A proper understanding of the problem requires, however, to find an interpretation of the coupling in terms of models without free parameters; a natural idea is to investigate the motion of a free quantum particle on a system of thin tubes which shrink towards the graph.

This is a longstanding and nontrivial problem and the answer depends substantially on the network dynamics we work with. In the case when the Hamiltonian (which can be identified with the Laplacian by a suitable choice of units) refers to tubes with Neumann boundary, the limit yields typically quantum graphs with free boundary conditions (often called not quite appropriately Kirchhoff) at the vertices, described by $U = \frac{2}{n} \mathcal{J} - I$, where \mathcal{J} is the $n \times n$ matrix whose all entries are equal to one—see [FW93, KZ01, RS01, Sa01], the same is true also for shrinking families of ‘sleeve’ manifolds without a boundary [EP05]; the convergence is norm-resolvent [Po05a] and the conclusion extends to resonances on such structures [EP07].

From the viewpoint of applications to semiconductor structures and similar objects, however, it is the case of Dirichlet (hard-wall) boundary which is more important. It is very different from its Neumann counterpart and more difficult and relevant results started appearing only recently. The main source of the difficulties is that in the Dirichlet case geometric perturbations such as bending, ‘swelling’, twisting or branching give rise to an effective interaction—see, e.g., [EŠ89, DE95, EEK05] and references therein—sometimes attractive, sometimes repulsive, which changes spectral and scattering properties of such networks. Even the statement of the problem is more complicated than in the Neumann case where we naturally investigate spectrum around the zero value which is the continuum threshold. The analogous quantity in a Dirichlet network blows up to infinity and one has to choose the reference point by a suitable energy renormalization. In most cases, with the notable exception of the recent paper [MV06], the attention is paid to the vicinity of the ‘running’ threshold.

It is generally conjectured that the generic limit in the vicinity of the threshold corresponds to a fully decoupled graph with Dirichlet conditions at the edge endpoints. This is obvious if the vertex regions squeeze faster than tubes as in [Po05b] but it is expected to hold even without such an additional effective repulsion. To see the reason one has to realize that the problem at hand can be by scaling rephrased as analysis of a network of tubes of constant cross section whose overall size grow. This means that the distances between the ‘vertices’ tend to infinity and the character of solutions to the Schrödinger equation is determined by its asymptotic properties around each single vertex. Away of the threshold, as in the particular case of [MV06] cited above, the limiting boundary conditions are generally nontrivial and given by the scattering properties of the ‘fat star’ region. Around the threshold, on the other hand, the scattering is generically suppressed in view of the mentioned effective interaction—see a brief discussion in [MV06], the forthcoming paper [Gr07] and also a related problem with Dirichlet boundary replaced by a confining potential in [DT06]—leading to the Dirichlet decoupling.

This limiting behaviour is not universal, though. The situation changes if the described system associated with the vertex possesses a threshold resonance. The simplest case where a nontrivial effect of this type can be observed is a bent tube which squeezes in the limit to a graph of two half-line edges joined in a single vertex [ACF07]: in the presence of a threshold resonance one gets the line with a point interaction of scale-invariant type, cf [HC06] and references therein. More general results of that type were announced in [Gr07].

Our main point in this paper is to argue that one can go further and construct classes of squeezing Dirichlet networks which produce wider families of vertex coupling when renormalized to the continuum threshold, including those with nonempty discrete spectrum or resonances. The procedure we propose consists of two steps:

- (i) choose a network collapsing to a graph in such a way that the limit Hamiltonian has a threshold resonance⁴,
- (ii) change the scaling properties of the vertex region slightly, typically by adding higher order terms in the scaling parameter.

The modification in point (ii) can be achieved in various ways; for instance, one can ‘wiggle’ the edges angles or scale the vertex volume region at a rate which differs from that of the ‘edge tubes’ by a higher order term, a combination of such perturbations, etc. Incidentally, the same effect can also be obtained by introducing suitable potentials into the vertex region, but a purely geometric way is probably the most interesting.

Note that the described approximation follows the same scheme which one uses when interpreting pseudopotentials, or point interactions in dimensions two and three, by a suitable nonlinear scaling starting from a threshold resonance [AGHH05]. On the other hand, there is a large difference between the two cases coming from the fact that the geometric approximation discussed here covers not a single operator class but a broad variety of systems. Consequently, the proposal made above has a status of a *conjecture* and the corresponding procedure must be made concrete and worked out properly in each particular case.

To show that this programme is not void we are going in the rest of the paper to perform this task for the bent-waveguide system studied in [ACF07]. We will show that by modifying the bending angle around a threshold resonance value we can arrive in the limit at a two-parameter class of point interactions on the line including the important particular case of the δ interaction. In our model, the system has multiple threshold resonances and the limiting procedure described above can be associated with each of them—we expect that this is a standard behaviour of networks with Dirichlet boundary. We will describe the model and state the results in the next section. Then we extend the analysis of short-range potentials from [ACF07] to more general scaling, and in the last section we will prove our main theorem.

2. The bent-waveguide model and the results

As we have said we use the model studied in [ACF07], namely a planar waveguide of constant width obtained by ‘fattening’ a fixed smooth curve along its normal. We denote by C a curve embedded in \mathbb{R}^2 , i.e. $C := \{(x, y) \in \mathbb{R}^2 | x = \gamma_1(s), y = \gamma_2(s), s \in \mathbb{R}\}$, assuming that it is parameterized by its arc length, $\gamma_1'^2 + \gamma_2'^2 = 1$. Moreover, we denote by $\gamma(s)$ the signed curvature of C ,

$$\gamma(s) := \gamma_2'(s)\gamma_1''(s) - \gamma_1'(s)\gamma_2''(s);$$

it completely characterizes the curve C up to Euclidean transformations and the curvature radius at a given point is given by $r = |\gamma|^{-1}$. We suppose that the curve is not self-intersecting, i.e., it has no loops, and for simplicity we consider only curves with a compactly supported curvature. The last assumption means that the curve C is made up of two straight half lines joined by a smooth curve. In particular, the (overall) bending angle of C is the angle between the two half lines, which is equal to $\theta = \int_{\mathbb{R}} \gamma(s) ds$.

The above-mentioned fat curve which is our waveguide is the open set $\Omega \subset \mathbb{R}^2$ defined as $\Omega := \{(x, y) \in \mathbb{R}^2 | x = \gamma_1(s) - u\gamma_2'(s), y = \gamma_2(s) + u\gamma_1'(s), s \in \mathbb{R}, u \in (-d, d)\}$,

where s and u represent a global system of coordinates in strip, with s being the coordinate along the curve while u is the distance along the normal to C . The width of the waveguide

⁴ The original network Hamiltonian at that may or may not have such a resonance, depending on the limiting procedure used. A threshold resonance may be present, e.g., in the leading term of the perturbation expansion w.r.t. the squeezing parameter as in the model discussed below.

is constant and equal to $2d$ where $d > 0$. Another standard assumption we made is that d is smaller than the curvature radius, $d\|\gamma\|_\infty < 1$. The closure of Ω is conventionally denoted by $\overline{\Omega}$. The (negative) Laplacian with Dirichlet boundary conditions on $\partial\overline{\Omega}$, denoted as $-\Delta_\Omega^D$, is the Friedrichs extension of the positive, symmetric operator $L_0 := -\Delta$ with $\mathcal{D}(L_0) := C_0^\infty(\Omega)$.

The main geometric object of our study will be a family of waveguides whose shape and width depend on a scaling parameter $0 < \varepsilon \leq 1$ according to

$$\gamma_\varepsilon(s) := \frac{\sqrt{\lambda(\varepsilon)}}{\varepsilon} \gamma\left(\frac{s}{\varepsilon}\right); \quad d_\varepsilon := \varepsilon^\alpha d, \quad \text{with } \alpha > 1, \quad (2.1)$$

where $\lambda(\varepsilon)$ is a fixed function to be specified below; its presence is the main difference comparing to [ACF07] because of the term $\sqrt{\lambda(\varepsilon)}$. We suppose that $\lambda(\varepsilon)$ is real, positive and analytic near the origin and, moreover, that it expands as

$$\lambda(\varepsilon) = 1 + \lambda_1 \varepsilon + \mathcal{O}(\varepsilon^2) \quad \text{with } \lambda_1 = \lambda'(0). \quad (2.2)$$

We assume that the condition $d_\varepsilon\|\gamma_\varepsilon\|_\infty < 1$ is satisfied for every $0 < \varepsilon \leq \varepsilon_0$ with some $\varepsilon_0 > 0$. The scaling (2.1) gives rise to a family of curves, $C_\varepsilon := \{(x, y) \in \mathbb{R}^2 \mid x = \gamma_{\varepsilon,1}(s), y = \gamma_{\varepsilon,2}(s), s \in \mathbb{R}\}$, the bending angle θ_ε of which changes slightly with respect to ε ,

$$\theta_\varepsilon = \int_{\mathbb{R}} \gamma_\varepsilon(s) \, ds = \theta \sqrt{\lambda(\varepsilon)} = \theta \left(1 + \frac{1}{2} \lambda_1 \varepsilon\right) + \mathcal{O}(\varepsilon^2).$$

They in turn generate a family of bent waveguides, i.e. domains Ω_ε defined by

$$\Omega_\varepsilon := \{(x, y) \in \mathbb{R}^2 \mid x = \gamma_{\varepsilon,1}(s) - u\gamma'_{\varepsilon,2}(s), y = \gamma_{\varepsilon,2}(s) + u\gamma'_{\varepsilon,1}(s), s \in \mathbb{R}, u \in (-d_\varepsilon, d_\varepsilon)\}.$$

In the limit $\varepsilon \rightarrow 0$, the strip family shrinks to a graph, denoted by \mathcal{G} , made up of two edges and one vertex. Our aim is to investigate the limit of the respective operator family $-\Delta_{\Omega_\varepsilon}^D$ when $\varepsilon \rightarrow 0$. We will show that it approximates in a suitable sense an operator on \mathcal{G} , namely the Schrödinger operator on the line with a point interaction depending on γ and $\lambda(\varepsilon)$.

Before to state our main theorem, let us introduce some notation and mention some preliminary facts. Writing for brevity $\Omega' = \mathbb{R} \times (-d, d)$, we recall the following result [DE95, EŠ89]:

Proposition 1. *For any $0 < \varepsilon \leq \varepsilon_0$ let C_ε be as described above, with γ piecewise C^2 and compactly supported, such that γ', γ'' are bounded. Then $-\Delta_{\Omega_\varepsilon}^D$ is unitarily equivalent to the operator H_ε defined as the closure of the e.s.a. operator $H_{0\varepsilon}$ acting on $L^2(\Omega')$ as*

$$H_{0\varepsilon} = -\frac{\partial}{\partial s} \frac{1}{(1 + \varepsilon^{\alpha-1} u \sqrt{\lambda(\varepsilon)} \gamma(s/\varepsilon))^2} \frac{\partial}{\partial s} - \frac{1}{\varepsilon^{2\alpha}} \frac{\partial^2}{\partial u^2} + \frac{1}{\varepsilon^2} V_\varepsilon(s, u),$$

with the effective potential

$$V_\varepsilon(s, u) = -\frac{\lambda(\varepsilon) \gamma(s/\varepsilon)^2}{4(1 + \varepsilon^{\alpha-1} u \sqrt{\lambda(\varepsilon)} \gamma(s/\varepsilon))^2} + \frac{\varepsilon^{\alpha-1} u \sqrt{\lambda(\varepsilon)} \gamma''(s/\varepsilon)}{2(1 + \varepsilon^{\alpha-1} u \sqrt{\lambda(\varepsilon)} \gamma(s/\varepsilon))^3} - \frac{5}{4} \frac{\varepsilon^{2\alpha-2} u^2 \lambda(\varepsilon) \gamma'(s/\varepsilon)^2}{(1 + \varepsilon^{\alpha-1} u \sqrt{\lambda(\varepsilon)} \gamma(s/\varepsilon))^4}$$

and $\mathcal{D}(H_{0,\varepsilon}) = \{\psi \in L^2(\Omega') \mid \psi \in C^\infty(\Omega'), \psi(s, d) = \psi(s, -d) = 0, H_{0\varepsilon} \psi \in L^2(\Omega')\}$.

Let us next introduce the transversal modes, i.e., the normalized functions $\phi_n(u)$ which solve the equation $-\varepsilon^{-2\alpha} \phi_n''(u) = E_{\varepsilon,n} \phi_n(u)$ with the boundary conditions $\phi_n(\varepsilon^\alpha d) = \phi_n(-\varepsilon^\alpha d) = 0$. In particular, the corresponding eigenvalues $E_{\varepsilon,n}$ are explicitly given by

$$E_{\varepsilon,n} = \left(\frac{n\pi}{2d\varepsilon^\alpha}\right)^2 \quad \text{with } n = 1, 2, \dots$$

The resolvent of H_ε admits an integral representation with the kernel $(H_\varepsilon - z)^{-1}(s, u, s', u')$ for every $z \in \rho(H_\varepsilon)$ with $\text{Im} \sqrt{z} > 0$, where $\rho(H_\varepsilon)$ is the resolvent set of H_ε , cf theorem II.37 in [Si71]. Using it we define the projection of the resolvent on the normal modes eigenspaces as

$$\overline{R}_{n,m}^\varepsilon(k^2, s, s') := \int_{-d}^d du du' \phi_n(u)(H_\varepsilon - k^2 - E_{\varepsilon,m})^{-1}(s, u, s', u')\phi_m(u').$$

The operators $\overline{R}_{n,m}^\varepsilon(k^2) : L^2(\mathbb{R}) \rightarrow \text{Ran}[\overline{R}_{n,m}^\varepsilon(k^2)] \subset L^2(\mathbb{R})$ introduced in this way are bounded operator-valued analytic functions of k^2 for all $k^2 \in \mathbb{C} \setminus \mathbb{R}$ and $\text{Im} k > 0$.

Next we have to recall some facts about one-dimensional Schrödinger operators. We say that the Hamiltonian

$$\overline{H} = -\frac{d^2}{ds^2} + \overline{V}(s) \tag{2.3}$$

has a zero energy resonance if there exists a function $\psi_r \in L^\infty(\mathbb{R})$, $\psi_r \notin L^2(\mathbb{R})$, such that $\overline{H}\psi_r = 0$ holds in the sense of distributions. In particular, if

$$\int_{\mathbb{R}} \overline{V}(s) ds \neq 0 \quad \text{and} \quad e^{a|\cdot|} \overline{V} \in L^1(\mathbb{R}) \tag{2.4}$$

holds for some $a > 0$, then exactly one of the following situations can occur [BGW85]:

Case I. The Hamiltonian \overline{H} does not have a zero energy resonance.

Case II. The Hamiltonian \overline{H} has a zero energy resonance; in such a case the function ψ_r can be chosen real and two real constants can be defined,

$$\begin{aligned} c_1 &= \left[\int_{\mathbb{R}} \overline{V}(s) ds \right]^{-1} \int_{\mathbb{R}} \int_{\mathbb{R}} \overline{V}(s) \frac{|s-s'|}{2} \overline{V}(s') \psi_r(s') ds ds', \\ c_2 &= -\frac{1}{2} \int_{\mathbb{R}} s \overline{V}(s) \psi_r(s) ds, \end{aligned} \tag{2.5}$$

and moreover, c_1 and c_2 cannot vanish simultaneously. Let us stress that the constants c_1 and c_2 defined in (2.5) coincide with those employed in [ACF07].

Let us next introduce a pair of Hamiltonians on \mathcal{G} both acting as $f \mapsto -f''$ but differing by the boundary conditions in the vertex. The first is the Dirichlet-decoupled operator \overline{H}^d with the domain $\mathcal{D}(\overline{H}^d) := \{f \in H^2(\mathbb{R} \setminus \{0\}) \cap H^1(\mathbb{R}) \mid f(0) = 0\}$. The other is a point-interaction Hamiltonian \overline{H}^r , which again acts as $\overline{H}^r f = -f''$ but on the domain

$$\begin{aligned} \mathcal{D}(\overline{H}^r) &:= \left\{ f \in H^2(\mathbb{R} \setminus \{0\}) \mid (c_1 + c_2)f(0^+) = (c_1 - c_2)f(0^-), \right. \\ &\quad \left. (c_1 - c_2)f'(0^+) = (c_1 + c_2)f'(0^-) + \frac{\hat{\lambda}}{c_1 + c_2} f(0^-) \right\} \quad \text{for } c_2 \neq -c_1; \\ \mathcal{D}(\overline{H}^r) &:= \left\{ f \in H^2(\mathbb{R} \setminus \{0\}) \mid f(0^-) = 0, f'(0^+) = \frac{\hat{\lambda}}{4c_1^2} f(0^+) \right\} \quad \text{for } c_2 = -c_1, \end{aligned}$$

where we put

$$\hat{\lambda} := \lambda_1 \int_{\mathbb{R}} \overline{V}(s) (\psi_r(s))^2 ds. \tag{2.6}$$

The graph \mathcal{G} identifies naturally with a line and both \overline{H}^d and \overline{H}^r belong to the family of self-adjoint extensions of the symmetric operator $\overline{L}_0 f := -f''$ with the domain $\mathcal{D}(\overline{L}_0) := C_0^\infty(\mathbb{R} \setminus \{0\})$ [ABD95].

Let us say a few more words on the family \overline{H}^r which obviously depends on two real parameters. It is a straightforward exercise to check that the boundary conditions appearing in the definition of $\mathcal{D}(\overline{H}^r)$ can be rewritten in the form ((1.1)) with $\Psi := (f(0^+), f(0^-))^T$, $\Psi' := (f'(0^+), -f'(0^-))^T$ and the 2×2 unitary matrix

$$U := \frac{1}{2(c_1^2 + c_2^2) + i\hat{\lambda}} \begin{pmatrix} -4c_1c_2 - i\hat{\lambda} & 2(c_1^2 - c_2^2) \\ 2(c_1^2 - c_2^2) & 4c_1c_2 - i\hat{\lambda} \end{pmatrix}. \tag{2.7}$$

In particular, for $\lambda_1 = 0$ the boundary conditions define the ‘scale-invariant’ Hamiltonian obtained in [ACF07]. Applications of this point interaction were discussed recently in [HC06], and it is worth mentioning that it appears also in the theory of regular tree graphs [So04]. On the other hand, in distinction to [ACF07] we have here a wider class which contains, in particular, the standard δ interaction of coupling strength $\hat{\lambda}$ [AGHH05] corresponding to $c_1 = 1$ and $c_2 = 0$. Spectral and scattering properties of \overline{H}^r are well known [EG99] and we recall them only briefly:

Proposition 2. *For any $-\infty < \hat{\lambda} \leq \infty$, the essential spectrum of \overline{H}^r is absolutely continuous and coincides with the interval $[0, \infty)$. Furthermore, for $\hat{\lambda} > 0$ there are no eigenvalues, while for $\hat{\lambda} < 0$ there is just one negative eigenvalue given by $k^2 = k_0^2 = -\frac{1}{4}\hat{\lambda}^2(c_1^2 + c_2^2)^{-1}$ and the corresponding normalized eigenfunction is*

$$\psi_0 = \sqrt{\frac{|\hat{\lambda}|}{2} \frac{1}{c_1^2 + c_2^2}} \begin{cases} (c_1 - c_2) e^{ik_0s} & s > 0 \\ (c_1 + c_2) e^{-ik_0s} & s < 0 \end{cases}, \quad k_0 = \frac{i|\hat{\lambda}|}{2(c_1^2 + c_2^2)}, \quad \hat{\lambda} < 0.$$

Finally, for $\hat{\lambda} = 0$ the operator \overline{H}^r has a zero energy resonance. The on-shell scattering matrix at energy k^2 , $k \geq 0$, is given by $\mathcal{S}(k) = \begin{bmatrix} \mathcal{S}^l(k) & \mathcal{R}^r(k) \\ \mathcal{R}^l(k) & \mathcal{S}^r(k) \end{bmatrix}$ with the amplitudes

$$\mathcal{S}^{(l,r)}(k) = \frac{2k(c_1^2 - c_2^2)}{2k(c_1^2 + c_2^2) + i\hat{\lambda}}, \quad \mathcal{R}^{(l,r)}(k) = \pm \frac{4kc_1c_2 \mp i\hat{\lambda}}{2k(c_1^2 + c_2^2) + i\hat{\lambda}}.$$

Let now G_k be the resolvent of the free Laplacian on \mathbb{R} , it is a bounded operator-valued analytical function of k^2 for $k^2 \in \mathbb{C} \setminus \mathbb{R}^+$ and $\text{Im } k > 0$, with the integral kernel given by

$$G_k(s - s') = \frac{i}{2k} e^{ik|s-s'|} \quad k^2 \in \mathbb{C} \setminus \mathbb{R}^+, \quad \text{Im } k > 0.$$

By Krein’s formula [ABD95, EG99] the integral kernel of the resolvent $\overline{R}^d(k^2) := (\overline{H}^d - k^2)^{-1}$ is

$$\overline{R}^d(k^2, s, s') = G_k(s - s') + 2ikG_k(s)G_k(s'), \quad k^2 \in \mathbb{C} \setminus \mathbb{R}^+, \quad \text{Im } k > 0,$$

while the integral kernel of the resolvent $\overline{R}^r(k^2) := (\overline{H}^r - k^2)^{-1}$ equals

$$\begin{aligned} \overline{R}^r(k^2; s, s') &= G_k(s - s') + 2ik \frac{2kc_2^2 + i\hat{\lambda}}{2k(c_1^2 + c_2^2) + i\hat{\lambda}} G_k(s)G_k(s') + \frac{4ic_2^2}{2k(c_1^2 + c_2^2) + i\hat{\lambda}} G'_k(s)G'_k(s') \\ &+ \frac{4kc_1c_2}{2k(c_1^2 + c_2^2) + i\hat{\lambda}} [G_k(s)G'_k(s') + G'_k(s)G_k(s')], \quad k^2 \in \rho(\overline{H}^r), \quad \text{Im } k > 0. \end{aligned}$$

Our main result can be now stated as follows:

Theorem 2.1. *Suppose that for every $0 < \varepsilon \leq \varepsilon_0$ the curve C_ε has no self-intersections, γ is piecewise C^2 with a compact support and γ', γ'' are bounded. Assuming $\alpha > 5/2$, we have*

(i) if $-\frac{d^2}{ds^2} - \frac{1}{4}\gamma^2(s)$ does not have a zero energy resonance, then we have in the uniform (in other words, operator-norm) limit

$$u - \lim_{\varepsilon \rightarrow 0} \overline{R}_{n,m}^\varepsilon(k^2) = \delta_{n,m} \overline{R}^d(k^2) \quad k^2 \in \mathbb{C} \setminus \mathbb{R}, \quad \text{Im } k > 0;$$

(ii) if, on the other hand, $-\frac{d^2}{ds^2} - \frac{1}{4}\gamma^2(s)$ has a zero energy resonance, then

$$u - \lim_{\varepsilon \rightarrow 0} \overline{R}_{n,m}^\varepsilon(k^2) = \delta_{n,m} \overline{R}^r(k^2) \quad k^2 \in \mathbb{C} \setminus \mathbb{R}, \quad \text{Im } k > 0,$$

where the constants c_1, c_2 and $\hat{\lambda}$, defined in (2.5) and (2.6), are obtained by setting $\overline{V} = -\frac{1}{4}\gamma^2$ and $\delta_{n,m}$ indicates the Kronecker symbol, $\delta_{n,m} = 0$, if $n \neq m$ and $\delta_{n,n} = 1$.

3. The limit of short-range potentials in dimension one

The main ingredient in the proof of theorem 2.1 is the analysis of scaling properties of one-dimensional Hamiltonians. Specifically, we will find the limiting behaviour as $\varepsilon \rightarrow 0$ for

$$\overline{H}_\varepsilon := -\frac{d^2}{ds^2} + \frac{\lambda(\varepsilon)}{\varepsilon^2} \overline{V}\left(\frac{s}{\varepsilon}\right), \quad s \in \mathbb{R}.$$

Recall that this problem is well studied if the limit is considered, roughly speaking, around the free operator [AGHH05]. The case which involves threshold resonances is different and in a sense similar to approximations of point interactions in dimensions two and three mentioned above. It is useful to discuss this issue separately because in our opinion it is of independent interest as approximation of a class of point perturbations of the Laplacian in dimension one with scaled potentials. Let us stress that the δ' -type interactions do not belong to this class—a way to approximate them by regular potentials can be found in [ENZ01]. The main idea of our analysis comes from the work [BGW85].

In the following we suppose that the conditions (2.4) are satisfied. As $\lambda(\varepsilon)$ is real analytic near the origin by assumption we can make the expansion ((2.2)) for small ε more specific writing

$$\lambda(\varepsilon) = 1 + \sum_{n=1}^{\infty} \lambda_n \varepsilon^n. \tag{3.1}$$

For every $\varepsilon > 0$, the resolvent of \overline{H}_ε is a bounded operator-valued analytic function of k^2 as long as $k^2 \in \mathbb{C} \setminus \mathbb{R}^+$, $k^2 \notin \sigma_p(\overline{H}_\varepsilon)$ and $\text{Im } k > 0$, where $\sigma_p(\overline{H}_\varepsilon)$ denotes the point spectrum of \overline{H}_ε . As usual we factorize the interaction using the functions

$$v(s) := |\overline{V}(s)|^{1/2}, \quad u(s) := \text{sgn}[\overline{V}(s)] |\overline{V}(s)|^{1/2},$$

which allows us to write $(\overline{H}_\varepsilon - k^2)^{-1}$ as in [AGHH05], namely

$$(\overline{H}_\varepsilon - k^2)^{-1} = G_k - \frac{\lambda(\varepsilon)}{\varepsilon} A_\varepsilon(k) T_\varepsilon(k) C_\varepsilon(k), \tag{3.2}$$

where

$$T_\varepsilon(k) = [1 + \lambda(\varepsilon) u G_{\varepsilon k} v]^{-1} \quad \text{Im } k \geq 0, k \neq 0, k^2 \notin \sigma_p(\overline{H}_\varepsilon)$$

and $A_\varepsilon(k)$, $C_\varepsilon(k)$ are defined via their integral kernels, $A_\varepsilon(k; s, s') = G_k(s - \varepsilon s') v(s')$ and $C_\varepsilon(k; s, s') = u(s) G_k(\varepsilon s - s')$, respectively. We are interested in the behaviour of $T_\varepsilon(k)$ as $\varepsilon \rightarrow 0$. To this aim we define the operators P and Q by

$$P := \frac{1}{(v, u)} (v, \cdot) u, \quad Q := 1 - P$$

where (\cdot, \cdot) denotes the standard scalar product in $L^2(\mathbb{R})$; let us note that by assumption we have $(v, u) = \int_{\mathbb{R}} \bar{V}(s) ds \neq 0$. The operator $T_\varepsilon(k)$ can be written as in [BGW85],

$$T_\varepsilon(k) = \left[1 + \frac{i(v, u)}{2\varepsilon k} P + \tilde{M}_\varepsilon(k) \right]^{-1} \tag{3.3}$$

where $\tilde{M}_\varepsilon(k) \in \mathcal{B}(L^2, L^2)$, the Banach space of bounded operators from $L^2(\mathbb{R})$ to $L^2(\mathbb{R})$, for every $\varepsilon > 0$ and $\text{Im } k > 0$. Furthermore, if $e^{a|\cdot|} \bar{V} \in L^1(\mathbb{R})$ holds for some $a > 0$ then \tilde{M}_ε is analytic with respect to ε for $\varepsilon > -a/(2\text{Im } k)$ and the following series expansion converges in the $\mathcal{B}(L^2, L^2)$ -norm,

$$\tilde{M}_\varepsilon(k) = \sum_{n=0}^{\infty} \varepsilon^n \tilde{m}_n(k),$$

where

$$\tilde{m}_n(k) := (ik)^n m_n + \frac{i\lambda_{n+1}(v, u)}{2k} P + \sum_{j=0}^n \lambda_{n-j} (ik)^j m_j \quad n = 0, 1, 2, \dots$$

The operators m_n are Hilbert–Schmidt and do not depend on k , their integral kernels being given by the expressions

$$m_n(s, s') = -u(s) \frac{|s - s'|^{n+1}}{2(n+1)!} v(s').$$

The behaviour of $T_\varepsilon(k)$ as $\varepsilon \rightarrow 0$ depends strongly on the presence of a zero energy resonance for the Hamiltonian \bar{H} . Under the assumptions (2.4), the presence of such a resonance is equivalent to the existence of a function $\varphi_0 \in L^2(\mathbb{R})$ which satisfies the relation

$$\varphi_0 + Q M_0 Q \varphi_0 = 0. \tag{3.4}$$

Furthermore, if such a φ_0 exists, it can be chosen real, in which case the constants c_1, c_2 and $\hat{\lambda}$ defined in (2.5) and (2.6), respectively, are related to φ_0 by

$$c_1 = \frac{(v, m_0 \varphi_0)}{(v, u)}, \quad c_2 = \frac{1}{2} ((\cdot)v, \varphi_0), \quad \hat{\lambda} = \lambda_1(\tilde{\varphi}_0, \varphi_0).$$

with $\tilde{\varphi}_0(s) := \text{sgn}[\bar{V}(s)]\varphi_0$, and $u(s)\psi_r(s) = -\varphi_0(s)$ holds a.e., cf lemma 2.2 in [BGW85]. Let us introduce the operator

$$P_0 := \begin{cases} 0 & \text{in the case I} \\ \frac{(\tilde{\varphi}_0, \cdot)\varphi_0}{(\tilde{\varphi}_0, \varphi_0)} & \text{in the case II} \end{cases}$$

and the complementary projection $Q_0 := 1 - P_0$. From lemma 3.1 in [BGW85] we infer that for $\varepsilon \in \mathbb{C} \setminus \{0\}$ small enough the following norm convergent series expansion holds:

$$[1 + Q m_0 Q + \varepsilon]^{-1} = \frac{P_0}{\varepsilon} + \sum_{n=0}^{\infty} (-\varepsilon)^n T_{\text{red}}^{n+1},$$

where $T_{\text{red}} = u - \lim_{\varepsilon \rightarrow 0} [1 + Q m_0 Q + \varepsilon]^{-1} Q_0$ is the reduced resolvent. The following claim is a generalization of theorem 3.1 in [BGW85].

Lemma 1. *Suppose that \bar{V} satisfies the conditions (2.4) and take $\lambda(\varepsilon)$ real analytic near the origin and with the series expansion (3.1). Assume that $k^2 \notin \sigma_p(\bar{H}_\varepsilon)$, $\text{Im } k > 0$, and,*

additionally, that in the case II $k \neq -i\hat{\lambda}/(2(c_1^2+c_2^2))$. Then for all ε small enough the operator $T_\varepsilon(k)$ has the following norm-convergent series expansions:

$$T_\varepsilon(k) = \sum_{n=p}^{\infty} \varepsilon^n t_n(k), \tag{3.5}$$

where $p = 0$ in the case I and $p = 1$ in the case II. Moreover, we have

(i) in the case I

$$(v, t_0u) = 0; ((\cdot)v, t_0u) = (v, t_0(\cdot)u) = 0; \tag{3.6}$$

$$(v, t_1u) = -2ik; \tag{3.7}$$

(ii) in the case II

$$t_{-1}u = t_{-1}^*v = 0; ((\cdot)v, t_{-1}(\cdot)u) = -\frac{4ic_2^2}{2k(c_1^2+c_2^2)+i\hat{\lambda}}; \tag{3.8}$$

$$(v, t_0u) = 0; ((\cdot)v, t_0u) = (v, t_0(\cdot)u) = \frac{4kc_1c_2}{2k(c_1^2+c_2^2)+i\hat{\lambda}}; \tag{3.9}$$

$$(v, t_1u) = -2ik\frac{2kc_2^2+i\hat{\lambda}}{2k(c_1^2+c_2^2)+i\hat{\lambda}}. \tag{3.10}$$

Proof. We prove the lemma first in the case II. Let us assume that equation (3.4) is solved by $\varphi_0 \in L^2(\mathbb{R})$. By using the relation [BGW85]

$$1 + \frac{i(v, u)}{2\varepsilon k} P = \left[Q + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P \right]^{-1}$$

in formula (3.3) we obtain

$$T_\varepsilon = \left[1 + Q\tilde{M}_\varepsilon + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P\tilde{M}_\varepsilon \right]^{-1} \left[Q + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P \right]. \tag{3.11}$$

Since $Q\tilde{m}_0 = Qm_0$, the following norm convergent series expansion holds [BGW85]:

$$(1 + Q\tilde{m}_0 + \delta)^{-1} = \left[\frac{P_0}{\delta} + \sum_{n=0}^{\infty} (-\delta)^n T_{\text{red}}^{n+1} \right] \left[1 - \frac{Qm_0P}{1 + \delta} \right]. \tag{3.12}$$

Taking into account that $P_0 - P_0Qm_0P = -P_0m_0$ and performing a simple manipulation, we can set $\delta = -2i\varepsilon k/(v, u)$ and use relation (3.12) in formula (3.11) to obtain

$$T_\varepsilon = \left[1 + \left(\frac{(v, u)}{2i\varepsilon k} P_0m_0 + D_\varepsilon \right) \left(\frac{2i\varepsilon k}{(v, u)} + Q\tilde{M}_\varepsilon^{(1)} + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P\tilde{M}_\varepsilon \right) \right]^{-1} \\ \times \left[\frac{(v, u)}{2i\varepsilon k} P_0m_0 + D_\varepsilon \right] \left[Q + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P \right],$$

where

$$D_\varepsilon(k) := \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P_0Qm_0P + \sum_{n=0}^{\infty} \left(\frac{2i\varepsilon k}{(v, u)} \right)^n T_{\text{red}}^{n+1} \left[1 - \frac{Qm_0P}{1 - 2i\varepsilon k/(v, u)} \right]$$

and $\tilde{M}_\varepsilon^{(j)}(k) = \sum_{n=j}^\infty \varepsilon^n \tilde{m}_n(k)$ with $j = 1, 2, \dots$. After some computation we arrive at the following formula for the operator T_ε ,

$$T_\varepsilon = [1 + P_0 \tilde{B} + E_\varepsilon]^{-1} \left[\frac{(v, u)}{2i\varepsilon k} P_0 m_0 + D_\varepsilon \right] \left[Q + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P \right], \quad (3.13)$$

where

$$\tilde{B}(k) = m_0 + \frac{(v, u)}{2ik} m_0 Q \tilde{m}_1(k) - m_0 P \tilde{m}_0(k)$$

and

$$E_\varepsilon(k) = \frac{(v, u)}{2i\varepsilon k} P_0 m_0 Q \tilde{M}_\varepsilon^{(2)}(k) + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P_0 m_0 P \tilde{m}_0(k) - \frac{i(v, u)}{2\varepsilon k + i(v, u)} P_0 m_0 P \tilde{M}_\varepsilon^{(1)}(k) + D_\varepsilon(k) \left(\frac{2i\varepsilon k}{(v, u)} + Q \tilde{M}_\varepsilon^{(1)}(k) + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P \tilde{M}_\varepsilon(k) \right).$$

The operator $\tilde{B}(k)$ does not depend on ε while $D_\varepsilon(k)$ and $E_\varepsilon(k)$ have with respect to the parameter the following norm convergent series expansions:

$$D_\varepsilon(k) = \sum_{n=0}^\infty \varepsilon^n d_n(k), \quad E_\varepsilon(k) = \sum_{n=1}^\infty \varepsilon^n e_n(k). \quad (3.14)$$

Let us note that

$$\frac{(\tilde{\varphi}_0, \tilde{B}\varphi_0)}{(\tilde{\varphi}_0, \varphi_0)} = -1 - \frac{(v, u)}{(\tilde{\varphi}_0, \varphi_0)} (c_1^2 + c_2^2 + i\hat{\lambda}/(2k))$$

and $P_0 \tilde{B} P_0 = (\tilde{\varphi}_0, \tilde{B}\varphi_0)/(\tilde{\varphi}_0, \varphi_0) P_0$. In a similar way as in [BGW85] we can explicitly evaluate for $k \neq -i\hat{\lambda}/(2(c_1^2 + c_2^2))$ the inverse of $1 + P_0 \tilde{B}$ obtaining

$$[1 + P_0 \tilde{B}]^{-1} = 1 + \frac{(\tilde{\varphi}_0, \varphi_0)}{(v, u)} \frac{1}{c_1^2 + c_2^2 + i\hat{\lambda}/(2k)} P_0 \tilde{B}.$$

Formula (3.13) implies that in the case II the norm convergent series expansion (3.5) holds for ε small enough with $p = -1$.

Keeping only the corresponding terms of order ε^{-1} on the right-hand side of equation (3.13) and using the relation $P_0 m_0 Q = -P_0$ we obtain

$$t_{-1} = \frac{(v, u)}{2ik} [1 + P_0 \tilde{B}]^{-1} P_0 m_0 Q = \frac{(\tilde{\varphi}_0, \varphi_0)}{2ik} \frac{1}{(c_1^2 + c_2^2 + i\hat{\lambda}/(2k))} P_0.$$

Relations (3.8) follow from $P_0 u = P_0^* v = 0$ and $((\cdot)v, P_0(\cdot)u) = 4c_2^2/(\tilde{\varphi}_0, \varphi_0)$. Inspecting the terms of order zero in ε on the right-hand side of (3.13) we obtain

$$t_0 = [1 + P_0 \tilde{B}]^{-1} [-P_0 m_0 P + d_0 Q - e_1 t_{-1}].$$

The relation $(v, t_0 u) = 0$ is a consequence of the fact that $([1 + P_0 \tilde{B}]^{-1})^* v = v$. Moreover, by a direct calculation based on the relation $T_{\text{red}} P = P T_{\text{red}} = P$ similar to [BGW85] one can check that

$$((\cdot)v, t_0 u) = -((\cdot)v, [1 + P_0 \tilde{B}]^{-1} P_0 m_0 u)$$

and

$$(v, t_0(\cdot)u) = -(v, e_1 t_{-1}(\cdot)u)$$

from which relations (3.9) follow. Formula (3.10) is obtained by considering the terms of order ε on the right-hand side of equation (3.13) in combination with the relation

$$(v, t_1 u) = -\frac{2ik}{(v, u)} (v, d_0 u) - (v, e_1 [1 + P_0 \tilde{B}]^{-1} P_0 m_0 u).$$

It remains to deal with the case I. Then $P_0 = 0$ and equation (3.13) becomes

$$T_\varepsilon = [1 + E_\varepsilon]^{-1} D_\varepsilon \left[Q + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P \right], \quad (3.15)$$

where

$$D_\varepsilon(k) = \sum_{n=0}^{\infty} \left(\frac{2i\varepsilon k}{(v, u)} \right)^n T_{\text{red}}^{n+1} \left[1 - \frac{Qm_0 P}{1 - 2i\varepsilon k/(v, u)} \right]$$

and

$$E_\varepsilon(k) = D_\varepsilon(k) \left(\frac{2i\varepsilon k}{(v, u)} + Q\tilde{M}_\varepsilon^{(1)}(k) + \frac{2\varepsilon k}{2\varepsilon k + i(v, u)} P\tilde{M}_\varepsilon(k) \right).$$

The series expansions (3.14) still hold, and the norm convergent series expansion (3.5) in the case I is valid with $p = 0$. Let us note that $[1 + E_\varepsilon]$ is invertible for $\varepsilon \geq 0$ with ε small enough and, consequently, it is not necessary to assume $k \neq -i\hat{\lambda}/(2(c_1^2 + c_2^2))$. In the case I we thus have

$$t_0 = d_0 Q = T_{\text{red}}[1 - Qm_0 P]Q,$$

from which it easily follows that $(v, t_0 u) = 0$, and from $PT_{\text{red}} = T_{\text{red}}P = P$ one gets relations (3.6). The terms of order ε on the right-hand side of equation (3.15) give

$$t_1 = \frac{2k}{i(v, u)} d_0 P + d_1 Q - e_1 d_0 Q,$$

formula (3.7) then follows from $(v, d_0 u) = (v, u)$. \square

With this result at hand we can follow the argument line of [ACF07] to establish the norm-resolvent convergence of the Hamiltonian \overline{H}_ε to \overline{H}^d or \overline{H}^r , depending on the potential \overline{V} ; we omit the details. Using formulae (3.6)–(3.10) in the proof of lemma 1 of [ACF07] we arrive at the following conclusion:

Theorem 3.1. *Suppose that \overline{V} satisfies the conditions (2.4) and $\lambda(\varepsilon)$ is real analytic near the origin having the series expansion (3.1). Then we have*

(i) *in the case I*

$$u - \lim_{\varepsilon \rightarrow 0} (\overline{H}_\varepsilon - k^2)^{-1} = \overline{R}^d(k^2) \quad k^2 \in \mathbb{C} \setminus \mathbb{R}, \quad \text{Im } k > 0;$$

(ii) *in the case II*

$$u - \lim_{\varepsilon \rightarrow 0} (\overline{H}_\varepsilon - k^2)^{-1} = \overline{R}^r(k^2) \quad k^2 \in \mathbb{C} \setminus \mathbb{R}, \quad \text{Im } k > 0.$$

4. Proof of theorem 2.1

Also the rest of the proof of the main result now follows closely [ACF07] so it is sufficient to sketch the argument. It splits into two steps. The first was dealt with in the previous section, the second step consists of the proof of the claim given below. Since the latter is essentially as lemma 3 in [ACF07], we just state it omitting the details.

Lemma 4.1. *Suppose that C_ε has no self-intersections for every $0 < \varepsilon \leq \varepsilon_0$ and, moreover, that γ is piecewise C^2 , has compact support and γ', γ'' are bounded. Fix an $\alpha > 5/2$ and define H_ε^γ as the closure of the e.s.a. operator*

$$H_{0\varepsilon}^\gamma := -\frac{\partial^2}{\partial s^2} - \frac{1}{\varepsilon^{2\alpha}} \frac{\partial^2}{\partial u^2} - \frac{\lambda(\varepsilon)}{\varepsilon^2} \frac{\gamma(s/\varepsilon)^2}{4}$$

with the domain

$$\mathcal{D}(H_{0\varepsilon}^\gamma) := \{\psi \in L^2(\Omega') \mid \psi \in C^\infty(\Omega'), \psi(s, d) = \psi(s, -d) = 0, H_{0\varepsilon}^\gamma \psi \in L^2(\Omega')\}.$$

Defining the matrix elements $R_{n,m}^{\gamma,\varepsilon}$ with respect to the transverse modes ϕ_n and ϕ_m ,

$$R_{n,m}^{\gamma,\varepsilon}(k^2; s, s') = \int_{-d}^d du du' \phi_n(u) (H_\varepsilon^\gamma - k^2 - E_{\varepsilon,m})^{-1}(s, u, s', u') \phi_m(u'),$$

we have

$$\mathbf{u} - \lim_{\varepsilon \rightarrow 0} (R_{n,m}^\varepsilon(k^2) - R_{n,m}^{\gamma,\varepsilon}(k^2)) = 0 \quad k^2 \in \mathbb{C} \setminus \mathbb{R}, \quad \text{Im } k > 0.$$

In analogy with [ACF07], theorem 2.1 is now obtained by combination of theorem 3.1 and lemma 4.1.

Acknowledgments

The research was partially supported by GAAS and MEYS of the Czech Republic under projects A100480501 and LC06002, by the Collaborative Research Center (SFB) 611 ‘Singular Phenomena and Scaling in Mathematical Models’ and by the Deutscher Akademischer Austauschdienst (DAAD).

Note added in proof. At the same time as this paper was posted as a preprint, an extension of [MV06] was posted, 0704.2795v1 [Math-ph], showing that the ‘Kirchhoff’ conditions can be obtained in the case of a threshold resonance.

References

- [ABD95] Albeverio S, Brzeźniak Z and Dbrowski L 1995 Fundamental solutions of the heat and Schrödinger equations with point interactions *J. Funct. Anal.* **130** 220–54
- [ACF07] Albeverio S, Cacciapuoti C and Finco D 2007 Coupling in the singular limit of thin quantum waveguides *J. Math. Phys.* **48** 032103
- [AGHH05] Albeverio S, Gesztesy F, Høegh-Krohn R and Holden H 2005 *Solvable Models in Quantum Mechanics* 2nd edn (New York: Chelsea) (with an appendix by P Exner)
- [BCFK06] Berkolaiko G, Carlson R, Fulling S and Kuchment P (ed) 2006 *Quantum Graphs and Their Applications (Contemporary Math. vol 415)* (Providence, RI: American Mathematical Society)
- [BGW85] Bollé D, Gesztesy F and Wilk S F J 1985 A complete treatment of low energy scattering in one dimension *J. Operator Theory* **13** 3–32
- [CE04] Cheon T and Exner P 2004 An approximation to δ' couplings on graphs *J. Phys. A: Math. Gen.* **37** L329–35
- [DT06] Dell’Antonio G and Tenuta L 2006 Quantum graphs as holonomic constraints *J. Math. Phys.* **47** 072102
- [DE95] Duclos P and Exner P 1995 Curvature-induced bound states in quantum waveguides in two and three dimensions *Rev. Math. Phys.* **7** 73–102
- [EEK05] Ekholm T, Kovařík H and Krejčířík D 2005 A Hardy inequality in twisted waveguides *Preprint math-ph/0512050 (Arch. Rat. Mech. Anal. to appear)*
- [E96] Exner P 1996 Weakly coupled states on branching graphs *Lett. Math. Phys.* **38** 313–20
- [EG99] Exner P and Grosse H 1999 Some properties of the one-dimensional generalized point interactions (a torso) *Preprint math-ph/9910029*
- [ENZ01] Exner P, Neidhardt H and Zagrebnov V A 2001 Potential approximations to δ' : an inverse Klaunder phenomenon with norm-resolvent convergence *Commun. Math. Phys.* **224** 593–612
- [EP05] Exner P and Post O 2005 Convergence of spectra of graph-like thin manifolds *J. Geom. Phys.* **54** 77–115
- [EP07] Exner P and Post O 2007 Convergence of resonances on thin branched quantum wave guides *Preprint math-ph/0702075*
- [EŠ89] Exner P and Šeba P 1989 Bound states in curved quantum waveguides *J. Math. Phys.* **30** 2574–80
- [ET07] Exner P and Turek O 2007 Approximations of singular vertex couplings in quantum graphs *Preprint math-ph/0703051*

- [FW93] Freidlin M and Wentzell A 1993 Diffusion processes on graphs and the averaging principle *Ann. Prob.* **21** 2215–45
- [Gr07] Grieser D 2007 Spectral asymptotics of the Dirichlet Laplacian on fat graphs A Talk at *INIAGA Workshop (April 2007)* (in preparation)
- [Ha00] Harmer M 2000 Hermitian symplectic geometry and extension theory *J. Phys. A: Math. Gen.* **33** 9193–203
- [HC06] Hejčík P and Cheon T 2006 Irregular dynamics in a solvable one-dimensional quantum graph *Phys. Lett. A* **356** 290–3
- [KS00] Kostykin V and Schrader R 2000 Kirchhoff's rule for quantum wires: II. The inverse problem with possible applications to quantum computers *Fortschr. Phys.* **48** 703–16
- [KZ01] Kuchment P and Zeng H 2001 Convergence of spectra of mesoscopic systems collapsing onto a graph *J. Math. Anal. Appl.* **258** 671–700
- [MV06] Molchanov S and Vainberg B 2006 Scattering solutions in a network of thin fibers: small diameter asymptotics *Preprint* [math-ph/0609021](https://arxiv.org/abs/math-ph/0609021)
- [Po05a] Post O 2006 Spectral convergence of quasi-one-dimensional spaces *Ann. Henri Poincaré* **7** 933–73
- [Po05b] Post O 2005 Branched quantum wave guides with Dirichlet boundary conditions: the decoupling case *J. Phys. A: Math. Gen.* **38** 4917–31
- [RS01] Rubinstein J and Schatzman M 2001 Variational problems on multiply connected thin strips: I. Basic estimates and convergence of the Laplacian spectrum *Arch. Ration. Mech. Anal.* **160** 271–308
- [Sa01] Saitō T 2001 Convergence of the Neumann Laplacian on shrinking domains *Analysis* **21** 171–204
- [Si71] Simon B 1971 *Quantum Mechanics for Hamiltonians Defined as Quadratic Forms* (Princeton: Princeton University Press)
- [So04] Solomyak M 2004 On the spectrum of the Laplacian on regular metric trees *Waves Random Media* **14** S155–71