

Comment on the sign of the Casimir force

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2007 J. Phys. A: Math. Theor. 40 9089

(<http://iopscience.iop.org/1751-8121/40/30/028>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.144

The article was downloaded on 03/06/2010 at 06:06

Please note that [terms and conditions apply](#).

Comment on the sign of the Casimir force

C P Bachas

Laboratoire de Physique Théorique de l'Ecole Normale Supérieure¹, 24 rue Lhomond,
75231 Paris cedex, France

Received 3 April 2007, in final form 7 June 2007

Published 12 July 2007

Online at stacks.iop.org/JPhysA/40/9089

Abstract

I show that reflection positivity implies that the force between any mirror pair of charge-conjugate probes of the quantum vacuum is attractive. This generalizes a recent theorem of Kenneth and Klich (2006 *Phys. Rev. Lett.* **97** 160401 (*Preprint* quant-ph/0601011)) to interacting quantum fields, to arbitrary semiclassical bodies and to quantized probes with non-overlapping wavefunctions. I also prove that the torques on charge-conjugate probes tend always to rotate them into a mirror-symmetric position.

PACS numbers: 03.70.+k, 12.20.–m

The Casimir force [1] and related electromagnetic interactions [2, 3] are the dominant forces between neutral non-magnetic objects at scales ranging from a few nanometres to hundreds of microns. These forces play, therefore, an important role both in mesoscopic gravity experiments [4], and in the design and fabrication of micro- and nano-electromechanical systems [5, 6]. Of particular interest is the possibility of repulsive interactions, since a frequent cause of malfunction of miniature devices is the collapse and subsequent permanent adhesion of mechanical elements, known as ‘stiction’ [7, 8]. It is therefore important to understand under what conditions such residual electromagnetic forces may change sign.

Motivated by this question, Kenneth and Klich have recently shown that a mirror pair of dielectric bodies always attract [9]. Their proof is based on a detailed study of the determinant formula that enters in the leading-order semiclassical calculation of the force. Here I want to point out that this result follows from a general property of quantum field theory known as ‘reflection positivity’ [10, 11]. This extends the validity of the theorem to interacting quantum fields, arbitrary static vacuum probes, and (with assumptions to be discussed) beyond the semiclassical approximation of the probes, as well as in the presence of spectator bodies or material media. The proof applies equally well to electrostatic and magnetostatic forces, as to those arising from the quantum and thermal fluctuations of the electromagnetic field. A variant of the argument shows that torques always tend to rotate a pair of conjugate probes into a mirror-symmetric configuration. The arguments presented here have been used in the

¹ Unité mixte de recherche (UMR 8549) du CNRS et de l'ENS, associée à l'Université Pierre et Marie Curie et aux fédérations de recherche FR684 et FR2687.

past to establish the concavity of the quark/anti-quark potential [12, 13]. The main point of the present paper is to explain that they apply to a much broader class of physical situations.

Reflection positivity is a general property of Euclidean quantum field theory which guarantees the existence of a positive Hilbert space and a self-adjoint, non-negative Hamiltonian [10]. It can be stated as the following correlation inequality:

$$\langle f \Theta(f) \rangle \geq 0, \quad (1)$$

where Θ denotes the reflection with respect to a three-dimensional hyperplane in \mathbb{R}^4 , f is any functional of the fields with arguments to the left of the hyperplane, and the action of Θ on f is anti-unitary (i.e. $\Theta(cf) = c^* \Theta(f)$). Inequality (1) can be established easily in the path-integral formulation of quantum field theory. Consider, for example, a real scalar field defined on a hypercubic lattice Λ which admits Θ as a reflection symmetry. Let the functional-integral measure be

$$d\mu = \left(\prod_x d\phi_x \right) e^{-S_\phi} \quad \text{with} \quad S_\phi = \sum_{\langle x, y \rangle} \frac{1}{2} (\phi_x - \phi_y)^2 + \sum_x V(\phi_x), \quad (2)$$

where V is the scalar potential and $\langle x, y \rangle$ the lattice links. We also define Λ_+ , Λ_0 and Λ_- to be the collections of sites and links that lie to the left, on, or to the right of the mirror hyperplane. Splitting the integration measure accordingly, $d\mu = d\mu_+ d\mu_0 d\mu_-$, one finds

$$\langle f(\phi_x) f^*(\phi_{\Theta x}) \rangle = \int d\mu_0 \left| \int d\mu_+ f(\phi_x) \right|^2 \quad (3)$$

for any functional f that is defined entirely in Λ_+ . Inequality (1), with $\Theta(f(\phi_x)) \equiv f^*(\phi_{\Theta x})$, follows now from the positivity of the measure $d\mu_0$. This proof only applies, strictly speaking, to lattice symmetries, but it extends to all mirror reflections in the continuum limit where rotational invariance is restored.

Reflection positivity for gauge fields can be established in a similar way [11], starting, for instance, from the Wilson action

$$S_{\text{Wilson}} = -\frac{1}{2g^2} \sum_P \text{Re tr} \left(\prod_{\langle x, y \rangle \in P} U_{\langle x, y \rangle} \right). \quad (4)$$

Here P labels the lattice plaquettes, the product runs over the links of P in a path-ordered way, and the link variables $U_{\langle x, y \rangle} = U_{\langle y, x \rangle}^\dagger$ take values in the gauge group [14]. We define the action of Θ as follows, $\Theta(f(U_{\langle x, y \rangle})) \equiv f^*(U_{\langle \Theta x, \Theta y \rangle})$, and note that it leaves invariant the Wilson action. Inequality (1) follows now easily from positivity of the Haar measure, $\prod dU_{\langle x, y \rangle}$, by a similar argument as above. The extension to fermion fields is also straightforward, provided one defines the action of Θ appropriately [15, 11]. Thus (1) is a general property of relativistic quantum field theories, and in particular of quantum electrodynamics and of the standard model.

It is important here to emphasize that the existence of a reflection operator Θ satisfying (1) is only a consequence of unitarity, and makes no assumptions about the discrete symmetries P , C and T .² This is made clear by the following facts:

- A bounded-from-below potential for charged scalar fields respects reflection positivity, even though it can lead to explicit or spontaneous breaking of the charge-conjugation symmetry C . Note that, unlike Θ , the operation C does not complex-conjugate the constant parameters of the scalar potential.

² The issue was raised by Adam Schwimmer. I am grateful to him, as well as to Henri Epstein and John Iliopoulos, for helping to clarify it.

- A θ -term respects reflection positivity, but breaks explicitly parity and time reversal. Note that $\theta \int \epsilon_{\mu\nu\rho\sigma} \text{tr} F^{\mu\nu} F^{\rho\sigma}$ is odd under P or T , but it makes an imaginary contribution to the Euclidean action. Thus the combination of reflection and complex conjugation leaves invariant the functional-integral measure, and inequality (1) continues to hold in the presence of the θ -term³.

In some of the situations discussed below, Θ will act in the same way as the combination CP . This is however only a coincidence, as the above examples clearly illustrate.

There is a second point that needs clarification: positivity of the theory requires, strictly speaking, inequality (1) only for reflections in the direction of (Euclidean) time. These are indistinguishable from space reflections if we consider probes of a relativistic vacuum. In the presence of a medium, on the other hand, the measure need not be invariant under an operation Θ that reflects space. Only when it is will we be allowed to extend the validity of our theorem.

Let us now concentrate on functionals f which, when inserted in the path integral, describe static semiclassical probes interacting with the electromagnetic vacuum. Examples are a superconducting shell, a dielectric body, or (a collection of) infinitely heavy electric charges. The corresponding functionals read.

Probe	Conductor	Dielectric	Point charges
f	$\prod_{P \in \mathcal{W}} \delta(U_P - 1)$	$\prod_{P_0 \in \mathcal{W}} \exp(\alpha \text{Re } U_{P_0})$	$\prod_j (U_j)^{q_j}$

In these formulae U_P are the plaquette variables, \mathcal{W} is the world-volume of the dielectric body or the conducting shell, P_0 labels timelike plaquettes (i.e. those with one link in the time direction) and U_j are Wilson lines along the worldline of the j th particle whose electric charge is q_j . Note that $U_P \simeq \exp(iF_P)$, where F_P is the electromagnetic field strength along the plaquette. Thus the δ -functions in the first entry impose the vanishing of the electric and magnetic fields tangent, respectively normal, to the conducting shell, as appropriate. Note also that the material of the dielectric body has been assumed isotropic and non-dispersive, with relative permittivity ϵ given by $\alpha = (1 - \epsilon)/2g^2$. We will comment on dispersive media later on. Note finally that the use of lattice variables is not essential. It shows, however, that the relevant inequalities survive the (non-perturbative) regularization of the theory. One can still describe geometrically smooth probes by using linear combinations of plaquette field strengths in the definition of the above functionals.

For static probes the functionals f are time-translation invariant. We may thus impose periodic boundary conditions in the time direction ($x^0 = x^0 + \beta$) so that

$$\langle f_1 \dots f_N \rangle = e^{-\beta E(1, \dots, N)} \tag{5}$$

where β is the inverse temperature, and E is the free energy of the system in the presence of the probes $1, \dots, N$. The existence of a continuum limit for the correlator (5) raises, in fact, some subtle mathematical questions. Careful subtraction procedures have been proposed for Wilson-loop operators [17] and, at the one loop-level, for perfectly conducting shells [18]. A weak assumption, verified explicitly in the above two examples, is that (5) can be made finite by a multiplicative renormalization of the functionals, $f_a^{\text{ren}} = \exp(-\beta E_a) f_a$, where E_a is the (generally divergent) self-energy of the a th probe. This will be sufficient for our purposes here, since we will focus on the dependence of the energy on the relative position and orientation of the probes, rather than on their detailed composition and shape.

³ Defining the θ -term on the lattice so as to make the argument rigorous is tricky [16].




Figure 1. Two identical probes with centres of mass aligned along the Oz axis, and their mirror images in the (xz) plane. Probe 2 is in general rotated with respect to probe 1. Inequalities (7) and (10) are obtained in the special limits of zero rotation, or for $z_1 = z_2$.

We come now to the main point of this paper. Reflection positivity implies that $(f_1, f_2) \equiv \langle f_1 \Theta(f_2) \rangle$ is a non-negative inner product on the space of functionals that are defined on one side of a reflection hyperplane. This implies, in turn, the Schwarz inequality,

$$|\langle f_1 \Theta(f_2) \rangle|^2 \leq \langle f_1 \Theta(f_1) \rangle \langle f_2 \Theta(f_2) \rangle, \quad (6)$$

for any two probes 1 and 2 that live on the same side of the hyperplane. Consider the special case of identical probes, with the same orientation and with centres of mass aligned on the Oz axis. Let Θ act by reflecting the coordinate z (see figure 1). Combining inequality (6) with equation (5) we find

$$2E_{\text{mir}}(z_1 + z_2) \geq E_{\text{mir}}(2z_1) + E_{\text{mir}}(2z_2), \quad (7)$$

where $E_{\text{mir}}(z)$ is the interaction energy of the mirror pair of conjugate probes, and z is the separation of their centres. It follows from (7) that $E_{\text{mir}}(z)$ is a concave function, i.e., its second derivative is nowhere positive. Thus dE_{mir}/dz is monotone non-increasing. Since, furthermore, the energy is bounded at infinite separation from below, we conclude that dE_{mir}/dz is everywhere non-negative, so that the force is either attractive or exactly zero. This generalizes the proof of [9] to arbitrary static external probes, and beyond the one-loop approximation⁴. An extra immediate corollary is that the binding energy of a pair of conjugate probes can never grow faster than linearly with distance [12].

Note that it is crucial for the theorem's validity that the operation Θ involves a complex conjugation. This does not affect neutral probes, such as the conducting shell or the dielectric bodies, but it does flip the sign of electrostatic forces. To illustrate the point, consider the classical interaction energy between a pair of electric dipoles:

$$E(\mathbf{d}_1, \mathbf{d}_2 | \mathbf{r}) = \frac{\mathbf{d}_1 \cdot \mathbf{d}_2 - 3(\mathbf{d}_1 \cdot \hat{\mathbf{r}})(\mathbf{d}_2 \cdot \hat{\mathbf{r}})}{|\mathbf{r}|^3}. \quad (8)$$

⁴ The fact that a spherical perfectly conducting thin shell feels an outward Casimir pressure [19] does not contradict the theorem. It has indeed been recognized that separating an elastic shell into two rigid hemispheres is a mathematically singular operation, which introduces divergent edge contributions to the energy [20, 18].

The action of Θ on an electric dipole is $\Theta(\mathbf{d}) = -\mathbf{d} + 2\hat{\mathbf{n}}(\mathbf{d} \cdot \hat{\mathbf{n}})$, where $\hat{\mathbf{n}}$ is a unit normal to the reflection plane, and the overall minus sign comes from charge conjugation. Setting $\hat{\mathbf{n}} = \hat{\mathbf{r}}$ gives the interaction energy between a mirror pair of conjugate dipoles

$$E(\mathbf{d}, \Theta(\mathbf{d})|\mathbf{r}) = -\frac{\mathbf{d} \cdot \mathbf{d} + (\mathbf{d} \cdot \hat{\mathbf{r}})^2}{|\mathbf{r}|^3}. \tag{9}$$

The force between the pair is thus attractive, consistently with reflection positivity. Note that for a magnetic dipole created by a persistent current $\Theta(\mathbf{m}) = \mathbf{m} - 2\hat{\mathbf{n}}(\mathbf{m} \cdot \hat{\mathbf{n}})$. The theorem nevertheless continues to hold, because the interaction energy of two such magnetic dipoles is minus the expression in (8); see, for example, [21].

As a further application of inequality (6) we can consider the dependence of the energy on the spatial orientation of the probes. Let 1 and 2 be obtained from some reference probe by rotations \mathcal{R}_1 and \mathcal{R}_2 around its centre of mass (see figure 1). Inequality (6) then reads in a self-explanatory notation

$$2E(\mathcal{R}_1, \Theta(\mathcal{R}_2)) \geq E(\mathcal{R}_1, \Theta(\mathcal{R}_1)) + E(\mathcal{R}_2, \Theta(\mathcal{R}_2)). \tag{10}$$

It follows that $E(\mathcal{R}_1, \Theta(\mathcal{R}_2))$ is greater than the smaller of the two terms on the right-hand side, so that the minimum of the energy is obtained when the relative orientation of conjugate probes corresponds precisely to that of a mirror pair⁵. Since in this configuration the force is, as we saw, attractive, such systems will always tend to collapse unless stabilized by external torques and forces.

Let us now turn attention to situations in which a material medium, or a spectator body, break Lorentz invariance even before the insertion of external probes. Spatial and Euclidean-time reflections are no more equivalent, so that the theorem need not always hold. Its (partial) validity requires that the functional measure be invariant under (at least a subset of) *spatial* reflections and complex conjugation. We have already used this fact above, in our discussion of the thermal vacuum. The following examples further illustrate the point:

- Two identical pistons inside an infinite conducting cylinder will attract, in agreement with the conclusion of [22]. This follows from Θ -invariance for all reflections that are geometric symmetries of the cylinder. For a *finite* cylinder the energy is not only a function of the distance between the pistons, and the arguments of this paper do not apply. More generally, for a spectator body of finite extent, our argument can at best be used to fix the sign of torques.
- The proof continues to apply in the presence of a classical homogeneous and isotropic dispersive medium, as discussed already by Kenneth and Klich [9]. The Euclidean Maxwell action in this case can be written as

$$S_{\text{Max}} = \int d^3\mathbf{x} \int_0^\infty \frac{d\omega}{4\pi} (\epsilon(i\omega)|\mathbf{E}(\omega, \mathbf{x})|^2 + |\mathbf{B}(\omega, \mathbf{x})|^2), \tag{11}$$

where $\epsilon(\omega)$ is the dielectric permittivity, which must be analytic in the upper-half complex plane, and real and positive on the imaginary ω axis [23]. Although this action is non-local in Euclidean time, it does satisfy reflection positivity for space reflections, so that a mirror pair of conjugate probes still attracts⁶.

- A case where the theorem does not hold, consistently with claims in the literature [24], is that of mirror probes in a Fermi sea. Indeed a non-zero chemical-potential term μj^0 , where j^0 is the relevant number density, is invariant under charge conjugation and

⁵ The absolute preferred orientation depends on the details of the probes, and cannot be determined by such general arguments.

⁶ Note that the dielectric or superconducting probes discussed so far can be thought of as special limits of a spatially varying permittivity $\epsilon(\omega, \mathbf{x})$.

time reflection, but not under charge conjugation and *space* reflection. This seems to contradict our first example, since a conducting material could be after all modelled by a free-electron gas. The point is, however, the following: the theorem holds to the extent that the conductor can be described by a classical permittivity for the photon field, but does not apply if the electron-gas fluctuations become important⁷.

This last point can be made more clear by considering the extension of the theorem to quantum probes. A simple example is a charged non-relativistic quantum particle trapped inside a potential well. The corresponding functional of the electromagnetic field (that replaces the Wilson line of a classical charge) reads

$$f(A^\mu) = \int [D\mathbf{x}(t)] e^{-I} e^{iq \int (A^0 dt - \mathbf{A} \cdot d\mathbf{x})}. \quad (12)$$

Here $[D\mathbf{x}(t)]$ is the integration measure over particle trajectories,

$$I = \int_0^\beta dt \left[\frac{m}{2} \left(\frac{d\mathbf{x}}{dt} \right)^2 + V(\mathbf{x}) \right] \quad (13)$$

is the Euclidean non-relativistic particle action, and $V(\mathbf{x})$ the localizing potential. Plugging the functionals (12) in equation (5) gives the free energy of the combined system of particles and fields at inverse temperature β . If we could use inequality (6), we would then conclude as before that the force between a pair of conjugate probes is attractive. The problem with this argument is that the quantized particle does not, strictly speaking, live in one side of the reflecting plane. Forcing it to do so (e.g. by adding to V an artificial hard-wall term) will, however, introduce only a tiny error if the particle–mirror separation is much bigger than the localization length. For a harmonic-well potential the error will, for instance, decay exponentially fast. For sufficiently large probe separations the conclusions of the theorem should thus still apply.

We can easily extend the above argument to collections of particles, such as the protons, neutrons and electrons of a heavy atom. Note, however, that because Θ acts as charge conjugation, the mirror partner must be made of antimatter! Thus this last extension of the theorem to truly microscopic quantum probes is of little relevance for applications to realistic electromechanical systems.

Acknowledgments

I thank Bertrand Duplantier and Giuseppe Policastro for useful conversations. I also thank Israel Klich, Erhard Seiler, Adam Schwimmer and one of the PRL referees for comments that significantly improved the original manuscript. This research was partially supported by the European Superstring Theory Network MRTN-CT-2004-512194.

References

- [1] Casimir H B G 1948 On the attraction between two perfectly conducting plates *Proc. K. Ned. Akad. Wet.* **51** 793
- [2] Casimir H B G and Polder D 1948 The influence of retardation on the London-van der Waals forces *Phys. Rev.* **73** 360
- Lifshitz E M 1956 *Sov. Phys.—JETP* **2** 73
- Dzyaloshinskii I E, Lifshitz E M and Pitaevskii L P 1961 The general theory of van der Waals forces *Adv. Phys.* **10** 165

⁷ Other examples that violate the assumptions of our theorem are periodic boundary conditions for fermions [15], or the situations discussed in [25].

- [3] Some recent (and less recent) reviews are Levin F S and Micha D A (ed) 1993 *Long-range Casimir Forces: Theory and Recent Experiments on Atomic Systems* (New York: Plenum)
Milonni P 1994 *The Quantum Vacuum. An Introduction to Quantum Electrodynamics* (New York: Academic)
Jaekel M-T and Reynaud S 1997 Movement and fluctuations of the vacuum *Rep. Prog. Phys.* **60** 863
Bordag M, Mohideen U and Mostepanenko V M 2001 New developments in the Casimir effect *Phys. Rep.* **353** 1 (Preprint [quant-ph/0106045](#))
Duplantier B and Rivasseau V (ed) 2003 *Vacuum Energy: Renormalization (Poincaré Seminar 2002)* (Berlin: Birkhäuser)
- [4] For recent experiments, see Long J C, Chan H W, Churnside A B, Gulbis E A, Varney M C M and Price J C 2002 New experimental limits on macroscopic forces below 100-microns Preprint [hep-ph/0210004](#)
Hoyle C D, Kapner D J, Heckel B R, Adelberger E G, Gundlach J H, Schmidt U and Swanson H E 2004 Sub-millimeter tests of the gravitational inverse-square law *Phys. Rev. D* **70** 042004 (Preprint [hep-ph/0405262](#))
Decca R S, Lopez D, Chan H B, Fischbach E, Krause D E and Jamell C R 2005 Constraining new forces in the Casimir regime using the isoelectronic technique *Phys. Rev. Lett.* **94** 240401 (Preprint [hep-ph/0502025](#))
Smullin S J, Geraci A A, Weld D M, Chiaverini J, Holmes S P and Kapitulnik A 2005 New constraints on Yukawa-type deviations from Newtonian gravity at 20-microns *Phys. Rev. D* **72** 122001
Smullin S J, Geraci A A, Weld D M, Chiaverini J, Holmes S P and Kapitulnik A 2005 *Phys. Rev. D* **72** 129901 (Preprint [hep-ph/0508204](#)) (erratum)
- [5] Srivastava Y, Widom A and Friedman M H 1985 Microchips as precision quantum electrodynamic probes *Phys. Rev. Lett.* **55** 2246
- [6] Chan H B, Aksyuk V A, Kleiman R N, Bishop D J and Capasso F 2001 Quantum mechanical actuation of microelectromechanical systems by the Casimir force *Science* **291** 1941
Also Chan H B, Aksyuk V A, Kleiman R N, Bishop D J and Capasso F 2001 Nonlinear micromechanical Casimir oscillator *Phys. Rev. Lett.* **87** 211801 (Preprint [quant-ph/0109046](#))
- [7] Serry M, Walliser D and Maclay J 1998 The role of the Casimir effect in the static deflection and stiction of membrane strips in microelectromechanical systems (MEMS) *J. Appl. Phys.* **84** 2501
Serry M, Walliser D and Maclay J 1995 The anharmonic Casimir oscillator *J. Microelectromech. Syst.* **4** 193
- [8] Buks E and Roukes M L 2001 Stiction, adhesion energy, and the Casimir effect in micromechanical systems *Phys. Rev. B* **63** 033402 (Preprint [cond-mat/0008051](#))
Buks E and Roukes M L 2001 Metastability and the Casimir effect in micromechanical systems *Europhys. Lett.* **54** 220 (Preprint [cond-mat/0008096](#))
- [9] Kenneth O and Klich I 2006 Opposites attract—a theorem about the Casimir force *Phys. Rev. Lett.* **97** 160401 (Preprint [quant-ph/0601011](#))
- [10] Osterwalder K and Schrader R 1973 Axioms for Euclidean Green's functions *Commun. Math. Phys.* **31** 83
Osterwalder K and Schrader R 1975 Axioms for Euclidean Green's functions: 2 *Commun. Math. Phys.* **42** 281
- [11] Osterwalder K and Seiler E 1978 Gauge field theories on the lattice *Ann. Phys.* **110** 440
Brydges D, Fröhlich J and Seiler E 1979 On the construction of quantized gauge fields: I. General results *Ann. Phys.* **121** 227
- [12] Seiler E 1978 Upper bound on the color confining potential *Phys. Rev. D* **18** 482
Borgs C and Seiler E 1983 Lattice Yang–Mills theory at nonzero temperature and the confinement problem *Commun. Math. Phys.* **91** 329
- [13] Bachas C 1986 Convexity of the quarkonium potential *Phys. Rev. D* **33** 2723 (note that this title makes a non-standard use of the words 'convex' and 'concave')
- [14] For a review of lattice gauge theory see e.g. Creutz M 1983 *Quarks, Gluons and Lattices (Cambridge Monographs on Mathematical Physics)* (Cambridge: Cambridge University Press)
- [15] Lüscher M 1977 Construction of a self-adjoint, strictly positive transfer matrix for Euclidean lattice gauge theories *Commun. Math. Phys.* **54** 283
- [16] Lüscher M 1982 Topology of lattice gauge fields *Commun. Math. Phys.* **85** 39
Woit P 1983 Topological charge in lattice gauge theory *Phys. Rev. Lett.* **51** 638
Alles B, D'Elia M, Giacomo A Di and Kirchner R 1998 A critical comparison of different definitions of topological charge on the lattice *Phys. Rev. D* **58** 114506 (Preprint [hep-lat/9711026](#))
- [17] Dotsenko V S and Vergeles S N 1980 Renormalizability of phase factors in the nonabelian gauge theory *Nucl. Phys. B* **169** 527
Fischler W 1977 Quark–anti-quark potential in QCD *Nucl. Phys. B* **129** 157
- [18] Balian R and Duplantier B 2004 Geometry of the Casimir effect Preprint [quant-ph/0408124](#)
Balian R and Duplantier B 1977 Electromagnetic waves near perfect conductors: 1. Multiple scattering expansions. Distribution of modes *Ann. Phys.* **104** 300

- Balian R and Duplantier B 1978 Electromagnetic waves near perfect conductors: 2. Casimir effect *Ann. Phys.* **112** 165
- [19] Boyer T H 1968 Quantum electromagnetic zero point energy of a conducting spherical shell and the Casimir model for a charged particle *Phys. Rev.* **174** 1764
- [20] Baacke J and Krusemann G 1986 Perturbative analysis of the divergent contributions to the Casimir energy *Z. Phys. C* **30** 413
- Barton G 2001 Perturbative Casimir energies of dispersive spheres, cubes and cylinders *J. Phys. A: Math. Gen.* **34** 4083
- Graham N, Jaffe R L, Khemani V, Quandt M, Schroeder O and Weigel H 2004 The Dirichlet Casimir problem *Nucl. Phys. B* **677** 379 (Preprint [hep-th/0309130](#))
- [21] Jackson J D 1962 *Classical Electrodynamics* (New York: Wiley) (for the relevant calculation, see section 6.2)
- [22] Cavalcanti R M 2004 Casimir force on a piston *Phys. Rev. D* **69** 065015 (Preprint [quant-ph/0310184](#))
- Hertzberg M P, Jaffe R L, Kardar M and Scardicchio A 2005 The Casimir piston Preprint [quant-ph/0509071](#)
- Edery A 2006 Casimir piston for massless scalar fields in three dimensions Preprint [hep-th/0610173](#)
- [23] Landau L D and Lifshitz E M 1960 *Electrodynamics of Continuous Media* (New York: Pergamon)
- [24] Wirzba A, Bulgac A and Magierski P 2006 Casimir interaction between normal or superfluid grains in the Fermi sea *J. Phys. A: Math. Gen.* **39** 6815 (Preprint [quant-ph/0511057](#))
- [25] Fulling S A and Wilson J H 2006 Repulsive Casimir Pistons Preprint [quant-ph/0608122](#)