

Comment on 'Position-dependent effective mass Dirac equations with  $PT$ -symmetric and non- $PT$ -symmetric potentials'

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2007 J. Phys. A: Math. Theor. 40 863

(<http://iopscience.iop.org/1751-8121/40/4/N01>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.146

The article was downloaded on 03/06/2010 at 06:20

Please note that [terms and conditions apply](#).

## COMMENT

## Comment on ‘Position-dependent effective mass Dirac equations with $PT$ -symmetric and non- $PT$ -symmetric potentials’

Omar Mustafa and S Habib Mazharimousavi

Department of Physics, Eastern Mediterranean University, G Magusa, North Cyprus, Mersin 10, Turkey

E-mail: [omar.mustafa@emu.edu.tr](mailto:omar.mustafa@emu.edu.tr) and [habib.mazhari@emu.edu.tr](mailto:habib.mazhari@emu.edu.tr)

Received 11 October 2006

Published 9 January 2007

Online at [stacks.iop.org/JPhysA/40/863](http://stacks.iop.org/JPhysA/40/863)

### Abstract

Jia and Dutra (2006 *J. Phys. A: Math. Gen.* **39** 11877) have considered the one-dimensional non-Hermitian complexified potentials with real spectra in the context of position-dependent mass in the Dirac equation. In their second example, a smooth step shape mass distribution is considered and a non-Hermitian non- $PT$ -symmetric Lorentz vector potential is obtained. They have mapped this problem into an exactly solvable Rosen–Morse Schrödinger model and claimed that the energy spectrum is real. The energy spectrum they have reported is pure imaginary or at best forms an empty set. Their claim on the reality of the energy spectrum is fragile, therefore.

PACS numbers: 03.65.Ge, 03.65.Pm, 03.65.Fd

(Some figures in this article are in colour only in the electronic version)

Very recently, Jia and Dutra [1] have presented a new procedure to construct a set of non-Hermitian potentials with real spectra for the one-dimensional Dirac particle endowed with position-dependent mass. They have reported that a Lorentz vector potential of the form

$$V(x) = \frac{i}{2} \frac{1}{M(x)} \frac{dM(x)}{dx} \quad (1)$$

would imply a Schrödinger-like Dirac equation

$$-\frac{d^2}{dx^2}\varphi(x) + V_{\text{eff}}(x)\varphi(x) = E^2\varphi(x), \quad (2)$$

where  $V_{\text{eff}}(x) = M(x)^2$ ,  $M(x)$  and  $E$  are the mass and the real energy of Dirac’s particle, respectively, and  $\varphi(x)$  is the upper component of the Dirac spinor (equations (1)–(9) in [1]).




Figure 1. A 3D plot of  $f(\eta, \Lambda)$  with  $\Lambda \in [0, 10]$  and  $-1 \leq \eta \leq 1$ .

In their second example (section 3.2 in [1]) they have considered a smooth step shape mass distribution

$$M(x) = M_0(1 + \eta \tanh \alpha x) \quad (3)$$

that led to a non- $\mathcal{PT}$ -symmetric non-Hermitian Lorentz vector potential

$$V(x) = \frac{i}{2} \frac{\alpha \eta \operatorname{sech}^2 \alpha x}{(1 + \eta \tanh \alpha x)} \quad (4)$$

where  $\eta$  is a small parameter that satisfies the condition  $|\eta| \leq 1$  to get a positive mass distribution. Under such settings, equation (2) reads a Rosen–Morse-type Schrödinger problem [2]. Employing the SUSQM method [3], Jia and Dutra [1] have reported that the reality of energy spectrum

$$E_n = \pm \sqrt{M_0^2(1 + \eta^2) - \frac{\eta^2 M_0^4}{\alpha^2(n + \delta_1)^2} - \alpha^2(n + \delta_1)^2}, \quad n = 0, 1, 2, \dots, \quad (5)$$

with

$$\delta_1 = \frac{1}{2} \left( 1 - \sqrt{1 + \frac{4\eta^2 M_0^2}{\alpha^2}} \right), \quad (6)$$

is guaranteed if and only if

$$M_0^2(1 + \eta^2) \geq \frac{\eta^2 M_0^4}{\alpha^2(n + \delta_1)^2} + \alpha^2(n + \delta_1)^2. \quad (7)$$

In this comment we shall show that even with the lowest possible value of the principle quantum number  $n$  (i.e.,  $n = 0$ ) the inequality in (7) cannot be satisfied. Hence, the reality of the spectrum in (5) is fragile (of course within the parameters' settings reported in [1] for this particular case).

To do so, we start with the introduction of a positive parameter  $\Lambda = \alpha/M_0 \geq 0$  to recast, with  $n = 0$ , the inequality (7) as

$$f(\eta, \Lambda) = (1 + \eta^2) - \frac{4\eta^2}{(\Lambda - \sqrt{\Lambda^2 + 4\eta^2})^2} - \frac{1}{4}(\Lambda - \sqrt{\Lambda^2 + 4\eta^2})^2 \geq 0. \quad (8)$$

Then we shall now test the validity of  $\Re \ni f(\eta, \Lambda) \geq 0$  for  $-1 \leq \eta \leq 1$  and  $\Lambda \geq 0$ , where  $\eta, \Lambda \in \mathbb{R}$ .

A 3D plot, in figure 1, of  $f(\eta, \Lambda)$  with  $\Lambda \in [0, 10]$  clearly indicates that  $0 \geq \Re \ni f(\eta, \Lambda) \not\geq 0$  over the physically acceptable parametric values. Therefore, the energy spectrum in (5) can only be pure imaginary (i.e.,  $E_n \in \mathbb{C}$  and  $E_n \notin \mathbb{R}$ ).

---

**References**

- [1] Jia C S and de Souza Dutra A 2006 *J. Phys. A: Math. Gen.* **39** 11877
- [2] Rosen N and Morse P M 1932 *Phys. Rev.* **42** 210
- [3] Cooper F, Khare A and Sukhatme U 1995 *Phys. Rep.* **251** 267