

## Spectral geometry, homogeneous spaces and differential forms with finite Fourier series

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys. A: Math. Theor. 41 135204

(<http://iopscience.iop.org/1751-8121/41/13/135204>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.147

The article was downloaded on 03/06/2010 at 06:38

Please note that [terms and conditions apply](#).

# Spectral geometry, homogeneous spaces and differential forms with finite Fourier series

C Dunn<sup>1</sup>, P Gilkey<sup>2</sup> and J H Park<sup>3,4</sup>

<sup>1</sup> Mathematics Department, California State University at San Bernardino, San Bernardino, CA 92407, USA

<sup>2</sup> Mathematics Department, University of Oregon, Eugene, OR 97403, USA

<sup>3</sup> Department of Mathematics, Sungkyunkwan University, Suwon 440-746, Korea

E-mail: [cmdunn@csusb.edu](mailto:cmdunn@csusb.edu), [gilkey@uoregon.edu](mailto:gilkey@uoregon.edu) and [parkj@skku.edu](mailto:parkj@skku.edu)

Received 9 August 2007, in final form 5 February 2008

Published 14 March 2008

Online at [stacks.iop.org/JPhysA/41/135204](http://stacks.iop.org/JPhysA/41/135204)

## Abstract

Let  $G$  be a compact Lie group acting transitively on Riemannian manifolds  $M_i$  and let  $\pi : M_1 \rightarrow M_2$  be a  $G$ -equivariant Riemannian submersion. We show that a smooth differential form  $\phi$  on  $M_2$  has finite Fourier series on  $M_2$  if and only if the pull back  $\pi^*\phi$  has finite Fourier series on  $M_1$ .

PACS numbers: 02.20.Qs, 02.30.Em, 02.30.Nw, 02.40.Vh

## 1. Introduction

The spectral geometry of homogeneous Riemannian submersions has been discussed by many authors in a variety of physical contexts. For example, Bérard-Bergery, and Bourguignon [4] study the Laplacian of a Riemannian submersion and provide an application to quantum physics. Also, Boiteux [5] studies the Coulomb potential via a fiber bundle formulation of mechanics, and we direct the reader who is interested in this particular physical application to remark 1.3; see [13, 17] for subsequent work. The spectral geometry of homogeneous Riemannian submersions also plays an important role in the study of non-bijective canonical transformations; we refer, for example, to the discussion in Lambert and Kibler [14] (see also [6, 11, 15, 16] for later work). Gilkey, Leahy and Park [9] study the spectral geometry of the Hopf fibration  $S^1 \rightarrow S^3 \rightarrow S^2$  and provide an overview of the potential physical applications. Their work continues in [10], where the authors give a more extensive discussion of these applications; see also Bao and Shen [1].

Let  $M$  be a compact smooth closed Riemannian manifold of dimension  $m$ , and let  $\Delta_M^p$  be the Laplace–Beltrami operator acting on the space  $C^\infty(\Lambda^p M)$  of smooth  $p$ -forms. Let  $\text{Spec}(\Delta_M^p)$  be the spectrum of  $\Delta_M^p$ ; this is a discrete countable set of non-negative real

<sup>4</sup> Corresponding author.

numbers. The associated eigenspaces  $E(\lambda, \Delta_M^p)$  are finite dimensional and there is a complete orthonormal decomposition

$$L^2(\Lambda^p M) = \bigoplus_{\lambda \in \text{Spec}(\Delta_M^p)} E(\lambda, \Delta_M^p) \tag{1a}$$

which we may use to decompose a smooth  $p$ -form  $\phi$  on  $M$  in the form  $\phi = \sum_{\lambda} \phi_{\lambda}$ , where  $\phi_{\lambda} \in E(\lambda, \Delta_M^p)$ . We say  $\phi$  has *finite Fourier series on  $M$*  if this is a finite sum. If  $p = 0$  and if  $M = S^1$ , then this yields, modulo a slight change of notation, the classical Fourier series decomposition  $f(\theta) = \sum_n a_n e^{in\theta}$  and a function has a finite Fourier series on the circle if and only if it is a trigonometric polynomial. There is an extensive literature on the subject with appropriate physical applications, several representative items being [2, 3, 7, 8, 18].

We say that  $M$  is a *homogeneous space* if there is a compact Lie group  $G$  which acts transitively on  $M$  by isometries; if  $H$  is the isotropy subgroup associated with some point  $P \in M$ , then we may identify  $M = G/H$ . We may choose a left-invariant metric  $\tilde{g}$  on  $G$  so  $g$  is the induced metric or, equivalently, that  $\pi : (G, \tilde{g}) \rightarrow (M, g)$  is a Riemannian submersion. The following is the main result of this paper:

**Theorem 1.1.** *Let  $\pi : G \rightarrow G/H$ , where  $H$  is a Lie subgroup of a compact Lie group  $G$ . Let  $\tilde{g}$  be a left-invariant Riemannian metric on  $G$  and let  $g$  be the induced Riemannian metric on  $G/H$ . Then a  $p$ -form  $\phi$  on  $G/H$  has finite Fourier series on  $G/H$  if and only if  $\pi^*\phi$  has finite Fourier series on  $G$ .*

There is an associated corollary which is useful in applications.

**Corollary 1.2.** *Let  $G$  be a compact Lie group acting transitively on Riemannian manifolds  $M_1$  and  $M_2$ . Let  $\pi : M_1 \rightarrow M_2$  be a  $G$ -equivariant Riemannian submersion. If  $\phi$  is a smooth  $p$ -form on  $M_2$ , then  $\phi$  has finite Fourier series on  $M_2$  if and only if  $\pi^*\phi$  has finite Fourier series on  $M_1$ .*

**Remark 1.3.** The Hopf fibration  $\pi : S^{2n+1} \rightarrow \mathbb{C}\mathbb{P}^n$  is a  $U(n+1)$  equivariant Riemannian submersion which is an important non-canonical transformation used to study the Coulomb problem, see, for example, the discussion in [5]. Corollary 1.2 shows  $\phi$  has finite Fourier series on  $\mathbb{C}\mathbb{P}^n$  if and only if  $\pi^*\phi$  has finite Fourier series on  $S^{2n+1}$ .

## 2. The proof of theorem 1.1

The central ingredient in our discussion is the classical Peter–Weyl theorem [12]. Let  $\text{Irr}(G)$  be the collection of equivalence classes of irreducible finite dimensional representations of  $G$ ; if  $\rho \in \text{Irr}(G)$ , let  $V_{\rho}$  be the associated representation space. The Hilbert space structure on  $L^2(\Lambda^p(G))$  depends on the particular Riemannian metric which is chosen. However different Riemannian metrics give rise to equivalent norms so the Banach space structure on this space is invariantly defined; this is a minor observation which will be useful in section 4. Left multiplication defines an action of  $G$  on  $L^2(\Lambda^p(G))$ . This action decomposes as a direct sum

$$L^2(\Lambda^p G) = \bigoplus_{\rho \in \text{Irr}(G)} W_{\rho}, \tag{2a}$$

where each  $W_{\rho}$  is a finite dimensional invariant subspace of  $L^2(\Lambda^p(G))$  which is isomorphic to the direct sum of finite number of copies of  $V_{\rho}$ . If  $\Phi$  is a smooth  $p$ -form on  $G$ , we may use equation (2a) to decompose  $\Phi = \sum_{\rho} \Phi_{\rho}$  for  $\Phi_{\rho} \in W_{\rho}$ . We say that  $\Phi$  has *finite  $G$ -representation series* on  $G$  if this sum is finite; we emphasize that this notion is independent of the particular Riemannian metric chosen.

Since we have taken the induced metric on  $G/H$ , the map  $\pi$  is a Riemannian submersion. Thus we have a pointwise estimate  $|\pi^*\phi(g)| = |\phi(\pi g)|$ . Since the volume form on  $G$  is bi-invariant, the fibers have constant volume. Thus

$$|\pi^*\phi|_{L^2(\Lambda^p(G))} = \sqrt{\text{vol}(H)}|\phi|_{L^2(\Lambda^p(G/H))}.$$

Consequently  $\pi^*$  is an injective  $G$ -equivariant map from  $L^2(\Lambda^p(G/H))$  to  $L^2(\Lambda^p(G))$  with closed image. The decomposition

$$L^2(\Lambda^p G) = \pi^*(L^2(\Lambda^p(G/H))) \oplus \{\pi^*(L^2(\Lambda^p(G/H)))\}^\perp$$

is  $G$ -equivariant. We therefore have an orthogonal direct sum decomposition of  $L^2(\Lambda^p(G/H))$  as a representation space for  $G$  in the form

$$L^2(\Lambda^p(G/H)) = \bigoplus_{\rho \in \text{Irr}(G)} X_\rho, \tag{2b}$$

where

$$\pi^* X_\rho = W_\rho \cap \pi^*(L^2(\Lambda^p(G/H))). \tag{2c}$$

We say that a  $p$ -form  $\phi$  on  $G/H$  has *finite  $G$ -representation series* if the expansion  $\phi = \sum_\rho \phi_\rho$  given by equation (2a) is finite. Theorem 1.1 will follow from the following:

**Lemma 2.1.** *Adopt the notation established above. Let  $\phi$  be a  $p$ -form on  $G/H$ . Fix a left-invariant  $\tilde{g}$  metric on  $G$  and let  $g$  be the induced metric on  $G/H$ . The following assertions are equivalent:*

- (i)  $\phi$  has finite Fourier series on  $G/H$ .
- (ii)  $\phi$  has finite  $G$ -representation series on  $G/H$ .
- (iii)  $\pi^*\phi$  has finite Fourier series on  $G$ .
- (iv)  $\pi^*\phi$  has finite  $G$ -representation series on  $G$ .

We remark that elliptic regularity shows such a  $\phi$  is necessarily smooth.

**Proof.** The equivalence of assertions (ii) and (iv) is immediate from equation (2a). We argue as follows to prove that assertion (i) implies assertion (ii). Suppose that  $\phi$  has finite Fourier series on  $G/H$ . Since  $G$  acts by isometries,  $G$  commutes with the Laplacian. Thus  $E(\lambda, \Delta_{G/H}^p)$  is a finite dimensional representation space for  $G$ . Only a finite number of representations occur in the representation decomposition of  $E(\lambda, \Delta_{G/H}^p)$  and thus any eigen  $p$ -form on  $G/H$  has finite  $G$ -representation series on  $G/H$ ; more generally, of course, any finite sum of eigen  $p$ -forms on  $G/H$  has finite  $G$ -representation series on  $G/H$ . This shows that assertion (i) implies assertion (ii); a similar argument shows assertion (iii) implies assertion (iv).

Each representation appears with finite multiplicity in  $L^2(\Lambda^p(G/H))$ . Thus each representation appears in the decomposition of  $E(\lambda, \Delta_{G/H}^p)$  for only a finite number of  $\lambda$ . Thus any element of  $X_\rho$  has finite Fourier series and more generally any  $p$ -form on  $G/H$  with finite  $G$ -representation series has finite Fourier series. Thus assertion (ii) implies assertion (i); similarly, assertion (iv) implies assertion (iii).  $\square$

### 3. The proof of corollary 1.2

Let  $\pi : M_1 \rightarrow M_2$  be a  $G$ -equivariant Riemannian submersion; this means that we may express  $M_i = G/H_i$ , where  $H_1 \subset H_2 \subset G$ . Let  $\pi_i : G \rightarrow G/H_i$  be the natural projections. We then have  $\pi \pi_1 = \pi_2$  and thus  $\pi_2^* = \pi_1^* \pi^*$ . Let  $\phi$  be a smooth  $p$ -form on  $G/H_2$ . We apply theorem 1.1 to derive the following chain of equivalent statements from which corollary 1.2 will follow:

- (i)  $\phi$  has finite Fourier series on  $G/H_2$ .
- (ii)  $\pi_2^*\phi$  has finite Fourier series on  $G$ .
- (iii)  $\pi_1^*(\pi^*\phi)$  has finite Fourier series on  $G$ .
- (iv)  $\pi^*\phi$  has finite Fourier series on  $G/H_1$ .

#### 4. Conclusions and open problems

Our methods in fact show a bit more. Let  $g_i$  be two left-invariant metrics on  $G$  and let  $\phi$  be a smooth  $p$ -form on  $G$ . Then  $\phi$  has finite Fourier series with respect to  $g_1$  if and only if  $\phi$  has finite Fourier series with respect to  $g_2$  since both conditions are equivalent to  $\phi$  having finite representation series and this notion is independent of the particular metric chosen.

Cayley multiplication defines a Riemannian submersion  $\pi : S^7 \times S^7 \rightarrow S^7$ . The group of isometries commuting with this action does not, however, act transitively on  $S^7 \times S^7$  and theorem 1.1 is not applicable. Our research continues in this area as this example has important physical applications (see, for example, the discussion in Lambert and Kibler [14]).

#### Acknowledgments

Research of C Dunn partially supported by a CSUSB faculty research grant. Research of P Gilkey partially supported by the Max Planck Institute in the Mathematical Sciences (Leipzig, Germany). Research of both C Dunn and P Gilkey partially supported by the University of Santiago (Spain) (Project MTM2006-01432). Research of J H Park partially supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD) KRF-2007-531-C00008.

#### References

- [1] Bao D and Shen Z 2002 *J. London Math. Soc.* **66** 453
- [2] Blevins R D 1997 *J. Sound Vib.* **208** 617
- [3] Blevins R D 2002 *J. Appl. Mech.—Trans. ASME* **69** 317
- [4] Berard-Bergery L and Bourguignon J-P 1982 *Illinois J. Math.* **26** 181
- [5] Boiteux M 1982 *J. Math. Phys.* **23** 1311
- [6] Coffey M W 2007 *Phys. Lett. A* **362** 352
- [7] Dinur N and Wulich D 2001 *IEEE Trans. Commun.* **49** 1063
- [8] Fisher C and Schmidt B 2006 *J. Aust. Math. Soc.* **81** 21
- [9] Gilkey P, Leahy J and Park J H 1996 *J. Phys. A Math. Gen.* **29** 5645
- [10] Gilkey P, Leahy J and Park J H 1999 *Spectral Geometry, Riemannian Submersions, and the Gromov-Lawson Conjecture* (London: Chapman and Hall)
- [11] Hakobyan Y and Ter-Antonyan V 2005 *Phys. At. Nuclei* **68** 1709
- [12] Hall B 2003 *Lie Groups, Lie Algebras and Representations* (Berlin: Springer)
- [13] Kibler M 2004 *Mol. Phys.* **102** 1221
- [14] Lambert D and Kibler M 1988 *J. Phys. A. Math. Gen.* **21** 307
- [15] Leach P and Nucci M 2004 *J. Math. Phys.* **45** 3590
- [16] Mardoyan L, Pogosyan G and Sissakian A 2003 *Theor. Math. Phys.* **135** 808
- [17] Michel L and Zhilinskii B 2001 *Phys. Rep.* **341** 173
- [18] Reut Z 2000 *J. Sound Vib.* **232** 490