

## Characterization of the critical magnetic field in the Dirac–Coulomb equation

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys. A: Math. Theor. 41 185303

(<http://iopscience.iop.org/1751-8121/41/18/185303>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.148

The article was downloaded on 03/06/2010 at 06:47

Please note that [terms and conditions apply](#).

# Characterization of the critical magnetic field in the Dirac–Coulomb equation

J Dolbeault<sup>1</sup>, M Esteban<sup>1</sup> and M Loss<sup>2</sup>

<sup>1</sup> Ceremade (UMR CNRS 7534), Université Paris Dauphine, Place de Lattre de Tassigny, F-75775 Paris Cédex 16, France

<sup>2</sup> School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332, USA

E-mail: [dolbeaul@ceremade.dauphine.fr](mailto:dolbeaul@ceremade.dauphine.fr), [esteban@ceremade.dauphine.fr](mailto:esteban@ceremade.dauphine.fr) and [loss@math.gatech.edu](mailto:loss@math.gatech.edu)

Received 23 December 2007, in final form 17 March 2008

Published 18 April 2008

Online at [stacks.iop.org/JPhysA/41/185303](http://stacks.iop.org/JPhysA/41/185303)

## Abstract

We consider a relativistic hydrogenic atom in a strong magnetic field. The ground-state level depends on the strength of the magnetic field and reaches the lower end of the spectral gap of the Dirac–Coulomb operator for a certain critical value, the *critical magnetic field*. We also define a critical magnetic field in a Landau level ansatz. In both cases, when the charge  $Z$  of the nucleus is not too small, these critical magnetic fields are huge when measured in tesla, but not so big when the equation is written in dimensionless form. When computed in the Landau level ansatz, orders of magnitude of the critical field are correct, as well as the dependence in  $Z$ . The computed value is however significantly too big for a large  $Z$ , and the wavefunction is not well approximated. Hence, accurate numerical computations involving the Dirac equation cannot systematically rely on the Landau level ansatz. Our approach is based on a scaling property. The critical magnetic field is characterized in terms of an equivalent eigenvalue problem. This is our main analytical result, and also the starting point of our numerical scheme.

PACS number: 31.30.J

Mathematics Subject Classification: 35Q40, 35Q75, 46N50, 81Q10, 34L40, 35P05, 47A05, 47N50, 81V45

## 1. Introduction

It is widely accepted in physics that eigenstates in strong magnetic fields can be well approximated in a Landau level ansatz. In this paper, we investigate the validity of such an ansatz in the case of the magnetic Dirac–Coulomb equation for large magnetic fields. More specifically, we study the *critical threshold* for the magnetic field defined as the smallest value

of the magnetic field for which the lowest eigenvalue (the ground state) in the gap of the Dirac operator reaches its lower end. After a reformulation in terms of an equivalent minimization problem, we compare the values of the critical magnetic field with and without ansatz, both from an analytical and a numerical point of view.

Our main theorem is stated in section 2. It characterizes the critical magnetic field as a function of the lowest energy for an explicit eigenvalue problem. Section 3 will be devoted to its proof. A Landau level ansatz is then defined in section 4 and some comparison results for the critical magnetic field, with and without ansatz, are given there. A numerical method based on theorem 1 has been implemented in both cases. Results are given in section 5 and compared with earlier works, see Schlüter *et al* (1985). They show that for large coupling constants (in the electrostatic field), the critical threshold is well below the critical threshold in the Landau ansatz.

Critical magnetic fields are huge and can eventually be encountered only in some extreme situations such as *magnetars*, which are neutron stars with intense magnetic fields, see Kouveliotou *et al* (1999). This is the only known domain of physics for which our computations might eventually be relevant, see Dolbeault *et al* (2007) for a discussion. In the Dirac–Coulomb model, the value of the critical magnetic field only provides an order of magnitude of the field strength for which pair creation could eventually occur. Such a phenomenon should of course be studied in a full QED framework and the computations done using the Dirac–Coulomb operator are only an indication on the scales that should be taken into account. See Pickl (2005) for more details. For a review on large magnetic fields in physics, see Duncan (2000).

A description of the non-relativistic many-electron atom under extreme magnetic fields has been given in various papers among them Fushiki *et al* (1992), Lieb *et al* (1994a, 1994b) as well as Baumgartner *et al* (2000), Baumgartner and Seiringer (2001), Brummelhuis and Duclos (2006), Medin and Lai (2006), von Milczewski *et al* (1996). Most of these papers cover the many-body problem and none of them deal with the quantitative issue of determining the critical magnetic field.

## 2. Main results

The magnetic Dirac operator with Coulomb potential  $v/|x|$  can be written as

$$H_B := \begin{pmatrix} \mathbb{I} - v/|x| & -i\sigma \cdot (\nabla - iA) \\ -i\sigma \cdot (\nabla - iA) & -\mathbb{I} - v/|x| \end{pmatrix} \quad (1)$$

where  $A$  is a magnetic potential corresponding to  $B$ , and  $\mathbb{I}$  and  $\sigma_k$  are respectively the identity and the Pauli matrices

$$\mathbb{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2)$$

Let  $B = (0, 0, B)$  be a constant magnetic field and  $A_B$  the associated magnetic potential. For any  $x = (x_1, x_2, x_3) \in \mathbb{R}^3$ , define

$$P_B := -i\sigma \cdot (\nabla - iA_B(x)), \quad A_B(x) := \frac{B}{2} \begin{pmatrix} -x_2 \\ x_1 \\ 0 \end{pmatrix}, \quad (3)$$

and consider the functional

$$J[\phi, \lambda, v, B] := \int_{\mathbb{R}^3} \left( \frac{|P_B \phi|^2}{1 + \lambda + \frac{v}{|x|}} + (1 - \lambda)|\phi|^2 - \frac{v}{|x|}|\phi|^2 \right) d^3x \quad (4)$$

on the set of admissible functions  $\mathcal{A}(v, B) := \{\phi \in C_0^\infty(\mathbb{R}^3) : \|\phi\|_{L^2(\mathbb{R}^3)}^2 = 1, \lambda \mapsto J[\phi, \lambda, v, B] \text{ changes sign in } (-1, +\infty)\}$  which expresses the full theory in terms of the upper components of the full four-spinor usually used in Dirac theory. The essential spectrum of  $H_B$  is  $\mathbb{R} \setminus (-1, 1)$ . By Dolbeault *et al* (2007) (theorem 1), the smallest eigenvalue in  $(-1, 1)$  of  $H_B$  is

$$\lambda_1(v, B) := \inf_{\phi \in \mathcal{A}(v, B)} \lambda[\phi, v, B] \tag{5}$$

where  $\lambda = \lambda[\phi, v, B]$  is either the unique solution to  $J[\phi, \lambda, v, B] = 0$  if  $\phi \in \mathcal{A}(v, B)$ , or  $\lambda[\phi, v, B] = -1$  if  $J[\phi, -1, v, B] \leq 0$ . Also see Dolbeault *et al* (2007) for the relation of the two-component spinors in  $\mathcal{A}(v, B)$  with the four-component spinors and the action of  $H_B$  on them. It is important to note that the reduction of the Dirac eigenvalue problem to an eigenvalue problem on two-component spinors is done without approximations.

The *critical magnetic field* is defined by

$$B(v) := \inf\{B > 0 : \liminf_{b \nearrow B} \lambda_1(v, b) = -1\}. \tag{6}$$

Define the auxiliary functional

$$\mathcal{E}_{B,v}[\phi] := \int_{\mathbb{R}^3} \frac{|x|}{v} |P_B \phi|^2 d^3x - \int_{\mathbb{R}^3} \frac{v}{|x|} |\phi|^2 d^3x, \tag{7}$$

that is  $\mathcal{E}_{B,v}[\phi] + 2\|\phi\|_{L^2(\mathbb{R}^3)}^2 = J[\phi, -1, v, B]$ . The scaling  $\phi_B := B^{3/4}\phi(B^{1/2}x)$  preserves the  $L^2$  norm, and yields

$$\mathcal{E}_{B,v}[\phi_B] = \sqrt{B}\mathcal{E}_{1,v}[\phi]. \tag{8}$$

We define

$$\mu(v) := \inf_{0 \neq \phi \in C_0^\infty(\mathbb{R}^3)} \frac{\mathcal{E}_{1,v}[\phi]}{\|\phi\|_{L^2(\mathbb{R}^3)}^2}. \tag{9}$$

Recall that  $\lambda_1(v, B)$  is characterized as an eigenvalue of  $H_B$  only as long as it takes values in  $(-1, 1)$ . If we could take the limit  $B \rightarrow B(v)$ , we would formally get that  $-1 = \lambda_1(v, B(v))$ , which, still formally, amounts to  $\inf_{0 \neq \phi \in C_0^\infty(\mathbb{R}^3)} J[\phi, -1, v, B] = 0$ . It is therefore natural to expect that  $\sqrt{B(v)}\mu(v) + 2 = 0$ . Proving this is the purpose of our main result.

**Theorem 1.** *For all  $v \in (0, 1)$ ,  $\mu(v)$  is negative, finite,*

$$B(v) = \frac{4}{\mu(v)^2} \tag{10}$$

*and  $B(v)$  is a continuous, monotone decreasing function of  $v$  on  $(0, 1)$ .*

### 3. Proofs

#### 3.1. Preliminary results

**Lemma 2.** *On the interval  $(0, 1)$ , the function  $v \mapsto \mu(v)$  is continuous, monotone decreasing and takes only negative real values.*

**Proof.** The monotonicity of the function  $\mu(v)$  is a consequence of its definition. On the other hand, the functional

$$(v, \phi) \mapsto v\mathcal{E}_{1,v}[\phi] = \int_{\mathbb{R}^3} |x| |P_1 \phi|^2 d^3x - \int_{\mathbb{R}^3} \frac{v^2}{|x|} |\phi|^2 d^3x \tag{11}$$

is a concave, bounded function of  $\nu \in (0, 1)$ , for any  $\phi \in C_0^\infty(\mathbb{R}^3)$ , and so is its infimum with respect to  $\phi$ . A bounded concave function is continuous.

Next, consider the function

$$\phi(x) := \sqrt{\frac{B}{2\pi}} e^{-\frac{B}{4}(|x_1|^2 + |x_2|^2)} \begin{pmatrix} f(x_3) \\ 0 \end{pmatrix} \quad \forall x = (x_1, x_2, x_3) \in \mathbb{R}^3 \quad (12)$$

for some  $f \in C_0^\infty(\mathbb{R}, \mathbb{R})$  such that  $f \equiv 1$  for  $|x| \leq \delta$ ,  $\delta > 0$ , and  $\|f\|_{L^2(\mathbb{R}^*)} = 1$ . Note that  $\phi \in \text{Ker}(P_B + i\sigma_3 \partial_{x_3})$  and so,  $P_B \phi = -i\sigma_3 \partial_{x_3} \phi$ . Moreover, the function  $\phi$  is normalized in  $L^2(\mathbb{R}^3)$ . The same computations as in (Dolbeault et al 2007, proof of proposition 6) show that

$$\mathcal{E}_{B,\nu}[\phi] \leq \frac{C_1}{\nu} + C_2 \nu - C_3 \nu \log B, \quad (13)$$

where  $C_i, i = 1, 2, 3$ , are positive constants which depend only on  $f$ . For  $B \geq 1$  large enough,  $\mathcal{E}_{B,\nu}[\phi] < 0$ . With  $\phi_{1/B}(x) = B^{-3/4} \phi(B^{-1/2}x)$ ,  $\mathcal{E}_{1,\nu}[\phi_{1/B}] = B^{-1/2} \mathcal{E}_{B,\nu}[\phi] < 0$  by (8). This proves that  $\mu(\nu)$  is negative.  $\square$

**Lemma 3.** For any  $a, b \in \mathbb{R}^d$  and any  $\varepsilon > 0$ ,

$$|a|^2 \geq \frac{|a+b|^2}{1+\varepsilon} - \frac{|b|^2}{\varepsilon}. \quad (14)$$

**Proof .** A simple computation shows that

$$\varepsilon(1+\varepsilon) \left[ |a|^2 - \frac{|a+b|^2}{1+\varepsilon} + \frac{|b|^2}{\varepsilon} \right] = (\varepsilon a - b)^2 \geq 0. \quad (15) \quad \square$$

### 3.2. Proof of theorem 2.1

Let us first prove that  $B(\nu) \leq \tilde{B}(\nu) := 4\mu(\nu)^2$ . By definition of  $\mu(\nu)$  and scaling,

$$\tilde{B}(\nu) = \sup \{ B > 0 : \inf_{\phi} (\mathcal{E}_{B,\nu}[\phi] + 2\|\phi\|_{L^2(\mathbb{R}^3)}^2) \geq 0 \}. \quad (16)$$

By definition of  $B(\nu)$ ,  $B < B(\nu)$  means that  $\lambda_1(\nu, B) > -1$ . Since  $\lambda_1(\nu, B)$  is the infimum of  $\lambda[\phi, \nu, B]$  with respect to  $\phi \in C_0^\infty(\mathbb{R}^3)$  and  $J[\phi, \lambda, \nu, B]$  is decreasing in  $\lambda$ ,

$$\mathcal{E}_{B,\nu}[\phi] + 2\|\phi\|_{L^2(\mathbb{R}^3)}^2 \geq J[\phi, \lambda_1(\nu, B), \nu, B] \geq J[\phi, \lambda[\phi, \nu, B], \nu, B] = 0 \quad (17)$$

for all  $\phi$ , so that  $B \leq \tilde{B}(\nu)$ . This proves that  $B(\nu) \leq \tilde{B}(\nu)$ . To prove the opposite inequality, let  $B = B(\nu)$  and consider a sequence  $(\nu_n)_{n \in \mathbb{N}}$  such that  $\nu_n \in (0, \nu)$ ,  $\lim_{n \rightarrow \infty} \nu_n = \nu$ ,  $\lambda^n := \lambda_1(\nu_n, B) > -1$  and  $\lim_{n \rightarrow \infty} \lambda^n = -1$ . Let  $\phi_n$  be the optimal function associated with  $\lambda^n$ :  $J[\phi_n, \lambda^n, \nu_n, B] = 0$ .

We define a sequence of truncation functions  $(\chi_n)_{n \in \mathbb{N}}$  as follows. Consider first a nonnegative smooth radial function  $\chi$  on  $\mathbb{R}^+$  such that  $\chi \equiv 1$  on  $[0, 1]$ ,  $0 \leq \chi \leq 1$  and  $\chi \equiv 0$  on  $[2, \infty)$ . Then we set  $\chi_n(x) := \chi(|x|/R_n)$  for some increasing sequence  $(R_n)_{n \in \mathbb{N}}$  such that  $\lim_{n \rightarrow \infty} R_n = \infty$ . By applying lemma 3 to

$$P_B \phi_n = \underbrace{(P_B \tilde{\phi}_n) \chi_n}_{=a} + \underbrace{[-(P_0 \chi_n) \phi_n]}_{=b}, \quad (18)$$

where  $\tilde{\phi}_n := \phi_n \chi_n$ , we get

$$|P_B \phi_n|^2 \geq \frac{|(P_B \tilde{\phi}_n) \chi_n|^2}{1 + \varepsilon_n} - \frac{|(P_0 \chi_n) \phi_n|^2}{\varepsilon_n}, \quad (19)$$

for some sequence  $(\varepsilon_n)_{n \in \mathbb{N}}$  of positive numbers, to be fixed. Hence, using the fact that  $0 \leq \chi_n^2 \leq 1$ , we get

$$\int_{\mathbb{R}^3} \frac{|P_B \phi_n|^2}{1 + \lambda^n + \frac{\nu_n}{|x|}} d^3x \geq \frac{1}{1 + \varepsilon_n} \int_{\mathbb{R}^3} \frac{|P_B \tilde{\phi}_n|^2}{1 + \lambda^n + \frac{\nu_n}{|x|}} d^3x - \frac{1}{\varepsilon_n} \int_{\mathbb{R}^3} \frac{|(P_0 \chi_n) \phi_n|^2}{1 + \lambda^n + \frac{\nu_n}{|x|}} d^3x. \quad (20)$$

The function  $\tilde{\phi}_n$  is supported in the ball  $B(0, 2R_n)$ : with  $\mu_n := (1 + \varepsilon_n)[2(1 + \lambda^n)R_n + \nu_n]$ ,

$$\frac{1}{1 + \varepsilon_n} \int_{\mathbb{R}^3} \frac{|P_B \tilde{\phi}_n|^2}{1 + \lambda^n + \frac{\nu_n}{|x|}} d^3x \geq \frac{1}{\mu_n} \int_{\mathbb{R}^3} |x| |P_B \tilde{\phi}_n|^2 d^3x. \quad (21)$$

We choose  $\varepsilon_n$  and  $R_n$  such that

$$\lim_{n \rightarrow \infty} \varepsilon_n = 0, \quad \lim_{n \rightarrow \infty} R_n = \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} (1 + \lambda^n)R_n = 0, \quad (22)$$

so that

$$\lim_{n \rightarrow \infty} \mu_n = \nu. \quad (23)$$

The function  $(P_0 \chi_n)$  is supported in  $B(0, 2R_n) \setminus B(0, R_n)$ : there exists constant  $\kappa$  depending on  $\|\chi'\|_{L^\infty(1,2)}$  such that  $|P_0 \chi_n|^2 \leq \kappa R_n^{-2}$  and as a consequence,

$$\frac{1}{\varepsilon_n} \int_{\mathbb{R}^3} \frac{|(P_0 \chi_n) \phi_n|^2}{1 + \lambda^n + \frac{\nu_n}{|x|}} d^3x \leq \frac{\kappa}{\varepsilon_n R_n [(1 + \lambda^n)R_n + \frac{\nu_n}{2}]} \int_{\mathbb{R}^3} |\phi_n|^2 d^3x. \quad (24)$$

Moreover,

$$\nu_n \int_{\mathbb{R}^3} \frac{|\phi_n|^2}{|x|} d^3x - \nu_n \int_{\mathbb{R}^3} \frac{|\tilde{\phi}_n|^2}{|x|} d^3x \leq \frac{\nu_n}{R_n} \int_{\mathbb{R}^3} |\phi_n|^2 d^3x. \quad (25)$$

Thus, with  $\eta_n := \kappa(\varepsilon_n R_n((1 + \lambda^n)R_n + \frac{\nu_n}{2})) + \nu_n/R_n$ , we can write

$$\begin{aligned} 0 = J[\phi_n, \lambda^n, \nu_n, B] &\geq \frac{1}{\mu_n} \int_{\mathbb{R}^3} |x| |P_B \tilde{\phi}_n|^2 d^3x - \nu_n \int_{\mathbb{R}^3} \frac{|\tilde{\phi}_n|^2}{|x|} d^3x \\ &\quad + (1 - \lambda^n - \eta_n) \int_{\mathbb{R}^3} |\phi_n|^2 d^3x. \end{aligned} \quad (26)$$

Assume further that

$$\lim_{n \rightarrow \infty} \varepsilon_n R_n = \infty, \quad (27)$$

so that  $1 - \lambda^n \geq \eta_n \rightarrow 0$  as  $n \rightarrow \infty$ . Using again the fact that  $0 \leq \chi_n^2 \leq 1$ , we get

$$\begin{aligned} 0 = J[\phi_n, \lambda^n, \nu_n, B] &\geq \frac{1}{\mu_n} \int_{\mathbb{R}^3} |x| |P_B \tilde{\phi}_n|^2 d^3x - \nu_n \int_{\mathbb{R}^3} \frac{|\tilde{\phi}_n|^2}{|x|} d^3x \\ &\quad + (1 - \lambda^n - \eta_n) \int_{\mathbb{R}^3} |\tilde{\phi}_n|^2 d^3x. \end{aligned} \quad (28)$$

Let  $\tilde{\nu}_n = \sqrt{\mu_n \nu_n}$ . We have obtained

$$\begin{aligned} &\frac{1}{\tilde{\nu}_n} \int_{\mathbb{R}^3} |x| |P_B \tilde{\phi}_n|^2 d^3x - \tilde{\nu}_n \int_{\mathbb{R}^3} \frac{|\tilde{\phi}_n|^2}{|x|} d^3x + 2 \int_{\mathbb{R}^3} |\tilde{\phi}_n|^2 d^3x \\ &\leq \left[ 2 - \sqrt{\frac{\mu_n}{\nu_n}} (1 - \lambda^n - \eta_n) \right] \int_{\mathbb{R}^3} |\tilde{\phi}_n|^2 d^3x. \end{aligned} \quad (29)$$

The left-hand side is a decreasing function of  $\tilde{\nu}_n$ . Since  $\lim_{n \rightarrow \infty} \tilde{\nu}_n = \nu$ , for any  $\nu' > \nu$ , for  $n$  large enough,

$$\begin{aligned} &\frac{1}{\nu'} \int_{\mathbb{R}^3} |x| |P_B \tilde{\phi}_n|^2 d^3x - \nu' \int_{\mathbb{R}^3} \frac{|\tilde{\phi}_n|^2}{|x|} d^3x + 2 \int_{\mathbb{R}^3} |\tilde{\phi}_n|^2 d^3x \\ &\leq \left[ 2 - \sqrt{\frac{\mu_n}{\nu_n}} (1 - \lambda^n - \eta_n) \right] \int_{\mathbb{R}^3} |\tilde{\phi}_n|^2 d^3x. \end{aligned} \quad (30)$$

We observe that by construction  $\tilde{\phi}_n$  is non-trivial. By homogeneity, one can even assume that  $\|\tilde{\phi}_n\|_{L^2(\mathbb{R}^3)} = 1$ . Since  $\lim_{n \rightarrow \infty} \sqrt{\frac{\mu_n}{v_n}}(1 - \lambda^n - \eta_n) = 2$ ,

$$\tilde{B}(v') \leq B = B(v) \quad \forall v' > v. \tag{31}$$

By lemma 2,  $v' \mapsto \tilde{B}(v')$  is continuous. This proves that  $\tilde{B}(v) \leq B(v)$ .

#### 4. A Landau level ansatz

In analogy with what is done in non-relativistic quantum mechanics, we denote by *first Landau level* for a constant magnetic field of strength  $B$ , see Dolbeault *et al* (2007), the space of all functions  $\phi$  which are linear combinations of the functions

$$\phi_\ell := \frac{B}{\sqrt{2\pi 2^\ell \ell!}} (x_2 + ix_1)^\ell e^{-Bs^2/4} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \ell \in \mathbb{N}, \quad s^2 = x_1^2 + x_2^2, \tag{32}$$

where the coefficients depend only on  $x_3$ , i.e.,

$$\phi(x) = \sum_{\ell} f_\ell(x_3) \phi_\ell(x_1, x_2). \tag{33}$$

In this section, we shall restrict the functional  $\mathcal{E}_{B,v}$  to the first Landau level. In this framework, that we shall call the *Landau level ansatz*, we also define a critical field by

$$B_{\mathcal{L}}(v) := \inf \{ B > 0 : \liminf_{b \nearrow B} \lambda_1^{\mathcal{L}}(v, b) = -1 \}, \tag{34}$$

where

$$\lambda_1^{\mathcal{L}}(v, B) := \inf_{\phi \in \mathcal{A}(v, B), \Pi^\perp \phi = 0} \lambda[\phi, v, B]. \tag{35}$$

Here  $\Pi$  is the projection of  $\phi$  onto the first Landau level, and  $\Pi^\perp := \mathbb{I} - \Pi$ .

One can prove in the Landau level ansatz a result which is the exact counterpart of theorem 1. For any  $v \in (0, 1)$ , if

$$\mu_{\mathcal{L}}(v) := \inf_{\phi \in \mathcal{A}(v, B), \Pi^\perp \phi = 0} \mathcal{E}_{1,v}[\phi], \tag{36}$$

then

$$B_{\mathcal{L}}(v) = \frac{4}{\mu_{\mathcal{L}}(v)^2}. \tag{37}$$

The goal of this section is to compare  $\mu_{\mathcal{L}}(v)$  with  $\mu(v)$  given by (9). By definition of these quantities, we have

$$\mu(v) \leq \mu_{\mathcal{L}}(v). \tag{38}$$

With the notation  $s = \sqrt{x_1^2 + x_2^2}$  and  $z = x_3$ , if  $\phi$  is in the first Landau level, then

$$\mathcal{E}_{1,v}[\phi] = \sum_{\ell} \frac{1}{v} \int_0^\infty b_\ell f_\ell'^2 dz - v \int_0^\infty a_\ell f_\ell^2 dz, \tag{39}$$

where

$$a_\ell(z) := \left( \phi_\ell, \frac{1}{r} \phi_\ell \right)_{L^2(\mathbb{R}^2, \mathbb{C}^2)} = \frac{1}{2^\ell \ell!} \int_0^{+\infty} \frac{s^{2\ell+1} e^{-s^2/2}}{\sqrt{s^2 + z^2}} ds \tag{40}$$

and

$$b_\ell(z) := (\phi_\ell, r \phi_\ell)_{L^2(\mathbb{R}^2, \mathbb{C}^2)} = \frac{1}{2^\ell \ell!} \int_0^{+\infty} s^{2\ell+1} e^{-s^2/2} \sqrt{s^2 + z^2} ds. \tag{41}$$

A simple integration by parts shows that  $(\phi_\ell, F(r)\phi_\ell)_{L^2(\mathbb{R}^2, \mathbb{C}^2)}$  is increasing (respectively decreasing) in  $\ell$  whenever  $F(r)$  is increasing (respectively decreasing). Since  $a_\ell$  and  $b_\ell$  only depend on  $|z|$ ,  $\mu_{\mathcal{L}}(\nu)$  is also achieved by functions which only depend on  $|z|$  as well. It follows that

$$\sum_\ell \left( \frac{1}{\nu} \int_0^\infty b_\ell f_\ell'^2 dz - \nu \int_0^\infty a_\ell f_\ell^2 dz \right) \geq \frac{1}{\nu} \int_0^\infty b_0 \left( \sum_\ell f_\ell'^2 \right) dz - \nu \int_0^\infty a_0 \left( \sum_\ell f_\ell^2 \right) dz. \tag{42}$$

Using the inequality

$$\left| \frac{d}{dz} \sqrt{\sum_\ell f_\ell^2} \right|^2 \leq \sum_\ell \left| \frac{d}{dz} f_\ell \right|^2, \tag{43}$$

we find that

$$\mathcal{E}_{1,\nu}[\phi] \geq \frac{1}{\nu} \int_0^\infty b_0 |f'|^2 dz - \nu \int_0^\infty a_0 f^2 dz, \tag{44}$$

with  $f := \sqrt{\sum_\ell f_\ell^2}$ . In other words, for our minimization purpose, it is sufficient to consider functions of the form

$$\phi(x) = f(z) \frac{e^{-s^2/4}}{\sqrt{2\pi}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \tag{45}$$

We observe that

$$\frac{1}{2} \int_{\mathbb{R}^3} |\phi|^2 d^3\vec{x} = \int_0^\infty f^2 dz \tag{46}$$

and

$$\frac{1}{2} \mathcal{E}_{1,\nu}[\phi] = \frac{1}{\nu} \int_0^\infty b f'^2 dz - \nu \int_0^\infty a f^2 dz := \mathcal{L}_\nu[f], \tag{47}$$

with  $a = a_0, b = b_0$ , i.e.,

$$b(z) = \int_0^\infty \sqrt{s^2 + z^2} s e^{-s^2/2} ds \quad \text{and} \quad a(z) = \int_0^\infty \frac{s e^{-s^2/2}}{\sqrt{s^2 + z^2}} ds. \tag{48}$$

The minimization problem in the Landau level ansatz is now reduced to

$$\mu_{\mathcal{L}}(\nu) = \inf_f \frac{\mathcal{L}_\nu[f]}{\|f\|_{L^2(\mathbb{R}^+)}^2}. \tag{49}$$

It is a non-trivial problem to estimate how close  $\mu(\nu)$  and  $\mu_{\mathcal{L}}(\nu)$  are. Let

$$\mathcal{L}_\nu^-[f] := \frac{1}{\nu} \int_0^\infty \frac{1}{a} f'^2 dz - \nu \int_0^\infty a f^2 dz \tag{50}$$

and

$$\mathcal{L}_\nu^+[f] := \frac{1}{\nu} \int_0^\infty b f'^2 dz - \nu \int_0^\infty \frac{1}{b} f^2 dz, \tag{51}$$

with corresponding infima  $\mu_{\mathcal{L}}^-(\nu)$  and  $\mu_{\mathcal{L}}^+(\nu)$ .

**Lemma 4.** For any  $\nu \in (0, 1)$ ,

$$\mu_{\mathcal{L}}^-(\nu) \leq \mu_{\mathcal{L}}(\nu) \leq \mu_{\mathcal{L}}^+(\nu). \tag{52}$$



**Proof .** This follows from

$$b(z) \geq \frac{1}{a(z)} \tag{53}$$

which in turn follows from Jensen’s inequality, noting that  $s e^{-s^2/2} ds/2\pi$  is a probability measure.  $\square$

In Dolbeault *et al* (2007) it was proved that

$$\log |\mu_{\mathcal{L}}^-(\nu)| \approx -\frac{\pi}{2\nu} \tag{54}$$

as  $\nu \rightarrow 0_+$ . The methods in Dolbeault *et al* (2007) can be adapted to show that

$$\log |\mu_{\mathcal{L}}^+(\nu)| \approx -\frac{\pi}{2\nu} \tag{55}$$

as well, thus proving the following result.

**Lemma 5.** *With the above notations,  $\lim_{\nu \rightarrow 0_+} \nu \log |\mu_{\mathcal{L}}(\nu)| = -\frac{\pi}{2}$ .*

As  $\nu \rightarrow 0_+$ , this provides us with the leading asymptotics of the critical magnetic fields.

**Theorem 6.** *With the above notations,  $\lim_{\nu \rightarrow 0_+} \frac{\log B(\nu)}{\log B_{\mathcal{L}}(\nu)} = 1$ .*

**Remark .** This result does not prove that  $\frac{B(\nu)}{B_{\mathcal{L}}(\nu)}$  converges to some finite limit as  $\nu \rightarrow 0_+$ .

**Proof .** By (38), we already know that  $\mu(\nu) \leq \mu_{\mathcal{L}}(\nu)$ . To prove our result, we need an estimate of  $\mu(\nu)$  from below. Since for  $\nu$  small,  $B(\nu)$  is very large according to (Dolbeault *et al* 2007, corollary 11), we can assume that  $B(\nu) > 1$  for any  $\nu \in (0, \bar{\nu})$  for some  $\bar{\nu} > 0$ , so that, if  $\nu \in (0, \bar{\nu})$ , then  $\lambda_1(\nu, 1) > -1$  and therefore, for all  $\phi$ ,

$$\mathcal{E}_{1,\nu}[\phi] \geq \mathcal{F}_{\nu}[\phi] := \int_{\mathbb{R}^3} \frac{|\sigma \cdot \nabla_1 \phi|^2}{\lambda_1(\nu, 1) + 1 + \frac{\nu}{|x|}} d^3x - \int_{\mathbb{R}^3} \frac{\nu}{|x|} |\phi|^2 d^3x. \tag{56}$$

For completeness, we sketch the main steps of the proof, which is similar to that given in Dolbeault *et al* (2007). Since

$$\mathcal{G}_{\nu} \left( \begin{matrix} \phi \\ \chi \end{matrix} \right) := \left( H_B \left( \begin{matrix} \phi \\ \chi \end{matrix} \right), \begin{pmatrix} \phi \\ \chi \end{pmatrix} \right) \tag{57}$$

is concave in  $\chi$ ,

$$1 + \mathcal{F}_{\nu}[\phi] = \sup_{\chi} \frac{\mathcal{G}_{\nu} \left( \begin{matrix} \phi \\ \chi \end{matrix} \right)}{\|\phi\|_{L^2(\mathbb{R}^3)}^2 + \|\chi\|_{L^2(\mathbb{R}^3)}^2}. \tag{58}$$

Obviously,

$$\sup_{\chi} \frac{\mathcal{G}_{\nu} \left( \begin{matrix} \phi \\ \chi \end{matrix} \right)}{\|\phi\|_{L^2(\mathbb{R}^3)} + \|\chi\|_{L^2(\mathbb{R}^3)}} \geq \sup_{\Pi^{\perp} \chi = 0} \frac{\mathcal{G}_{\nu} \left( \begin{matrix} \phi \\ \chi \end{matrix} \right)}{\|\phi\|_{L^2(\mathbb{R}^3)} + \|\chi\|_{L^2(\mathbb{R}^3)}}. \tag{59}$$

This, by (Dolbeault *et al* 2007, proposition 14), is bounded below by

$$\sup_{\chi} \frac{\mathcal{G}_{\nu+\nu^{3/2}} \left( \begin{matrix} \Pi \phi \\ \Pi \chi \end{matrix} \right) + \mathcal{G}_{\nu+\sqrt{\nu}} \left( \begin{matrix} \Pi^{\perp} \phi \\ 0 \end{matrix} \right)}{\|\Pi \phi\|_{L^2(\mathbb{R}^3)} + \|\Pi^{\perp} \phi\|_{L^2(\mathbb{R}^3)} + \|\Pi \chi\|_{L^2(\mathbb{R}^3)}} \tag{60}$$




Figure 1. Plot of  $\lambda \mapsto \log[1 + g(\lambda, z_{\max})]$  with  $z_{\max} = 100$ , for  $\nu = 0.9$ .

Moreover, by (Dolbeault et al 2007, proposition 15), for  $\nu$  small enough, this is bounded below by

$$\sup_{\chi} \frac{\mathcal{G}_{\nu+\nu^{3/2}}\left(\frac{\Pi\phi}{\Pi\chi}\right) + d(\nu)\|\Pi^\perp\phi\|_{L^2(\mathbb{R}^3)}^2}{\|\Pi\phi\|_{L^2(\mathbb{R}^3)} + \|\Pi^\perp\phi\|_{L^2(\mathbb{R}^3)} + \|\Pi\chi\|_{L^2(\mathbb{R}^3)}}, \tag{61}$$

where  $d(\cdot)$  is a continuous function such that  $d(0) = \sqrt{2}$ . The inequality that leads to (61) displays the fact that being perpendicular to the lowest Landau level raises the energy.

Again by concavity, for all  $\phi$ , there is a unique  $\chi$  realizing

$$\sup_{\chi} \frac{\mathcal{G}_{\nu+\nu^{3/2}}\left(\frac{\Pi\phi}{\Pi\chi}\right)}{\|\Pi\phi\|_{L^2(\mathbb{R}^3)} + \|\Pi\chi\|_{L^2(\mathbb{R}^3)}} =: \lambda_{\nu+\nu^{3/2}}^-[\phi]. \tag{62}$$

Since  $\lambda_{\nu+\nu^{3/2}}^-[\phi] < 1 < \sqrt{2}$ , we finally have

$$1 + \mathcal{F}_\nu[\phi] \geq \lambda_{\nu+\nu^{3/2}}^-[\phi]. \tag{63}$$

By theorem 16 and corollary 17 of Dolbeault et al (2007),

$$\lambda_{\nu+\nu^{3/2}}^-[\phi] = 1 + \mu_{\mathcal{L}}^-(\nu + \nu^{3/2}) \tag{64}$$

and therefore, for  $\nu$  small enough, we have proved that

$$\mu_{\mathcal{L}}^-(\nu + \nu^{3/2}) \leq \mu(\nu) \leq \mu_{\mathcal{L}}(\nu), \tag{65}$$

where the upper estimate is given by inequality (38). Since  $\nu \rightarrow 0_+$ ,  $\nu^{3/2}$  becomes insignificant compared to  $\nu$ . The conclusion then holds by theorem 1.  $\square$

## 5. Numerical results


### 5.1. Computations in the Landau level ansatz

To compute  $\mu_{\mathcal{L}}$ , we minimize  $\mathcal{L}_\nu[f]\|f\|_{L^2(\mathbb{R}^+)}^2$  on the set of the solutions  $f_\lambda$  of


$$f'' + \frac{za(z)}{b(z)}f' + \frac{\nu}{b(z)}(\lambda + \nu a(z))f = 0, \quad f(0) = 1, \quad f'(0) = 0. \tag{66}$$

We note that  $b'(z) = za(z)$ , and, for any  $z > 0$ ,

$$a(z) = e^{\frac{z^2}{2}}\sqrt{\frac{\pi}{2}}\operatorname{erfc}\left(\frac{z}{\sqrt{2}}\right) \quad \text{and} \quad b(z) = e^{\frac{z^2}{2}}\sqrt{\frac{\pi}{2}}\operatorname{erfc}\left(\frac{z}{\sqrt{2}}\right) + z. \tag{67}$$



**Figure 2.** Left: values of the critical magnetic field in tesla ( $\log_{10}$  scale). Right: values in dimensionless units. Ground-state levels in the Landau level ansatz correspond to the upper curve, while those obtained without approximation are given by the lower curve. Dots correspond to the values computed by Schlüter *et al* (1985) in the Landau level ansatz.



**Figure 3.** Ratio of the ground-state levels computed in the Landau level ansatz versus ground-state levels obtained by a computation without any symmetry ansatz.


**Table 1.** Numerical values found in the Landau level ansatz.

$\nu$	$Z$	$\mu_{\mathcal{L}}$	$B_{\mathcal{L}}(\nu)$	$\log_{10}(bB_{\mathcal{L}}(\nu))$
0.409	56	-0.046 1591	1877.35	12.9184
0.5	68.52	-0.088 7408	507.941	12.3506
0.598	82	-0.145 25	189.596	11.9227
0.671	92	-0.192 837	107.567	11.6765
0.9	123.33	-0.363 773	30.2274	11.1252
1	137.037	-0.445 997	20.1093	10.9482

Numerically, we use a shooting method and minimize  $g(\lambda, z_{\max}) := |f_{\lambda}(z_{\max})|^2 + |f'_{\lambda}(z_{\max})|^2$  for some  $z_{\max}$  large enough. As  $z_{\max} \rightarrow \infty$ , the first minimum  $\mu_{\mathcal{L}}(\nu, z_{\max})$  of  $\lambda \mapsto g(\lambda, z_{\max})$  converges to 0 and thus determines  $\lambda = \mu_{\mathcal{L}}(\nu)$ . See figure 1.

Let  $b = \frac{m^2 c^2}{e \hbar} \approx 4.414 \cdot 10^9$  be the numerical factor to obtain the critical field in tesla. Corresponding values are given in  $\log_{10}$  scale. The minimum  $\mu_{\mathcal{L}}(\nu, z_{\max})$  is found by dichotomy. Results computed with Mathematica are given in table 1.

There is no significant difference with the results that were found in Schlüter *et al* (1985). See figure 2.



**Figure 4.** Plot of  $\log_{10}(10^{-4} + |\phi|^4)$  for  $\nu = 0.5$  (top) and  $\nu = 0.9$  (bottom). Left: Landau level ansatz. Right: computation without any symmetry constraint. Level lines are equidistributed.

**Table 2.** Results of the minimization method without symmetry ansatz.

$\nu$	$Z$	$\lambda_1$	$B(\nu)$	$\log_{10}(bB(\nu))$
0.50	68.5185	-0.087 4214	523.389	12.3637
0.55	75.3704	-0.119 458	280.305	12.0925
0.60	82.2222	-0.153 882	168.922	11.8725
0.65	89.0741	-0.191 037	109.604	11.6847
0.70	95.9259	-0.231 198	74.833	11.5189
0.75	102.778	-0.274 665	53.0216	11.3693
0.80	109.63	-0.321 875	38.6087	11.2315
0.85	116.481	-0.373 535	28.668	11.1022
0.90	123.333	-0.430 854	21.5476	10.9782
0.95	130.185	-0.496 005	16.2588	10.8559
1.00	137.037	-0.573 221	12.1735	10.7302

### 5.2. Computations in the unconstrained case

We numerically compute  $B(\nu)$  in the general case, without ansatz. For this purpose, we discretize the minimization problem defining  $\mu(\nu)$  using B-spline functions of degree 1. They are defined on a logarithmic, variable step-size grid, in order to capture the behavior of the eigenfunctions near the singularity. We also use cylindrical symmetry to lower the dimension from 3 to 2. These two choices provide us with very sparse matrices, even if large. We use Matlab routines to calculate all integrals and the eigenvalues of the corresponding discretized matrices. The size of the computing domain is adapted as  $\nu$  varies. Results of these computations are shown in table 2.

### 5.3. Discussion

When dealing with the physics of magnetars, one is interested only in the order of magnitude of the critical magnetic field. With this goal in mind, the values computed in the Landau level ansatz are quite satisfactory. The corresponding values are given in the right column of table 2,  $\log_{10}(bB(\nu))$ , where  $b$  is approximately  $4.414 \times 10^9$  tesla. See figure 2(left).

Except maybe in the limit  $\nu \rightarrow 0$ , it is however clear from theorem 1 that the equivalent eigenvalue problem (9) has nothing to do with its counterpart  $\mu_{\mathcal{L}}(\nu)$  in the Landau level ansatz. What our computations show is that the values of the computed critical fields significantly differ, see figure 3, and that the shapes of the corresponding ground state do not have much in common, see figure 4.

## 6. Conclusion

Orders of magnitude of the critical magnetic field given by the Landau level ansatz are similar to those obtained without constraint. Qualitatively, the curve of the critical magnetic field in terms of the charge, is also well reproduced by the Landau level ansatz.

However, the difference of the values of the critical magnetic fields is not small when comparison is done for values of  $\nu$  approaching 1. It turns out that values given by the Landau level ansatz are 50% higher than those found by a minimization approach without symmetry ansatz. Shapes of the density distributions also differ significantly. They are much more peaked around the singularity when computed unconstrained than in the Landau level ansatz.

The Landau level ansatz, which is commonly accepted in non-relativistic quantum mechanics as a good approximation for large magnetic fields, is a quite crude approximation for the computation of the critical magnetic field (that is the strength of the field at which the lowest eigenvalue in the gap reaches its lower end) in the Dirac–Coulomb model. Even for small values of  $\nu$ , which were out of reach in our numerical study, it is not clear that the Landau level ansatz gives the correct approximation at first order in terms of  $\nu$ . Hence, accurate numerical computations involving the Dirac equation cannot simply rely on the Landau level ansatz.

## Acknowledgments

JD and MJE acknowledge support from ANR Accquarel project. ML is partially supported by US National Science Foundation grant DMS 06-00037. The authors thank Joachim Reinhardt for pointing out Schlüter *et al* (1985) and the two referees for suggesting references and an interesting comment.

## References

- Baumgartner B and Seiringer R 2001 *Ann. Henri Poincaré* **2** 41–76
- Baumgartner B, Solovej J P and Yngvason J 2000 *Commun. Math. Phys.* **212** 703–24
- Brummelhuis R and Duclos P 2006 *J. Math. Phys.* **47** 032103, 41
- Dolbeault J, Esteban M J and Loss M 2007 *Ann. Henri Poincaré* **8** 749–79
- Duncan R C 2000 Physics in ultra-strong magnetic fields *Review talk given at 5th Huntsville Gamma-Ray Burst Symposium*
- Fushiki I, Gudmundsson E H, Pethick C J and Yngvason J 1992 *Ann. Phys.* **216** 29–72
- Kouveliotou C, Strohmayer T, Hurley K, van Paradijs J, Finger M H, Dieters S, Woods P, Thompson C and Duncan R C 1999 *Astrophys. J.* **510** L115–8

Lieb E H, Solovej J P and Yngvason J 1994a *Comm. Pure Appl. Math.* **47** 513–91

Lieb E H, Solovej J P and Yngvason J 1994b *Commun. Math. Phys.* **161** 77–124

Medin Z and Lai D 2006 *Phys. Rev.* **74** 062507

Pickl P 2005 Existence of spontaneous pair creation PhD Thesis Mathematisches Institut der Ludwig-Maximilians-Universität München

Schlüter P, Soff G, Wietschorke K H and Greiner W 1985 *J. Phys. B: At. Mol. Phys.* **18** 1685–700

von Milczewski J, Diercksen G H F and Uzer T 1996 *Phys. Rev. Lett.* **76** 2890–3