

Dispersionless limit of the noncommutative potential KP hierarchy and solutions of the pseudodual chiral model in $2 + 1$ dimensions

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys. A: Math. Theor. 41 265205

(<http://iopscience.iop.org/1751-8121/41/26/265205>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.149

The article was downloaded on 03/06/2010 at 06:56

Please note that [terms and conditions apply](#).

Dispersionless limit of the noncommutative potential KP hierarchy and solutions of the pseudodual chiral model in 2 + 1 dimensions

Aristophanes Dimakis¹ and Folkert Müller-Hoissen²

¹ Department of Financial and Management Engineering, University of the Aegean, 31 Fostini Str., GR-82100 Chios, Greece

² Max-Planck-Institute for Dynamics and Self-Organization, Bunsenstrasse 10, D-37073 Göttingen, Germany

E-mail: dimakis@aegean.gr and folkert.mueller-hoissen@ds.mpg.de

Received 17 March 2008

Published 11 June 2008

Online at stacks.iop.org/JPhysA/41/265205

Abstract

The usual dispersionless limit of the KP hierarchy does not work in the case where the dependent variable has values in a noncommutative (e.g. matrix) algebra. Passing over to the potential KP hierarchy, there is a corresponding scaling limit in the noncommutative case, which turns out to be the hierarchy of a ‘pseudodual chiral model’ in 2 + 1 dimensions (‘pseudodual’ to a hierarchy extending Ward’s (modified) integrable chiral model). Applying the scaling procedure to a method generating exact solutions of a matrix (potential) KP hierarchy from solutions of a matrix linear heat hierarchy, leads to a corresponding method that generates exact solutions of the matrix dispersionless potential KP hierarchy, i.e. the pseudodual chiral model hierarchy. We use this result to construct classes of exact solutions of the $su(m)$ pseudodual chiral model in 2 + 1 dimensions, including various multiple lump configurations.

PACS numbers: 02.30.Ik, 05.45.Yv

Mathematics Subject Classification: 35Q58, 37K10, 35Q51

(Some figures in this article are in colour only in the electronic version)

1. Introduction

Expressing the scalar KP hierarchy with the dependent variable $u(t_1, t_2, \dots)$ in terms of new evolution variables $T_n = \epsilon t_n$ with a parameter ϵ , the limit $\epsilon \rightarrow 0$ (keeping T_n fixed) leads to the so-called dispersionless KP hierarchy (see [1–23], for example). The same limit does not work, however, for the KP hierarchy with the dependent variable in a noncommutative (e.g.

matrix) algebra. In fact, different scaling limits of the matrix KP equation have already been explored in [24], where the multiscale expansion method has been used to relate different integrable systems.

In the present work, we formulate a dispersionless limit of the ‘noncommutative’ potential KP (ncpKP) hierarchy with dependent variable ϕ , where $u = \phi_{t_1}$. It turns out to be the hierarchy associated with a ‘pseudodual chiral model’ (pdCM) in $2 + 1$ dimensions, a well-known reduction of the self-dual Yang–Mills equation [25, 26]. Applying the scaling limit procedure to a method generating exact solutions of a matrix pKP hierarchy from solutions of a matrix linear heat hierarchy, then results in a method generating solutions of this pdCM hierarchy.

In section 2 we consider the dispersionless limit of the ncpKP equation. Section 3 generalizes this limit to the whole ncpKP hierarchy, explores some of its properties, and in particular establishes a pseudoduality relation with a hierarchy that extends Ward’s (modified) chiral model in $2 + 1$ dimensions [27–32]. The latter model has been studied extensively [33–52] (see also [53–55] for the Ward model in (anti-)de Sitter spacetime and [56–63] for explorations of a Moyal-deformed version), in particular concerning its (multi-)lump solutions, which are two-dimensional soliton-like objects. In this respect, its pseudodual received comparatively little attention. The dependent variables of the two equations are related by a kind of hetero-Bäcklund transformation. Given a solution of one of the two equation, this becomes a first-order system of partial differential equations, which determines a solution of the other equation. The necessary integration is typically difficult to carry out, however. Hence, although some properties of the pseudodual model can certainly be inferred from corresponding knowledge of the Ward model, there is no *explicit* translation of its solutions. In any case, in this work we present an independent approach to solutions of the pdCM and moreover to its hierarchy.

In section 4 we derive the abovementioned method to generate exact solutions of the pdCM hierarchy from the corresponding knowledge of the ncpKP hierarchy. The main result is independently verified in section 5 and then applied to construct some classes of exact solutions. This section is actually formulated in such a way that it can be accessed almost without any knowledge of the previous sections. We concentrate on solutions of the $su(m)$ pdCM hierarchy and restrict concrete examples to the $su(2)$ case. Some conclusions are collected in section 6.

2. The dispersionless limit of the noncommutative pKP equation

Let $\phi(\mathbf{t})$ with $\mathbf{t} = (t_1, t_2, \dots)$ be a function with values in some matrix space³ \mathcal{A} which is endowed with a product $A \cdot B = AQB$, where Q is a constant matrix, i.e. independent of \mathbf{t} . We consider the following ncpKP equation,

$$4\phi_{tx} - \phi_{xxxx} - 3\phi_{yy} = 6(\phi_x Q \phi_x)_x - 6[\phi_x, \phi_y]_Q, \tag{2.1}$$

where $x = t_1, y = t_2, t = t_3$ and

$$[A, B]_Q := AQB - BQA. \tag{2.2}$$

Let ϕ now also depend on a parameter ϵ in such a way that

$$\phi(\mathbf{t}, \epsilon) = \epsilon^a \Phi(\mathbf{T}) + \mathcal{O}(\epsilon^{a+1}) \tag{2.3}$$

³ The entries will be taken as complex functions of t_1, t_2, \dots , though large parts of this work also apply to the case where they are elements of any (possibly noncommutative) associative algebra, for which differentiability with respect to t_1, t_2, \dots can be defined.

with some integer a . Furthermore, we assume that Q has an expansion

$$Q = Q_{(0)} + \epsilon Q_{(1)} + \mathcal{O}(\epsilon^2). \tag{2.4}$$

Rewriting the ncpKP equation in terms of the rescaled variables $T_n = \epsilon t_n$, dividing the equation by the maximal power of ϵ common to all of its summands, and taking the limit $\epsilon \rightarrow 0$ while keeping T_1, T_2, \dots fixed, should result in an equation that still has linear as well as nonlinear terms (in Φ). This fixes the value of a , but we have to distinguish the following two cases.

If the algebra (\mathcal{A}, \cdot) is *commutative* at $\epsilon = 0$ with $Q_{(0)} \neq 0$, and hence the commutator $[\Phi_X, \Phi_Y]_{Q_{(0)}} = \Phi_X Q_{(0)} \Phi_Y - \Phi_Y Q_{(0)} \Phi_X$ vanishes, then our requirements lead to $a = -1$, and the scaling limit of the pKP equation, divided by ϵ , is

$$4\Phi_{TX} - 3\Phi_{YY} = 6(\Phi_X Q_{(0)} \Phi_X)_X - 6[\Phi_X, \Phi_Y]_{Q_{(1)}}, \tag{2.5}$$

where $X = T_1, Y = T_2, T = T_3$. If Φ is a scalar and $Q_{(0)} = 1$, the last equation reduces to

$$4\Phi_{TX} - 3\Phi_{YY} = 6(\Phi_X^2)_X. \tag{2.6}$$

This is the potential form of the dispersionless limit of the ('commutative') scalar KP equation, which is also known as the Khokhlov–Zabolotskaya equation (see [4] for instance).

If the algebra (\mathcal{A}, \cdot) is *noncommutative* at $\epsilon = 0$, we have to set

$$a = 0, \tag{2.7}$$

and this choice will be made throughout this work. Then we obtain the following *dispersionless limit of the ncpKP equation (2.1)*:

$$4\Phi_{TX} - 3\Phi_{YY} = -6[\Phi_X, \Phi_Y]_{Q_{(0)}}. \tag{2.8}$$

Up to the modified matrix product and rescalings of the coordinates, this is a well-known reduction of the self-dual Yang–Mills equation (see [25, 26, 64–68]). With the further dimensional reduction $\Phi_X = \Phi_T$, it becomes the *pseudodual chiral model* [69–71] (see also [64, 72–74]). Accordingly, we may call (2.8) a *pseudodual chiral model in 2 + 1 dimensions*, in the following abbreviated to *pdCM*. In fact, as explained in section 3.2, it is 'pseudodual' to an integrable (modified) chiral model in 2 + 1 dimensions.

3. The dispersionless limit of the ncpKP hierarchy

A functional representation of the ncpKP hierarchy is given by [75]

$$(\phi - \phi_{-[\lambda]})(\lambda^{-1} - Q\phi) - \phi_{t_1} = \theta - \theta_{-[\lambda]}, \tag{3.1}$$

where θ is an arbitrary \mathcal{A} -valued function, and $(\phi_{-[\lambda]})(\mathbf{t}) := \phi(\mathbf{t} - [\lambda])$ is a Miwa shift with $[\lambda] = (\lambda, \lambda^2/2, \lambda^3/3, \dots)$, λ an indeterminate. Eliminating θ from this equation, we get the following functional form of the ncpKP hierarchy:

$$\begin{aligned} & ((\phi - \phi_{-[\lambda]})(\lambda^{-1} - Q\phi) - \phi_{t_1}) - ((\phi - \phi_{-[\lambda]})(\lambda^{-1} - Q\phi) - \phi_{t_1})_{-[\mu]} \\ &= ((\phi - \phi_{-[\mu]})(\mu^{-1} - Q\phi) - \phi_{t_1}) - ((\phi - \phi_{-[\mu]})(\mu^{-1} - Q\phi) - \phi_{t_1})_{-[\lambda]}, \end{aligned} \tag{3.2}$$

where μ is another indeterminate.

If $\mathbf{p}_n, n = 1, 2, \dots$, denote the elementary Schur polynomials, then

$$\mathbf{p}_n(-\tilde{\partial}) = -\frac{\epsilon}{n} \partial_{T_n} + \mathcal{O}(\epsilon^2), \tag{3.3}$$

where $\tilde{\partial} = (\partial_{t_1}, \partial_{t_2}/2, \partial_{t_3}/3, \dots)$, and hence

$$\phi - \phi_{-[\lambda]} = \epsilon D(\lambda)\Phi + \mathcal{O}(\epsilon^2), \tag{3.4}$$

where

$$D(\lambda) := \sum_{n \geq 1} \frac{\lambda^n}{n} \partial_{T_n}. \quad (3.5)$$

In accordance with (2.3), where now $a = 0$, we shall assume

$$\theta(\mathbf{t}, \epsilon) = \Theta(\mathbf{T}) + \mathcal{O}(\epsilon). \quad (3.6)$$

Then we obtain

$$D(\lambda)(\Phi) (\lambda^{-1} - Q_{(0)}\Phi) - \Phi_{T_1} = D(\lambda)(\Theta). \quad (3.7)$$

Expanding this in powers of λ , we find

$$\frac{1}{n+1} \Phi_{T_{n+1}} - \frac{1}{n} \Phi_{T_n} Q_{(0)}\Phi = \frac{1}{n} \Theta_{T_n}, \quad n = 1, 2, \dots \quad (3.8)$$

Elimination of Θ results in the hierarchy equations

$$\frac{n}{(n+1)} \Phi_{T_{n+1}, T_m} - \frac{m}{(m+1)} \Phi_{T_{m+1}, T_n} = \Phi_{T_n} Q_{(0)}\Phi_{T_m} - \Phi_{T_m} Q_{(0)}\Phi_{T_n}. \quad (3.9)$$

Introducing

$$x_n := nT_n, \quad n = 1, 2, \dots, \quad (3.10)$$

this becomes

$$\Phi_{x_{n+1}, x_m} - \Phi_{x_{m+1}, x_n} = [\Phi_{x_n}, \Phi_{x_m}]_{Q_{(0)}}, \quad m, n = 1, 2, \dots \quad (3.11)$$

For $m = 1, n = 2$, we recover (2.8).

Expressing $Q_{(0)}$ as

$$Q_{(0)} = VU^\dagger \quad (3.12)$$

with matrices U, V and the adjoint (complex conjugate and transpose) U^\dagger of U , then

$$\varphi := U^\dagger \Phi V \quad (3.13)$$

(which includes the cases $\varphi = Q_{(0)}\Phi$ and $\varphi = \Phi Q_{(0)}$) solves

$$\varphi_{x_{n+1}, x_m} - \varphi_{x_{m+1}, x_n} = [\varphi_{x_n}, \varphi_{x_m}], \quad m, n = 1, 2, \dots, \quad (3.14)$$

if Φ solves (3.11). The power of this observation lies in the fact that any solution of (3.11) in some $M \times N$ matrix algebra, where $Q_{(0)} = VU^\dagger$ with an $M \times m$ matrix U and an $N \times m$ matrix V , determines in this way a solution of (3.14) in the $m \times m$ matrix algebra. For example, if we are looking for solutions of (3.14) in the algebra of 2×2 matrices, we may first look for solutions of (3.11) with any $M, N \geq 2$ and $Q_{(0)} = VU^\dagger$ with $M \times 2$ and $N \times 2$ matrices U and V . In this way (simple) solutions of (3.11) in arbitrarily large matrix algebras lead to (complicated) solutions of (3.14) in the algebra of 2×2 matrices. In particular, this explains the significance of $Q_{(0)}$ in our previous formulae. In section 5 we will substantiate this method. The hierarchy (3.14) is consistent with restricting φ to take values in any Lie algebra, e.g. $sl(N, \mathbb{R}), sl(N, \mathbb{C}), u(N)$ or $su(N)$. If φ solves (3.14), then also $\varphi + \varphi_0$, where φ_0 is a constant in the respective Lie algebra.

As a consequence of their origin, the hierarchies (3.11) and (3.14) are invariant under the scaling transformation $x_n \mapsto \lambda x_n, n = 1, 2, \dots$, with any constant $\lambda \neq 0$.

Remark. If g_1, g_2 are any two constant invertible matrices with size such that $g_1 \Phi g_2$ is defined, then

$$\Phi \mapsto g_1 \Phi g_2, \quad Q_{(0)} \mapsto g_2^{-1} Q_{(0)} g_1^{-1} \quad (3.15)$$

leaves (3.11) invariant. If $Q_{(0)}$ is given by (3.12), the latter transformation results from

$$V \mapsto g_2^{-1}V, \quad U \mapsto (g_1^\dagger)^{-1}U, \quad (3.16)$$

and φ is invariant. More generally, the transformation $V \mapsto g_2^{-1}V\sigma, U \mapsto (g_1^\dagger)^{-1}U(\sigma^\dagger)^{-1}$, with a constant $m \times m$ matrix σ , leads to $\varphi \mapsto \sigma^{-1}\varphi\sigma$. This leaves the hierarchy equations (3.14) invariant.

3.1. Some properties of the first dispersionless hierarchy equation

A Lagrangian for the first equation ($m = 1, n = 2$)

$$\varphi_{x_1, x_3} - \varphi_{x_2, x_2} = -[\varphi_{x_1}, \varphi_{x_2}] \quad (3.17)$$

of the hierarchy (3.14) is

$$\mathcal{L} = -\text{tr}(\varphi_{x_1}\varphi_{x_3} - \varphi_{x_2}^2 - \frac{2}{3}\varphi[\varphi_{x_1}, \varphi_{x_2}]) \quad (3.18)$$

(see also [64, 65]). After passage to the new coordinates x, y, t given by

$$x_1 = \frac{1}{2}(t - x), \quad x_2 = y, \quad x_3 = \frac{1}{2}(t + x), \quad (3.19)$$

equation (3.17) becomes

$$\varphi_{tt} - \varphi_{xx} - \varphi_{yy} + [\varphi_t - \varphi_x, \varphi_y] = 0, \quad (3.20)$$

and the Lagrangian takes the form

$$\begin{aligned} \mathcal{L} &= -\frac{1}{2} \text{tr}(\varphi_t^2 - \varphi_x^2 - \varphi_y^2 - \frac{2}{3}\varphi[\varphi_t - \varphi_x, \varphi_y]) \\ &= -\frac{1}{2} \text{tr}(\eta^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi + \frac{2}{3} \varphi v_\rho \epsilon^{\rho\mu\nu} \partial_\mu \varphi \partial_\nu \varphi), \end{aligned} \quad (3.21)$$

where we introduced the components $\eta^{\mu\nu}$ (with respect to the coordinates $(x^\mu) = (t, x, y)$) of the Minkowski metric in 2+1 dimensions, the totally antisymmetric Levi-Civita pseudo-tensor with $\epsilon^{012} = 1$, and a constant covector v_ρ with components $(1, 1, 0)$. As a consequence of the translational invariance of the Lagrangian, the energy–momentum tensor

$$T^\mu{}_\nu = \text{tr} \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu \varphi)} \partial_\nu \varphi - \delta_\nu^\mu \mathcal{L} \right) \quad (3.22)$$

provides us with the conserved densities

$$\begin{aligned} T^0{}_0 &= -\frac{1}{2} \text{tr}(\varphi_t^2 + \varphi_x^2 + \varphi_y^2 - \frac{2}{3}\varphi[\varphi_x, \varphi_y]), \\ T^0{}_1 &= -\text{tr}(\varphi_t \varphi_x - \frac{1}{3}\varphi[\varphi_x, \varphi_y]), \quad T^0{}_2 = -\text{tr}(\varphi_t \varphi_y). \end{aligned} \quad (3.23)$$

Then also

$$\mathcal{E} = T^0{}_0 - T^0{}_1 = -\frac{1}{2} \text{tr}[(\varphi_t - \varphi_x)^2 + \varphi_y^2] \quad (3.24)$$

is a conserved density. For any nonzero *anti-Hermitian* matrix, the trace of the square of the matrix is real and negative. Hence \mathcal{E} provides us with a *non-negative* ‘energy’ density in the case where φ takes values in the Lie algebra $u(m)$ of the unitary group.

For any infinitesimal symmetry $\delta\varphi = \frac{\partial\varphi}{\partial\alpha}\delta\alpha$ (with a parameter α) of the Lagrangian, there is a conserved current

$$J^\mu := \text{tr} \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu \varphi)} \frac{\partial \varphi}{\partial \alpha} \right), \quad (3.25)$$

i.e., $\partial_\mu J^\mu = 0$. A symmetry of the above Lagrangian is given by $\delta\varphi = [C, \varphi]\alpha$ with any constant (anti-Hermitian) matrix C . Hence

$$J_C^0 = -\text{tr} \left(([\varphi, \varphi_t] + \frac{1}{3}(\varphi^2 \varphi_y - 2\varphi \varphi_y \varphi + \varphi_y \varphi^2)) C \right) \quad (3.26)$$

is a conserved density.

3.2. Relation with Ward's chiral model in 2 + 1 dimensions

The hierarchy (3.14) is related to the hierarchy of an integrable (modified) chiral model in 2 + 1 dimensions. First we note that (3.14) is the integrability condition of the linear system

$$J_{x_{n+1}} = -J\varphi_{x_n}, \quad n = 1, 2, \dots \quad (3.27)$$

with some invertible J . Rewriting this as

$$\varphi_{x_n} = -J^{-1}J_{x_{n+1}}, \quad n = 1, 2, \dots, \quad (3.28)$$

we find that (3.14) is automatically satisfied and the integrability conditions now take the form

$$\left(J^{-1}J_{x_{n+1}}\right)_{x_m} - \left(J^{-1}J_{x_{m+1}}\right)_{x_n} = 0, \quad m, n = 1, 2, \dots \quad (3.29)$$

In conclusion, solutions J of (3.29) are in correspondence with solutions φ of (3.14) via (3.28). This correspondence is of a nonlocal nature. In particular, given a solution φ of (3.14), (3.28) does not directly determine $J^{-1}J_{x_1}$. We first have to solve (3.27) for J in order to be able to calculate this expression.

Equation (3.29) is immediately recognized as the dispersionless limit of the noncommutative *modified* KP hierarchy (see equation (4.12) in [76]).

For $m = 1$ and $n = 2$, (3.29) reads

$$\left(J^{-1}J_{x_3}\right)_{x_1} - \left(J^{-1}J_{x_2}\right)_{x_2} = 0. \quad (3.30)$$

This equation apparently first appeared in [77, 78]. It is a reduction of the self-dual Yang–Mills equation (see [77, 65, 26], for example). In terms of the coordinates x, y, t given by (3.19), it takes the form

$$\left(J^{-1}J_t\right)_t - \left(J^{-1}J_x\right)_x - \left(J^{-1}J_y\right)_y + \left[J^{-1}J_x, J^{-1}J_t\right] = 0, \quad (3.31)$$

or in tensor notation (using the summation convention)

$$\left(\eta^{\mu\nu} + \epsilon^{\mu\nu}\right)\partial_\mu\left(J^{-1}\partial_\nu J\right) = 0, \quad (3.32)$$

where $\mu, \nu = 0, 1, 2$, $\left(\eta^{\mu\nu}\right) = \text{diag}(1, -1, -1)$, and $\epsilon^{\mu\nu}$ is antisymmetric with $\epsilon^{01} = -\epsilon^{10} = 1$ and zero otherwise. We note that the bivector $\epsilon^{\mu\nu}$ breaks Lorentz invariance in 2 + 1 dimensions. Using the Lorentz invariant Levi-Civita pseudo-tensor and the constant unit covector v_α with components $(0, 0, 1)$, it can be expressed as $\epsilon^{\mu\nu} = v_\alpha\epsilon^{\alpha\mu\nu}$. Another integrable equation is obtained if we choose v_α to be timelike [27, 38, 78]. Equation (3.32) is Ward's (2+1)-dimensional generalization of the chiral (or sigma) model [27–32], see also [33–48, 50–52, 68]. J can be consistently restricted to any Lie group, e.g. $SL(N, \mathbb{R})$, $SL(N, \mathbb{C})$, $U(N)$ or $SU(N)$.

Remark. According to (3.28) we have $J^{-1}J_y = \varphi_x - \varphi_t$ and $J^{-1}J_t + J^{-1}J_x = -\varphi_y$, in terms of the variables x, y, t given by (3.19). Hence

$$\mathcal{E} = \mathcal{E}_{\text{Ward}} - \text{tr}\left(J^{-1}J_t J^{-1}J_x\right), \quad (3.33)$$

where

$$\mathcal{E}_{\text{Ward}} = -\frac{1}{2}\text{tr}\left(\left(J^{-1}J_t\right)^2 + \left(J^{-1}J_x\right)^2 + \left(J^{-1}J_y\right)^2\right) \quad (3.34)$$

is the energy density of Ward's chiral model. The difference between $\mathcal{E}_{\text{Ward}}$ and \mathcal{E} is *not* a local expression in terms of φ . The appendix attempts to further clarify the relation between Ward's chiral model and the pdCM hierarchy (and yet another version of it).

3.3. An associated bidifferential calculus

On the algebra \mathcal{A} of $m \times m$ matrices with entries depending smoothly on x_1, x_2, \dots , we introduce two linear maps d, \bar{d} by⁴

$$d\psi = \sum_{n \geq 1} \psi_{x_n} dx_n, \quad \bar{d}\psi = \sum_{n \geq 1} \psi_{x_{n+1}} dx_n. \quad (3.35)$$

By use of the graded Leibniz rule they extend to a (bi-)differential graded algebra and satisfy

$$d^2 = \bar{d}^2 = d\bar{d} + \bar{d}d = 0, \quad (3.36)$$

and hence we have a *bidifferential calculus*. Dressing \bar{d} by setting

$$\bar{D}\psi = \bar{d}\psi - A\psi, \quad (3.37)$$

with a 1-form $A = \sum_{n \geq 1} A_{n+1} dx_n$, we find that d, \bar{D} yields again a bidifferential calculus ($\bar{D}^2 = d\bar{D} + \bar{D}d = 0$), iff

$$dA = 0, \quad \bar{d}A = A \wedge A \quad (3.38)$$

(see also [81]). These equations cover Ward’s chiral model hierarchy as well as its pseudodual, which is the dispersionless ncpKP hierarchy. Indeed, solving the first equation by setting

$$A = d\varphi, \quad (3.39)$$

the second reproduces the pdCM hierarchy

$$\bar{d}d\varphi = d\varphi \wedge d\varphi. \quad (3.40)$$

Alternatively, solving the second of equations (3.38) by setting

$$A = -J^{-1}\bar{d}J, \quad (3.41)$$

we recover the hierarchy

$$d(J^{-1}\bar{d}J) = 0 \quad (3.42)$$

associated with Ward’s chiral model. The relation between both hierarchies is given by

$$J^{-1}\bar{d}J = -d\varphi \quad (3.43)$$

(which is (3.28)). This may be regarded as a ‘Miura transformation’. The linear system associated with the bidifferential calculus is

$$\bar{D}\psi - \lambda d\psi = 0, \quad (3.44)$$

with a parameter λ . Taking components of the differential forms, this reads

$$(\partial_{x_{n+1}} - A_{n+1} - \lambda \partial_{x_n})\psi = 0 \quad n = 1, 2, \dots \quad (3.45)$$

The integrability conditions now have the form

$$[\partial_{x_{n+1}} - A_{n+1} - \lambda \partial_{x_n}, \partial_{x_{m+1}} - A_{m+1} - \lambda \partial_{x_m}] = 0. \quad (3.46)$$

Its multicomponent version (and with $m, n \in \mathbb{Z}$) appeared in [82] (see (2.1), (2.2), and also the references therein).

Nonlocal conserved currents are obtained in the following way [81]. Let $d\chi_0 = 0$. As a consequence of the bidifferential calculus structure, there are $\chi_n, n = 1, 2, \dots$, such that

$$j_{n+1} := \bar{D}\chi_n = -d\chi_{n+1} \quad n = 0, 1, \dots \quad (3.47)$$

⁴ We note that $\bar{d} = \mathcal{R} \circ d$ where \mathcal{R} is the linear left \mathcal{A} -module map determined by $\mathcal{R}(dx_n) = dx_{n-1}$ for $n > 1$, and $\mathcal{R}(dx_1) = 0$. This makes contact with Frölicher–Nijenhuis theory [79], see also [80].

iteratively determines $\chi_n, n = 1, 2, \dots$. For example, starting with $\chi_0 = \mathcal{I}$ (the unit matrix), we get $j_1 = \bar{D}\mathcal{I} = -d\varphi$ (using (3.39)), hence $\chi_1 = \varphi + a$ with $da = 0$, and thus also $\bar{d}a = 0$. In the second step we have $j_2 = \bar{D}(\varphi + a) = \bar{d}\varphi - d\varphi(\varphi + a)$, and the construction of the next current requires the integration of $d\chi_2 = d\varphi(\varphi + a) - \bar{d}\varphi$. The constant a actually turns out to be redundant and should be set to zero.

A Bäcklund transformation is obtained from

$$(d - \lambda^{-1}\bar{D})(\mathcal{I} + \lambda^{-1}\mathcal{B}) = (\mathcal{I} + \lambda^{-1}\mathcal{B})(d - \lambda^{-1}\bar{D}'), \quad \bar{D}' = \bar{d} - A', \quad (3.48)$$

with an operator \mathcal{B} (see [83]). Expanding in powers of λ^{-1} , we find

$$[d, \mathcal{B}] = \bar{D} - \bar{D}', \quad \bar{D}\mathcal{B} = \mathcal{B}\bar{D}'. \quad (3.49)$$

Assuming $\mathcal{B}(\psi) = B\psi$ with a matrix B , this means

$$d(B) = A' - A, \quad \bar{d}(B) = AB - BA'. \quad (3.50)$$

Using (3.39) and solving the first of these equations by setting $B = \varphi' - \varphi - a$ with $da = 0$, we obtain from the second

$$\bar{d}(\varphi' - \varphi) = d\varphi(\varphi' - \varphi - a) - (\varphi' - \varphi - a)d\varphi', \quad (3.51)$$

a Bäcklund transformation of the pdCM hierarchy. Alternatively, using (3.41) and solving the second of equations (3.50) by setting $B = -J^{-1}\mathcal{K}J'$ with $\bar{d}\mathcal{K} = 0$, the first becomes

$$J'^{-1}\bar{d}J' - J^{-1}\bar{d}J = d(J^{-1}\mathcal{K}J'), \quad (3.52)$$

a Bäcklund transformation of the (modified) chiral model hierarchy (see also [84] for the case of the chiral model on a two-dimensional spacetime).

If $\mathcal{B}_{i,j}$ leads from an i th to a j th solution, a permutability relation is given by

$$B_{12} + B_{24} = B_{13} + B_{34}, \quad B_{12}B_{24} = B_{13}B_{34} \quad (3.53)$$

(see [83]). This determines algebraically a forth solution from a given (first) solution and two Bäcklund descendants of it (with different parameters).

4. Towards exact solutions of the dispersionless ncpKP hierarchy

In this section, we start with a result that determines a large class of exact solutions of an ncpKP hierarchy and use the scaling limit towards the dispersionless hierarchy in order to obtain from it a corresponding result that determines exact solutions of the latter, which is a pdCM hierarchy. Let us recall theorem 4.1 from [75].

Theorem 4.1. *Let (\mathcal{A}, \cdot) be the algebra of $M \times N$ matrices of functions of \mathfrak{t} with the product*

$$A \cdot B = AQB, \quad (4.1)$$

where the ordinary matrix product is used on the right-hand side, and Q is a constant $N \times M$ matrix. Let $\tilde{\mathcal{X}}$ be an invertible $N \times N$ matrix and $\tilde{\mathcal{Y}} \in \mathcal{A}$, such that $\tilde{\mathcal{X}}, \tilde{\mathcal{Y}}$ solve the linear heat hierarchy (i.e. $\partial_{t_n}(\tilde{\mathcal{X}}) = \partial_{t_1}^n(\tilde{\mathcal{X}})$, $n = 2, 3, \dots$, and correspondingly for $\tilde{\mathcal{Y}}$) and satisfy

$$\tilde{\mathcal{X}}_{t_1} = R\tilde{\mathcal{X}} + Q\tilde{\mathcal{Y}}, \quad (4.2)$$

with a constant $N \times N$ matrix R . The pKP hierarchy in (\mathcal{A}, \cdot) is then solved by

$$\phi := \tilde{\mathcal{Y}}\tilde{\mathcal{X}}^{-1}. \quad (4.3)$$

A functional representation of the heat hierarchy condition is

$$\lambda^{-1}(\tilde{\mathcal{X}} - \tilde{\mathcal{X}}_{-[\lambda]}) = \tilde{\mathcal{X}}_{t_1}, \quad (4.4)$$

and correspondingly for $\tilde{\mathcal{Y}}$ (with an indeterminate λ). The theorem provides us with a method to construct exact solutions of the ncpKP hierarchy in (\mathcal{A}, \cdot) . The idea is now to take the dispersionless limit of (4.2) and (4.4). This should then result in conditions that determine exact solutions of the pdCM hierarchy in (\mathcal{A}, \cdot) . However, assuming for $\tilde{\mathcal{X}}, \tilde{\mathcal{Y}}$ power series expansions in ϵ with nonvanishing terms of zeroth order, this results in too restrictive conditions. The way out is to note that a ‘gauge transformation’

$$\tilde{\mathcal{X}} = \mathcal{X}G, \quad \tilde{\mathcal{Y}} = \mathcal{Y}G, \tag{4.5}$$

with an $N \times N$ matrix G , leaves ϕ invariant. Choosing

$$G = \exp(\xi(\mathbf{t}, P)), \quad \xi(\mathbf{t}, P) := \sum_{n \geq 1} t_n P^n, \tag{4.6}$$

with a constant $N \times N$ matrix P , and using $\xi(\mathbf{t}, P)_{-[\lambda]} = \xi(\mathbf{t}, P) + \ln(\mathcal{I}_N - \lambda P)$ with the $N \times N$ unit matrix \mathcal{I}_N , the heat hierarchy equations are mapped to

$$(\mathcal{X} - \mathcal{X}_{-[\lambda]})(\lambda^{-1} - P) = \mathcal{X}_{t_1}, \quad (\mathcal{Y} - \mathcal{Y}_{-[\lambda]})(\lambda^{-1} - P) = \mathcal{Y}_{t_1}, \tag{4.7}$$

and (4.2) is converted into

$$\mathcal{X}_{t_1} + \mathcal{X}P = R\mathcal{X} + Q\mathcal{Y}. \tag{4.8}$$

Assuming that

$$\mathcal{X}(\mathbf{t}, \epsilon) = \mathcal{X}_{(0)}(\mathbf{T}) + \mathcal{O}(\epsilon), \quad \mathcal{Y}(\mathbf{t}, \epsilon) = \mathcal{Y}_{(0)}(\mathbf{T}) + \mathcal{O}(\epsilon), \tag{4.9}$$

$$R = R_{(0)} + \mathcal{O}(\epsilon), \quad Q = Q_{(0)} + \mathcal{O}(\epsilon), \tag{4.10}$$

and P independent of ϵ , then we obtain from (4.8)

$$\epsilon \mathcal{X}_{(0), T_1} + \mathcal{X}_{(0)}P = R_{(0)}\mathcal{X}_{(0)} + Q_{(0)}\mathcal{Y}_{(0)} + \mathcal{O}(\epsilon), \tag{4.11}$$

and from (4.7)

$$\epsilon D(\lambda)\mathcal{X}_{(0)}(\lambda^{-1} - P) = \epsilon \mathcal{X}_{(0), T_1} + \mathcal{O}(\epsilon^2), \tag{4.12}$$

together with the same equation for $\mathcal{Y}_{(0)}$. After dividing the last equation by ϵ , these equations have the dispersionless limits

$$R_{(0)}\mathcal{X}_{(0)} + Q_{(0)}\mathcal{Y}_{(0)} = \mathcal{X}_{(0)}P, \tag{4.13}$$

respectively,

$$D(\lambda)\mathcal{X}_{(0)}(\lambda^{-1} - P) = \mathcal{X}_{(0), T_1}, \quad D(\lambda)\mathcal{Y}_{(0)}(\lambda^{-1} - P) = \mathcal{Y}_{(0), T_1}, \tag{4.14}$$

which is

$$\frac{1}{n+1}\mathcal{X}_{(0), T_{n+1}} = \frac{1}{n}\mathcal{X}_{(0), T_n}P, \quad \frac{1}{n+1}\mathcal{Y}_{(0), T_{n+1}} = \frac{1}{n}\mathcal{Y}_{(0), T_n}P, \tag{4.15}$$

($n = 1, 2, \dots$), or in terms of the variables (3.10),

$$\mathcal{X}_{(0), x_{n+1}} = \mathcal{X}_{(0), x_n}P, \quad \mathcal{Y}_{(0), x_{n+1}} = \mathcal{Y}_{(0), x_n}P, \quad n = 1, 2, \dots \tag{4.16}$$

Under the stated conditions, we have an expansion

$$\phi(\mathbf{t}, \epsilon) = \Phi(x_1, x_2, \dots) + \mathcal{O}(\epsilon), \tag{4.17}$$

which determines an exact solution Φ of the dispersionless limit of the ncpKP hierarchy, i.e. the pdCM hierarchy (3.11). Proposition 5.1 in the following section confirms this directly, i.e. without reference to the scaling limit procedure applied to the ncpKP hierarchy and the above theorem.

5. Exact solutions of the pdCM hierarchy

The main result of the preceding section will be formulated in the next proposition, and we provide a direct proof. It will then be further elaborated and applied in order to construct some classes of exact solutions of the $(su(m))$ pdCM hierarchy. In this section, symbols like \mathcal{X} and Q , for example, correspond to $\mathcal{X}_{(0)}$ and $Q_{(0)}$ in the preceding sections. Since now we resolve our considerations from the dispersionless limit procedure, there is no need to carry these indices with us any more. In fact, this section can be accessed almost completely without reference to the previous ones.

Proposition 5.1. *Let \mathcal{X} be an invertible $N \times N$ and \mathcal{Y} an $M \times N$ matrix such that*

$$R\mathcal{X} + Q\mathcal{Y} = \mathcal{X}P \tag{5.1}$$

and

$$\mathcal{X}_{x_{n+1}} = \mathcal{X}_{x_1} P^n, \quad \mathcal{Y}_{x_{n+1}} = \mathcal{Y}_{x_1} P^n, \quad n = 1, 2, \dots, \tag{5.2}$$

with constant matrices P, R of size $N \times N$, and Q of size $N \times M$. Then

$$\Phi = \mathcal{Y}\mathcal{X}^{-1} \tag{5.3}$$

solves the pdCM hierarchy

$$\Phi_{x_{n+1}, x_m} - \Phi_{x_{m+1}, x_n} = [\Phi_{x_n}, \Phi_{x_m}]_Q \quad m, n = 1, 2, \dots \tag{5.4}$$

(which is (3.11) with $Q_{(0)}$ replaced by Q).

Proof. Equation (5.2) is equivalent to

$$\mathcal{X}_{x_{n+1}} = \mathcal{X}_{x_n} P, \quad \mathcal{Y}_{x_{n+1}} = \mathcal{Y}_{x_n} P, \quad n = 1, 2, \dots$$

In terms of the maps d, \bar{d} , defined in section 3.3, this can be expressed as

$$\bar{d}\mathcal{X} = d\mathcal{X}P, \quad \bar{d}\mathcal{Y} = d\mathcal{Y}P.$$

Hence

$$\begin{aligned} (d\Phi)\mathcal{X}P + \Phi d\mathcal{X}P &= d\mathcal{Y}P = \bar{d}\mathcal{Y} = (\bar{d}\Phi)\mathcal{X} + \Phi \bar{d}\mathcal{X} \\ &= (\bar{d}\Phi)\mathcal{X} + \Phi d\mathcal{X}P, \end{aligned}$$

and thus⁵

$$\bar{d}\Phi = (d\Phi)W,$$

where

$$W := \mathcal{X}P\mathcal{X}^{-1} = Q\Phi + R,$$

using (5.1). Since d and \bar{d} satisfy (3.36), and since Q and R are constant, we obtain

$$\bar{d}d\Phi = -d\bar{d}\Phi = (d\Phi) \wedge dW = (d\Phi) \wedge Qd\Phi,$$

which is the hierarchy (5.4). □

The next result shows how to obtain via proposition 5.1 solutions of the pdCM hierarchy in the algebra of $m \times m$ matrices with the *usual* matrix product (i.e. without the modification by a matrix Q different from the unit matrix). If $\text{tr}(R) = \text{tr}(P)$, these solutions have values in $sl(m, \mathbb{C})$.

⁵ We note that this equation can be written as $\bar{d}\Phi - (d\Phi)Q\Phi = d\Theta$ with $\Theta := \Phi R$, which is (3.8).

Proposition 5.2. *Let U, V be $N \times m$ matrices and*

$$\varphi = U^\dagger \Phi V, \quad Q = VU^\dagger, \tag{5.5}$$

where $\Phi = \mathcal{Y}\mathcal{X}^{-1}$ with \mathcal{X}, \mathcal{Y} solving (5.1). Then

$$\text{tr}(\varphi) = \text{tr}(P) - \text{tr}(R). \tag{5.6}$$

Under the conditions of proposition 5.1, and if U and V are constant, φ solves the pdCM hierarchy (3.14).

Proof. We have

$$\text{tr}(\varphi) = \text{tr}(U^\dagger \mathcal{Y}\mathcal{X}^{-1} V) = \text{tr}(VU^\dagger \mathcal{Y}\mathcal{X}^{-1}) = \text{tr}(Q\mathcal{Y}\mathcal{X}^{-1}).$$

Using (5.1), this can be rewritten as

$$\text{tr}(\varphi) = \text{tr}(\mathcal{X}P\mathcal{X}^{-1} - R),$$

which is (5.6). The last statement of the proposition is easily verified (see also section 3). \square

It is helpful to extend (5.1) to

$$H\mathcal{Z} = \mathcal{Z}P, \tag{5.7}$$

where

$$\mathcal{Z} = \begin{pmatrix} \mathcal{X} \\ \mathcal{Y} \end{pmatrix}, \quad H = \begin{pmatrix} R & Q \\ S & L \end{pmatrix}, \tag{5.8}$$

with constant matrices L, S . Indeed, the upper component of (5.7) reproduces (5.1). But now we have an additional equation, namely $S\mathcal{X} + L\mathcal{Y} = \mathcal{Y}P$ (which together with (5.1) implies the algebraic Riccati equation $S + L\Phi - \Phi R - \Phi Q\Phi = 0$ for Φ). Although the latter appears to impose an unnecessary restriction, it will be helpful in order to determine interesting classes of exact solutions. The two equations (5.2) can be combined into

$$\mathcal{Z}_{x_{n+1}} = \mathcal{Z}_{x_1} P^n \quad n = 1, 2, \dots \tag{5.9}$$

Obviously, a transformation

$$\mathcal{Z} = \Gamma \mathcal{Z}', \quad H = \Gamma H' \Gamma^{-1}, \tag{5.10}$$

with a constant matrix

$$\Gamma = \begin{pmatrix} \Gamma_{11} & \Gamma_{12} \\ \Gamma_{21} & \Gamma_{22} \end{pmatrix}, \tag{5.11}$$

preserves the form of equations (5.7) and (5.9) with the same P . Consequently, if \mathcal{Z}' solves (5.7) and (5.9) with H' , and hence $\Phi' = \mathcal{Y}'\mathcal{X}'^{-1}$ solves the pdCM hierarchy with Q' , then \mathcal{Z} solves the corresponding equations with H , and according to proposition 5.1

$$\Phi = \mathcal{Y}\mathcal{X}^{-1} = (\Gamma_{21} + \Gamma_{22}\Phi')(\Gamma_{11} + \Gamma_{12}\Phi')^{-1} \tag{5.12}$$

solves the pdCM hierarchy with Q .⁶ Such a transformation thus relates solutions of different versions of the pdCM hierarchy, i.e. with different Q (which means different products). Since Q and Q' may have different rank, via (5.5) one obtains corresponding solutions of a pdCM hierarchy in a different matrix algebra. An extreme case is $Q' = 0$. Then the hierarchy (5.4) reduces to the system of linear equations

$$\Phi'_{x_m x_{n+1}} - \Phi'_{x_n x_{m+1}} = 0 \quad m, n = 1, 2, \dots \tag{5.13}$$

⁶ We note that the transformation (3.15) corresponds to the block-diagonal choice $\Gamma = \text{diag}(g_2, g_1^{-1})$. Such a transformation does not change a solution Φ in an essential way.

The above observation now suggests to first construct a solution Φ' of these linear equations, and then use such a transformation (as a ‘dressing transformation’) to generate a solution of a nonlinear hierarchy.

Proposition 5.3. *Let P, L, R be constant $N \times N$ matrices. Let $\mathcal{X}', \mathcal{Y}'$ solve (5.2) (which is (5.9)) and (5.7) with⁷*

$$H' = \begin{pmatrix} R & 0 \\ 0 & L \end{pmatrix}. \tag{5.14}$$

Then

$$\Phi = \mathcal{Y}' (\mathcal{X}' - K\mathcal{Y}')^{-1} \tag{5.15}$$

with any constant $N \times M$ matrix K , provided that the inverse in (5.15) exists, solves the pdCM hierarchy (5.4) with

$$Q = RK - KL. \tag{5.16}$$

Proof. Choosing in (5.10) the transformation matrix

$$\Gamma = \begin{pmatrix} \mathcal{I}_N & -K \\ 0 & \mathcal{I}_M \end{pmatrix},$$

where \mathcal{I}_N is the $N \times N$ unit matrix, we have

$$H = \Gamma H' \Gamma^{-1} = \begin{pmatrix} R & RK - KL \\ 0 & L \end{pmatrix},$$

and hence $Q = RK - KL$. Since $\mathcal{Z} = \Gamma \mathcal{Z}'$ again satisfies (5.7) and (5.9), proposition 5.1 tells us that Φ given by (5.12), which is (5.15), solves (5.4) with Q given by (5.16). \square

A special case of proposition 5.3 is formulated next. This will turn out to be particularly useful in the following.

Corollary 5.1. *Let P, K be constant $N \times N$ matrices, and \mathcal{X}' an $N \times N$ matrix solution of*

$$\mathcal{X}'_{x_{n+1}} = \mathcal{X}'_{x_1} P^n \quad n = 1, 2, \dots, \tag{5.17}$$

such that

$$[P, \mathcal{X}'] = 0. \tag{5.18}$$

Then

$$\Phi = (\mathcal{X}' - K)^{-1}, \tag{5.19}$$

provided that the inverse exists, solves the pdCM hierarchy with Q given by

$$Q = [P, K]. \tag{5.20}$$

If moreover (5.5) holds, then φ solves the pdCM hierarchy (3.14) in $sl(m, \mathbb{C})$.

Proof. We check that the assumptions of this corollary constitute a special case of those of proposition 5.3. Equation (5.7) decomposes into

$$R\mathcal{X}' = \mathcal{X}'P, \quad L\mathcal{Y}' = \mathcal{Y}'P.$$

Choosing

$$R = L = P, \quad \mathcal{Y}' = \mathcal{I}_N,$$

⁷ If \mathcal{X}' is invertible, then $\Phi' = \mathcal{Y}'\mathcal{X}'^{-1}$ solves the linear hierarchy (5.13). This follows from proposition 5.1, since $Q' = 0$.

this reduces to (5.18), and (5.2) reduces to (5.17). Since $\Phi'^{-1} = \mathcal{X}'$, (5.15) becomes (5.19), and (5.16) becomes (5.20). As a consequence of $R = P$ and proposition 5.2, φ has vanishing trace, hence takes values in $sl(m, \mathbb{C})$. \square

Example. Let us choose

$$P = \begin{pmatrix} p_1 & 0 \\ 0 & p_2 \end{pmatrix}, \quad \mathcal{X}' = \begin{pmatrix} f_1 & 0 \\ 0 & f_2 \end{pmatrix}, \quad Q = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (5.21)$$

with real constants $p_1 \neq p_2$ and real functions f_i . Then (5.18) holds and (5.17) requires that the function f_i depends on the variables x_1, x_2, \dots only through the combination $\omega_i = \sum_{n \geq 1} p_i^{n-1} x_n$. Equation (5.20) is solved by

$$K = \begin{pmatrix} 0 & (p_1 - p_2)^{-1} \\ (p_2 - p_1)^{-1} & 0 \end{pmatrix}. \quad (5.22)$$

A diagonal part of K can be absorbed in (5.19) by redefinition of f_1, f_2 . We obtain

$$\Phi = \frac{1}{D} \begin{pmatrix} f_2 & (p_2 - p_1)^{-1} \\ (p_1 - p_2)^{-1} & f_1 \end{pmatrix}, \quad (5.23)$$

where

$$D = f_1 f_2 + (p_1 - p_2)^{-2}, \quad (5.24)$$

and then the following solution of the $sl(2, \mathbb{R})$ pdCM hierarchy:

$$\varphi = \Phi Q = \frac{1}{D} \begin{pmatrix} (p_2 - p_1)^{-1} & f_2 \\ f_1 & (p_1 - p_2)^{-1} \end{pmatrix}. \quad (5.25)$$

The corresponding conserved density \mathcal{E} is given by

$$\mathcal{E} = -\frac{1 + p_1 p_2}{(f_1 f_2 + (p_1 - p_2)^{-2})^2} \frac{df_1}{d\omega_1} \frac{df_2}{d\omega_2}, \quad (5.26)$$

which can take both signs, depending on the values of the parameters. Choosing

$$f_i = \exp(q_i \omega_i) + c_i \quad i = 1, 2, \quad (5.27)$$

with non-negative constants c_i and real constants $q_i \neq 0$, the solution is regular (for all x_1, x_2, \dots). For positive c_i , \mathcal{E} is exponentially localized, a sort of soliton. The first derivatives of the components of φ are *not* localized, however. If c_1 or c_2 tends to zero, it stretches into a half-infinitely extended ‘line soliton’, the location of which is determined by $q_1 \omega_1 + q_2 \omega_2 = 0$.

As pointed out in section 3.1, the case where φ given by (5.5) has values in the Lie algebra of a unitary group is distinguished by the fact that there is a non-negative ‘energy’ functional, with density given by \mathcal{E} defined in (3.24). We will therefore concentrate on this case in the following. We further restrict our considerations to the case $M = N$, hence $\mathcal{X}, \mathcal{Y}, \Phi$ are all $N \times N$ matrices. Let U and V be constant $N \times m$ matrices. If Φ has the property

$$\Phi^\dagger = T \Phi T^{-1} \quad (5.28)$$

with a constant invertible $N \times N$ matrix T which is anti-Hermitian, i.e. $T^\dagger = -T$, then by setting

$$U = T V \quad (5.29)$$

we achieve that $\varphi = U^\dagger \Phi V$ is *anti-Hermitian*, i.e.,

$$\varphi^\dagger = -\varphi. \quad (5.30)$$

As a consequence of these conditions, we have

$$\varphi = -V^\dagger T \Phi V, \quad (5.31)$$

and

$$Q = VU^\dagger = -VV^\dagger T, \quad (5.32)$$

which has the property

$$Q^\dagger = -TQT^{-1}. \quad (5.33)$$

We note that $V \mapsto V\sigma$, with a constant unitary $m \times m$ matrix σ , leaves Q invariant and induces a gauge transformation $\varphi \mapsto \sigma^\dagger \varphi \sigma$. This can be used to reduce the freedom in the choice of V .

In the following, we address exact solutions of the $su(m)$ pdCM hierarchy by using the recipe of corollary 5.1. Accordingly we should arrange that the solution \mathcal{X}' of the linear hierarchy (5.17) satisfies

$$\mathcal{X}'^\dagger = T\mathcal{X}'T^{-1}. \quad (5.34)$$

If also

$$K^\dagger = TKT^{-1}, \quad (5.35)$$

then Φ given by (5.19) satisfies the same relation, i.e. (5.28). As a further consequence, (5.31) is then anti-Hermitian.

Together with (5.34), (5.18) implies $[T^{-1}P^\dagger T, \mathcal{X}'] = 0$, which is identically satisfied as a consequence of (5.18) if P has the property

$$P^\dagger = TPT^{-1}. \quad (5.36)$$

We note that (5.20) is consistent with (5.33), (5.35) and (5.36). Basically the problem of constructing solutions of (3.14) in $su(m)$ (on the basis of corollary 5.1) is reduced to the problem of satisfying the algebraic equation (5.20) with Q given by (5.32). We summarize our results.

Proposition 5.4. *Let (P, \mathcal{X}', T, V) be data consisting of a constant $N \times N$ matrix P , an $N \times N$ matrix \mathcal{X}' , which solves (5.17) and (5.18), a constant anti-Hermitian $N \times N$ matrix T , and a constant $N \times m$ matrix V . Furthermore, let (5.34) and (5.36) be satisfied, Q be defined by (5.32), and suppose that a solution K of (5.20) and (5.35) exists. Then $\varphi = -V^\dagger T \Phi V$, with Φ given by (5.19), is a solution of the pdCM hierarchy (3.14) in the Lie algebra $su(m)$.*

By application of proposition 5.4, some classes of exact solutions of the $su(m)$ pdCM hierarchy will be derived in the following subsections. Examples are worked out for the $su(2)$ case. Corresponding plots are restricted to the three variables entering the first hierarchy equation, and we will always use the coordinates t, x, y related to the variables x_1, x_2, x_3 by the transformation (3.19). This is mainly done in order to ease a comparison with solutions of Ward's modified chiral model (cf section 3.2).

5.1. A class of solutions of the $su(m)$ pdCM hierarchy

Assuming that P is diagonal, i.e.,

$$P = \text{diag}(p_1, \dots, p_N), \quad (5.37)$$

with complex constants $p_i \neq p_j$ for $i \neq j$, (5.18) requires \mathcal{X}' to be diagonal. Writing

$$\mathcal{X}' = \text{diag}(f_1, \dots, f_N), \quad (5.38)$$

where the entries are functions of x_1, x_2, \dots , (5.17) becomes

$$f_{j,x_n} = p_j^{n-1} f_{j,x_1} \quad j = 1, \dots, N, \quad n = 1, 2, \dots \quad (5.39)$$

This is solved if f_j is a holomorphic⁸ function of

$$\omega_j := \sum_{n \geq 1} x_n p_j^{n-1} \quad (5.40)$$

(with the same j both for the function and its argument). In particular, f_j depends on the variables x_1, x_2, \dots only through the combination (5.40). Condition (5.34) with an invertible anti-Hermitian matrix T imposes restrictions on the set of functions $\{f_j\}_{j=1,\dots,N}$, see section 5.1.1.

According to corollary 5.1, Φ given by (5.19) solves the pdCM hierarchy with Q given by (5.20), which implies $Q_{ii} = 0, i = 1, \dots, N$, and

$$K_{ij} = \frac{Q_{ij}}{p_i - p_j} = - \sum_{a=1}^m \sum_{k=1}^N \frac{V_{ia} V_{ka}^* T_{kj}}{p_i - p_j}, \quad i \neq j, \quad (5.41)$$

where we took (5.32) into account. Equation (5.19) shows that a diagonal part of K can be absorbed by redefinition of the functions f_j . Hence it is no restriction to assume that $K_{ii} = 0$ for $i = 1, \dots, N$. Equation (5.35) is then satisfied as a consequence of (5.36). Now (5.19) can be expressed as

$$(\Phi^{-1})_{ij} = f_i \delta_{ij} + \sum_{k=1}^N \sum_{a=1}^m \frac{V_{ia} V_{ka}^* T_{kj}}{p_i - p_j}, \quad (5.42)$$

and $\varphi = -V^\dagger T \Phi V$ solves the $su(m)$ pdCM hierarchy (3.14).

5.1.1. Some regular and localized solutions of the $su(2)$ pdCM hierarchy. Choosing N even and for T the following block-diagonal form:

$$T = \begin{pmatrix} 0 & -1 & & & & \\ 1 & 0 & & & & \\ & & \ddots & & & \\ & & & 0 & -1 & \\ & & & 1 & 0 & \end{pmatrix}, \quad (5.43)$$

conditions (5.34) and (5.36) read

$$f_2(\omega_2) = f_1(\omega_1)^*, \dots, f_N(\omega_N) = f_{N-1}(\omega_{N-1})^*, \quad (5.44)$$

$$p_2 = p_1^*, \dots, p_N = p_{N-1}^*. \quad (5.45)$$

We also have $\omega_2 = \omega_1^*, \dots, \omega_N = \omega_{N-1}^*$.

Example 1. Let $N = 2$ and $V = \mathcal{I}_2$, which leads to $Q = -T$. Then we obtain

$$\varphi = \frac{\beta}{1 + \beta^2 |f(\omega)|^2} \begin{pmatrix} -i & \beta f(\omega) \\ -\beta f(\omega)^* & i \end{pmatrix}, \quad (5.46)$$

⁸ More generally, the function f_j is allowed to have singularities in the complex ω_j -plane, but we will not consider such solutions in this work. See also, e.g., [44] in the case of Ward's chiral model.




Figure 1. Plots of \mathcal{E} for stationary lump solutions of the $su(2)$ pdCM equation, according to example 1 of section 5.1.1. Here we chose $p = i$ and $f(\omega) = \omega/2$ (left), $f(\omega) = \omega^2/2$ (middle) and $f(\omega) = (\omega^6 - 1)/8$ (right).

where $i = \sqrt{-1}$, $\beta = 2\Im(p)$, $p = p_1$, and $f = f_1$ is an arbitrary holomorphic function of $\omega = \sum_{n \geq 1} p^{n-1} x_n$. This solution is regular for all x_1, x_2, \dots . The corresponding ‘energy density’ is given by

$$\mathcal{E} = \beta^4 (1 + |p|^2)(1 + \beta^2 |f(\omega)|^2)^{-2} \left| \frac{df}{d\omega} \right|^2. \tag{5.47}$$

Choosing for f a non-constant polynomial in ω , the solution is rational and localized, thus a field configuration that is often called a ‘lump’. The shape of \mathcal{E} depends on the degree of the polynomial and in particular on its zeros.

Let $x_4, x_5, \dots = 0$ for the moment, so we concentrate on the first hierarchy equation. Applying the coordinate transformation (3.19), we have

$$\omega = \frac{1}{2}(t - x + 2py + p^2(t + x)). \tag{5.48}$$

We note that this becomes t -independent if $p = \pm i$ (i.e. $\beta = \pm 2$), in which case $\omega = -x \pm iy$, and the solution φ is stationary. For $f(\omega) = q\omega + c$, we obtain a simple lump. For example, choosing $p = i$ and $f(\omega) = \omega/2$, we have

$$\mathcal{E} = \frac{8}{(1 + x^2 + y^2)^2}, \tag{5.49}$$

see figure 1. $c \neq 0$ causes a displacement of the lump in the xy -plane.

For $f(\omega) = q(\omega - c)^n$, $n > 1$, with a zero of n th order, \mathcal{E} is bowl shaped. In particular, if $p = i$ and $f(\omega) = \omega^2/2$, we have

$$\mathcal{E} = \frac{32(x^2 + y^2)}{(1 + (x^2 + y^2)^2)^2}, \tag{5.50}$$

which is shown in figure 1 (second plot). The third plot in figure 1 displays another example.

Configurations with M lumps are obtained by choosing f as a product of (powers of) factors $\omega - c_i$, $i = 1, \dots, M$, with pairwise different complex constants c_i .

Example 2. Let $N = 4$ and

$$V = \begin{pmatrix} \mathcal{I}_2 \\ \mathcal{I}_2 \end{pmatrix}. \tag{5.51}$$

Then φ has the following components:

$$\begin{aligned} \varphi_{11} = -\varphi_{22} &= \frac{1}{\mathcal{D}}(\beta_1\beta_2(ah_1^*h_2 - a^*h_1h_2^*) - i[(\beta_1 + \beta_2)|b|^4 + \beta_1|ah_2|^2 + \beta_2|ah_1|^2]), \\ \varphi_{12} = -\varphi_{21}^* &= \frac{1}{\mathcal{D}}((b^*)^2(a\beta_1h_1 + a^*\beta_2h_2) + a|h_1|^2\beta_2h_2 + a^*\beta_1h_1|h_2|^2), \end{aligned} \tag{5.52}$$




Figure 2. Plots of \mathcal{E} at $t = -2, 0, 2$ for a three lump solution of the $su(2)$ pdCM equation, according to example 2 of section 5.1.1. Here we chose $p_1 = i, p_2 = 2i$, and $f_1(\omega_1) = \omega_1^2 + 2, f_2(\omega_2) = \omega_2/4$. Due to the special choice of p_1 , a pair of lumps is stationary. The positions of the latter are given by the zeros of $f_1(\omega_1) = x^2 - y^2 + 2 - 2ixy$, which are located at $(x, y) = (0, \pm\sqrt{2})$. The position of the third lump corresponds to the zero of $f_2(\omega_2)$, which is given by $(x, y) = (-3t/5, 0)$. Choosing -2 instead of $+2$ in $f_1(\omega_1)$, all three lumps are located on the line $y = 0$, and the third lump moves through both members of the pair (which then reside at $x = \pm\sqrt{2}$).

where

$$\begin{aligned} \beta_i &= 2\Im(p_i), & a &= p_1 - p_2^*, & b &= p_1 - p_2, \\ h_1 &= a\beta_1 f_1, & h_2 &= a^* \beta_2 f_2, \\ \mathcal{D} &= (|b|^2 + |h_1|^2)(|b|^2 + |h_2|^2) + \beta_1 \beta_2 |h_1 - h_2|^2. \end{aligned} \tag{5.53}$$

This solution is regular since \mathcal{D} is positive (note that $|b| > 0$ since $p_1 \neq p_2, |b|^2 \geq -\beta_1 \beta_2$, and use $|h_1|^2 + |h_2|^2 \geq |h_1 - h_2|^2$). Figure 2 shows an example. For generic parameter values, plots of \mathcal{E} show lumps with apparently trivial interaction. But if p_1, p_2 are close to the values $\pm i$ (that correspond to the stationary single lump solutions), a non-trivial interaction is observed in a compact space region, see figure 3. The scalar KP-I equation possesses solutions with the same behaviour [85]. Moreover, also dipolar vortices (modons) of a barotropic equation [86] and BPS monopoles [87] show such a behaviour in head-on collisions.

5.2. Another class of solutions of the $su(m)$ pdCM hierarchy

Let N be even. We introduce the commuting matrices

$$\mathcal{X}_I = \begin{pmatrix} f_I & \tilde{h}_I \\ 0 & f_I \end{pmatrix}, \quad P_I = \begin{pmatrix} p_I & 1 \\ 0 & p_I \end{pmatrix}, \tag{5.54}$$

with pairwise different complex constants p_I , and functions $f_I, h_I, I = 1, \dots, N/2$, and construct in terms of them the block-diagonal matrices

$$\mathcal{X}' = \begin{pmatrix} \mathcal{X}_1 & & & \\ & \mathcal{X}_2 & & \\ & & \ddots & \\ & & & \mathcal{X}_{N/2} \end{pmatrix}, \quad P = \begin{pmatrix} P_1 & & & \\ & P_2 & & \\ & & \ddots & \\ & & & P_{N/2} \end{pmatrix}, \tag{5.55}$$

which then obviously also commute. Now (5.17) becomes

$$f_{I,x_n} = p_I^{n-1} f_{I,x_1}, \quad \tilde{h}_{I,x_n} = p_I^{n-1} \tilde{h}_{I,x_1} + (n-1)p_I^{n-2} f_{I,x_1}, \tag{5.56}$$

where $I = 1, \dots, N/2$ and $n = 1, 2, \dots$. Writing

$$\tilde{h}_I = h_I + \frac{\partial f}{\partial p_I}, \tag{5.57}$$




Figure 3. Plots of \mathcal{E} at $t = -110, -100, -99, -90, 0, 90, 99, 100, 110$ for the solution in example 2 of section 5.1.1 with the data $p_1 = -19i/20, p_2 = 21i/20, f_1 = 2i\omega_1, f_2 = 2i\omega_2$. Two lumps approach each other in the x -direction, merge, move away from one another in the y -direction up to some maximal distance, return to each other and merge again, and then separate in the x -direction.

the second equation is turned into $h_{I,x_n} = p_I^{n-1}h_{I,x_1}$, by use of the first. Hence (5.17) is satisfied if, for $I = 1, \dots, N/2$, f_I and h_I are holomorphic functions of

$$\omega_I = \sum_{n \geq 1} x_n p_I^{n-1} \tag{5.58}$$

(which is (5.40)), and in particular only depend on the variables x_1, x_2, \dots through this combination. In order to explore the consequences of (5.20), we write K and Q as $N/2 \times N/2$ matrices, where the components K_{IJ} , respectively Q_{IJ} , are 2×2 matrices.

Proposition 5.5. *With the matrix P defined in (5.55), and any Q , the solution of (5.20) is given by*

$$K_{IJ} = \frac{Q_{IJ}}{p_I - p_J} - \frac{[\Pi_2, Q_{IJ}]}{(p_I - p_J)^2} + \frac{[\Pi_2, [\Pi_2, Q_{IJ}]]}{(p_I - p_J)^3} \tag{5.59}$$

for $I \neq J$, and

$$[\Pi_2, K_{JJ}] = Q_{JJ} \quad J = 1, \dots, N/2, \tag{5.60}$$

where

$$\Pi_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}. \tag{5.61}$$

Proof. We write $P_I = p_I \mathcal{I}_2 + \Pi_2$. Then (5.20), restricted to components with $I \neq J$, takes the form

$$\left(\text{id} + \frac{1}{p_I - p_J} \text{ad}_{\Pi_2} \right) K_{IJ} = \frac{Q_{IJ}}{p_I - p_J},$$

where $\text{ad}_{\Pi} K = [\Pi, K]$. Now (5.59) follows from

$$\left(\text{id} + \frac{1}{p_I - p_J} \text{ad}_{\Pi_2} \right)^{-1} = \sum_{k=0}^2 (-1)^k (p_I - p_J)^{-k} \text{ad}_{\Pi_2}^k,$$

since $\text{ad}_{\Pi_2}^3 = 0$. The diagonal components of (5.20) are $\text{ad}_{\Pi_2} K_{JJ} = Q_{JJ}$, which is (5.60). □

Remark. In view of (5.19), we may always assume that the two upper entries of K_{JJ} vanish (since non-vanishing entries can be absorbed into \mathcal{X}'). Using the matrix T given below in (5.62), condition (5.35) then implies $K_{2J-1,2J-1}^\dagger = \tau_2 K_{2J,2J} \tau_2$, $J = 1, \dots, N/2$, and this requires that K_{JJ} can only have a nonzero entry in the lower left corner. As a consequence of (5.60), Q_{JJ} is then diagonal and has vanishing trace.

A simple way of satisfying (5.60) is to choose V such that the diagonal blocks Q_{JJ} vanish, and then set $K_{JJ} = 0$, $J = 1, \dots, N/2$. This will be done in section 5.2.1.

It remains to satisfy the further anti-hermiticity conditions.

5.2.1. *su(2) lumps with ‘anomalous’ scattering.* Let N now be a multiple of 4. In analogy with (5.43) we set

$$T = \begin{pmatrix} 0 & -\tau_2 & & & & \\ \tau_2 & 0 & & & & \\ & & \ddots & & & \\ & & & 0 & -\tau_2 & \\ & & & \tau_2 & 0 & \end{pmatrix} \quad \text{where} \quad \tau_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (5.62)$$

Then (5.36) means $P_1^\dagger = \tau_2 P_2 \tau_2, \dots, P_{N/2-1}^\dagger = \tau_2 P_{N/2} \tau_2$, which by use of (5.54) amounts to

$$p_{2J} = p_{2J-1}^* \quad J = 1, \dots, N/2. \quad (5.63)$$

Since we address the case $m = 2$, V has to be chosen as an $N \times 2$ matrix, which we subdivide into 2×2 blocks V_I , $I = 1, \dots, N/2$. It follows that

$$Q_{JJ} = \begin{cases} -V_J V_{J+1}^\dagger \tau_2 & \text{if } J \text{ is odd} \\ V_J V_{J-1}^\dagger \tau_2 & \text{even.} \end{cases} \quad (5.64)$$

Thus, in order to achieve that $Q_{JJ} = 0$, we must arrange that

$$V_{2J-1} V_{2J}^\dagger = 0 \quad J = 1, \dots, N/2. \quad (5.65)$$

Then Q has the following structure:

$$Q = \begin{pmatrix} 0 & V_1 V_1^\dagger & -V_1 V_4^\dagger & V_1 V_3^\dagger & \cdots \\ -V_2 V_2^\dagger & 0 & -V_2 V_4^\dagger & V_2 V_3^\dagger & \cdots \\ -V_3 V_2^\dagger & V_3 V_1^\dagger & 0 & V_3 V_3^\dagger & \cdots \\ -V_4 V_2^\dagger & V_4 V_1^\dagger & -V_4 V_4^\dagger & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \tau_2. \quad (5.66)$$

Example. The simplest case is $N = 4$. Excluding degenerate cases, the two blocks V_1, V_2 of V should both have rank 1. Hence $V_1 = v_1 u_1^\dagger, V_2 = v_2 u_2^\dagger$ with vectors $u_J, v_J, J = 1, 2$, satisfying $u_1^\dagger u_2 = 0$. With a unitary transformation σ we can achieve that the lower component of u_1 vanishes. It follows that the upper component of u_2 also vanishes. By a redefinition of v_1, v_2 , we obtain $u_1^\dagger = (1, 0)$ and $u_2^\dagger = (0, 1)$, and thus

$$\begin{aligned} Q_{12} &= v_1(\tau_2 v_1)^\dagger = \begin{pmatrix} v_{11} v_{12}^* & |v_{11}|^2 \\ |v_{12}|^2 & v_{12} v_{11}^* \end{pmatrix}, \\ Q_{21} &= -v_2(\tau_2 v_2)^\dagger = -\begin{pmatrix} v_{21} v_{22}^* & |v_{21}|^2 \\ |v_{22}|^2 & v_{22} v_{21}^* \end{pmatrix}, \end{aligned} \tag{5.67}$$

with an obvious notation for the components of v_1 and v_2 . We should exclude the case when the expression (5.59) for K reduces to the first term on the right-hand side, since this leads back to the solution of example 1 in section 5.1. This case is ruled out if v_{12} or v_{22} is different from zero, which suggests to choose $v_J = (0, 1), J = 1, 2$, and thus

$$V_1 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad V_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}. \tag{5.68}$$

Then proposition 5.5 yields

$$K = \begin{pmatrix} 0 & K_{12} \\ -K_{21} & 0 \end{pmatrix}, \tag{5.69}$$

where

$$K_{IJ} = \begin{pmatrix} -(p_I - p_J)^{-2} & -2(p_I - p_J)^{-3} \\ (p_I - p_J)^{-1} & (p_I - p_J)^{-2} \end{pmatrix} \tag{5.70}$$

for $I \neq J$. This in turn allows us to compute Φ and then also φ . The anti-hermiticity conditions are then satisfied by setting

$$p_2 = p_1^*, \quad f_2(\omega_2) = f_1(\omega_1)^*, \quad h_2(\omega_2) = h_1(\omega_1)^*, \tag{5.71}$$

and we have $\omega := \omega_1 = \omega_2^*$. The result is

$$\begin{aligned} \varphi_{11} &= -\varphi_{22} = -\frac{i}{\beta^7 \mathcal{D}} (2 + \beta^4 |f|^2 + \beta^4 |f + i\beta \tilde{h}|^2), \\ \varphi_{12} &= -\varphi_{21}^* = \frac{1}{\beta^5 \mathcal{D}} (4if - \beta \tilde{h} - \beta^5 f^2 \tilde{h}^*), \end{aligned} \tag{5.72}$$

with $\beta = 2\Im(p_1), f = f_1(\omega), \tilde{h} = \tilde{h}_1$ given by (5.57) in terms of f and $h = h_1(\omega)$, and

$$\mathcal{D} = \beta^{-8} (1 + \beta^4 |f|^2)^2 + \beta^{-4} |2f + i\beta \tilde{h}|^2. \tag{5.73}$$

The solution φ is thus regular for any choice of p_1 , with non-vanishing imaginary part, and the holomorphic functions f, h . An example with $h = 0$ is shown in figure 4. More interesting structures appear for non-constant h . Indeed, figure 5 shows two lumps that scatter at an angle of 90° . Choosing f linear in ω and h proportional to ω^n , we observe a π/n scattering. Figures 6 and 7 show examples of 60° , respectively, 45° scattering.

Solutions with π/n scattering have also been found in Ward's chiral model numerically [35, 37], and analytically as certain limits of families of non-interacting lumps [32, 39, 43, 50]. Moreover, also the scalar KP equation (with positive dispersion, i.e. KP I) possesses solutions with this behaviour [88–93]. In fact, π/n scattering in head-on collisions of soliton-like objects is a familiar feature of many models (see [94–96], in particular). It occurs in dipolar




Figure 4. Plots of \mathcal{E} at $t = -50, 0, 50$ for the solution of section 5.2.1 with the data $p_1 = i$ (i.e. $\beta = 2$), $f = 2\omega$ and $h = 0$.




Figure 5. Plots of \mathcal{E} at $t = -5, 0, 5$ for a 2-lump solution, exhibiting ‘scattering at right angle’, see section 5.2.1. Here we chose $p_1 = i$ and $f = -i\omega/32, h = -\omega^2/4$.




Figure 6. Plots of \mathcal{E} at $t = -500, 0, 500$ for a 3-lump configuration exhibiting 60° scattering. This is obtained from the solution of the example in section 5.2.1 with the data $p_1 = i$ and $f = -i\omega, h = \omega^3/8$.




Figure 7. Plots of \mathcal{E} at $t = -20, 0, 20$ for a 4-lump configuration exhibiting 45° scattering. This is obtained from the solution of the example in section 5.2.1 with the data $p_1 = i$ and $f = -i\omega, h = \omega^4/8$.

vortex collisions [86, 97–99], in $O(3)$ and $\mathbb{C}P^1$ models [37, 100–106], in Skyrme models [95, 106–109], for vortices of the Abelian Higgs (or Ginzburg–Landau) model [106, 110–119], and BPS monopoles of a $SU(m)$ Yang–Mills–Higgs system [87, 106, 120–122]. Another integrable system that possesses solutions with this behaviour is the Davey–Stewartson II




Figure 8. Plots of \mathcal{E} at $t = -20, 0, 20$ for the solution of the example in section 5.3 with the data $p = i, f = -i\omega, h = \omega^4/8$ and $g = 0$. At $t = 0$ we have cut off an extremely large lump in the centre.

and (5.2) becomes

$$\begin{aligned} f_{x_n} &= p^{n-1} f_{x_1}, & \tilde{h}_{x_n} &= p^{n-1} \tilde{h}_{x_1} + (n-1)p^{n-2} f_{x_1}, \\ \tilde{g}_{x_n} &= p^{n-1} \tilde{g}_{x_1} + (n-1)p^{n-1} \tilde{h}_{x_1} + \frac{1}{2}(n-1)(n-2)p^{n-3} f_{x_1}. \end{aligned} \tag{5.78}$$

Writing

$$\tilde{h} = h + \frac{\partial f}{\partial p}, \quad \tilde{g} = g + \frac{\partial h}{\partial p} + \frac{1}{2} \frac{\partial^2 f}{\partial p^2}, \tag{5.79}$$

with functions f, g, h , it follows that these equations are satisfied if the latter are arbitrary holomorphic functions of $\omega = \sum_{n \geq 1} p^n x_n$. Furthermore, we find that

$$K = \begin{pmatrix} 0 & K_{12} \\ -K_{12}^* & 0 \end{pmatrix} \quad \text{with} \quad K_{12} = \begin{pmatrix} i/\beta^3 & 3/\beta^4 & -6i/\beta^5 \\ 1/\beta^2 & -2i/\beta^3 & -3/\beta^4 \\ -i/\beta & -1/\beta^2 & i/\beta^3 \end{pmatrix} \tag{5.80}$$

and $\beta = 2\Im(p)$, solves $[P, K] = Q$ with $Q = -VV^\dagger T$. The resulting class of solutions is regular since

$$\begin{aligned} \det(\mathcal{X}' - K) &= \beta^{-18} (1 + \beta^6 |f|^2)^3 + 2\beta^{-12} (1 + \beta^6 |f|^2) |3if - \beta \tilde{h}|^2 \\ &\quad + \beta^{-6} |2f^2 + (f + i\beta \tilde{h})^2 + \beta^2 f \tilde{g}|^2 + \beta^{-12} |6f + \beta(4i\tilde{h} - \beta \tilde{g})|^2. \end{aligned} \tag{5.81}$$

Now we have three arbitrary holomorphic functions at our disposal, so this class exhibits quite a variety of different structures. Figures 8 and 9 show some examples. If $h = g = 0$, the typical behaviour is similar to that shown in figure 4.

Comparing the data that determine the class of solutions in the example in section 5.2.1, based on a conjugate pair of 2×2 Jordan normal form matrices P_I , with those of the last example, which is based on a conjugate pair of 3×3 Jordan normal form matrices, there is an obvious generalization to the case of conjugate pairs of larger Jordan normal form matrices P_I . We note in particular that proposition 5.5 can be generalized. Since the solutions turned out to be automatically *regular* in the 2×2 and 3×3 case, it may well be that this holds in general. But a proof of this conjecture is out of reach so far.

5.4. Superposing solutions

The data that determine solutions of the $su(m)$ pdCM hierarchy on the basis of proposition 5.4 are given by a set of matrices (P, \mathcal{X}', T, V) . Let two such sets be given, $(P_i, \mathcal{X}'_i, T_i, V_i), i = 1, 2$, with associated matrices $Q_i = -V_i V_i^\dagger T_i, K_i$ (as solutions of (5.20)), and Φ_i given by




Figure 9. Plots of \mathcal{E} at $t = -1000, 0, 1000$ for the solution of the example in section 5.3 with the data $p = i, f = -i\omega/40, h = \omega^4/10$ and $g = 0$. At $t = 0$ we have cut off the lumps beyond a certain height.

(5.19). P_i, \mathcal{X}_i, T_i are $N_i \times N_i$ matrices and V_i is an $N_i \times m$ matrix. We can combine them into the larger matrices

$$\begin{aligned}
 P &= \begin{pmatrix} P_1 & 0 \\ 0 & P_2 \end{pmatrix}, & \mathcal{X}' &= \begin{pmatrix} \mathcal{X}'_1 & 0 \\ 0 & \mathcal{X}'_2 \end{pmatrix}, \\
 T &= \begin{pmatrix} T_1 & 0 \\ 0 & T_2 \end{pmatrix}, & V &= \begin{pmatrix} V_1 \\ V_2 \end{pmatrix}.
 \end{aligned}
 \tag{5.82}$$

Obviously, (P, \mathcal{X}', T, V) again satisfies (5.17), (5.18), (5.34), (5.36) and $T^\dagger = -T$. Equation (5.32) becomes

$$Q = \begin{pmatrix} Q_1 & Q_{12} \\ Q_{21} & Q_2 \end{pmatrix}, \quad Q_{12} = -V_1 V_2^\dagger T_2, \quad Q_{21} = -V_2 V_1^\dagger T_1, \tag{5.83}$$

and (5.20) with

$$K = \begin{pmatrix} K_1 & K_{12} \\ K_{21} & K_2 \end{pmatrix} \tag{5.84}$$

yields the equations

$$P_1 K_{12} - K_{12} P_2 = Q_{12}, \quad P_2 K_{21} - K_{21} P_1 = Q_{21}. \tag{5.85}$$

The off-diagonal blocks of K are a source of complexity and non-triviality of the resulting superposition. Equation (5.35) then determines K_{21} in terms of K_{12} (or vice versa),

$$K_{21} = T_2^{-1} K_{12}^\dagger T_1. \tag{5.86}$$

As a consequence, the second of equations (5.85) follows from the first. If we find a solution⁹ K_{12} of the remaining equation, then we obtain

$$\Phi = \begin{pmatrix} \Phi_1 & \Phi_1 K_{12} \Phi_2 \\ \Phi_2 K_{21} \Phi_1 & \Phi_2 \end{pmatrix} \begin{pmatrix} A_1^{-1} & 0 \\ 0 & A_2^{-1} \end{pmatrix}, \tag{5.87}$$

⁹ Choosing V_1 and V_2 such that $V_1 V_2^\dagger = 0$, we have $Q_{12} = 0$ and (5.85) is solved by $K_{12} = 0$. It follows that φ is simply the sum of the solutions φ_1 and φ_2 . But $V_1 V_2^\dagger = 0$ also implies that $\varphi_1 \varphi_2 = 0$, hence both constituent solutions φ_1, φ_2 must be degenerate, i.e. cannot have full rank.

where

$$A_1 = \mathcal{I}_{N_1} - K_{12}\Phi_2 K_{21}\Phi_1, \quad A_2 = \mathcal{I}_{N_2} - K_{21}\Phi_1 K_{12}\Phi_2, \quad (5.88)$$

and φ given by (5.31) solves the $su(m)$ pdCM hierarchy, provided that the inverses of A_1 and A_2 exist. If the two matrices Φ_i are regular (and thus also the corresponding solutions φ_i), then Φ and thus also φ is regular if and only if $\det(A_1)\det(A_2) \neq 0$ (for all values of x_1, x_2, \dots). Since $\det(A_1) = \det(A_2)$ by an application of Sylvester's determinant theorem, this reduces to the condition

$$\det(A_1) \neq 0. \quad (5.89)$$

We note also that $\det(A_1)$ is real since

$$\begin{aligned} \det(A_1)^* &= \det(A_1^\dagger) = \det(\mathcal{I}_{N_1} - K_{21}^\dagger \Phi_2^\dagger K_{12}^\dagger \Phi_1^\dagger) \\ &= \det(\mathcal{I}_{N_1} - T_1 K_{12} \Phi_2 K_{21} \Phi_1 T_1^{-1}) = \det(A_1). \end{aligned} \quad (5.90)$$

Example. We choose

$$\begin{aligned} P_1 &= \begin{pmatrix} p_1 & 1 & & \\ 0 & p_1 & & \\ & & p_1^* & 1 \\ & & 0 & p_1^* \end{pmatrix}, \quad \mathcal{X}'_1 = \begin{pmatrix} f_1 & h_1 + \partial f_1 / \partial p_1 & & \\ 0 & f_1 & & \\ & & f_1^* & h_1^* + (\partial f_1 / \partial p_1)^* \\ & & 0 & f_1^* \end{pmatrix}, \\ T_1 &= \begin{pmatrix} & & -1 & \\ & & & \\ & 1 & & \\ 1 & & & \end{pmatrix}, \quad V_1 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix}, \end{aligned} \quad (5.91)$$

where f_1, h_1 are arbitrary holomorphic functions of ω_1 (with ω_1 defined in (5.40)), and

$$P_2 = \begin{pmatrix} p_2 & 0 \\ 0 & p_2^* \end{pmatrix}, \quad \mathcal{X}'_2 = \begin{pmatrix} f_2 & 0 \\ 0 & f_2^* \end{pmatrix}, \quad T_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad V_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (5.92)$$

with an arbitrary holomorphic function f_2 of ω_2 . Thus we superpose data corresponding to a regular solution of the kind treated in the example in section 5.2.1 and data corresponding to a regular solution as given in example 1 of section 5.1. In the following we assume that

$$p_1 \neq p_2, \quad p_1 \neq p_2^*. \quad (5.93)$$

Together with the conditions $p_i \neq p_i^*, i = 1, 2$, which the data of the components have to satisfy, this means that the constants p_i and their complex conjugates are pairwise different. The second condition in (5.93) is in fact needed for the matrix K to exist. K has the form (5.84), where K_1 is given by the 4×4 matrix K in (5.69) with the pair (p_1, p_1^*) . Furthermore,

$$K_2 = \begin{pmatrix} 0 & \frac{1}{p_2 - p_2^*} \\ \frac{1}{p_2^* - p_2} & 0 \end{pmatrix}, \quad K_{21} = \begin{pmatrix} 0 & 0 & -\frac{1}{p_1^* - p_2} & \frac{1}{(p_1^* - p_2)^2} \\ \frac{1}{p_1 - p_2^*} & -\frac{1}{(p_1 - p_2^*)^2} & 0 & 0 \end{pmatrix}, \quad (5.94)$$




Figure 10. Plots of \mathcal{E} at $t = -2, 0, 2$ for a superposition of a 2-lump configuration, with ‘anomalous scattering’, and a single lump (which is at the top of the left plot and at the bottom of the right plot), according to the example of section 5.4. Here we chose $p_1 = i, p_2 = -3i/8, f_1 = -i\omega_1/32, f_2 = 24\omega_2$ and $h_1 = -\omega^2/4$.

and K_{12} is then determined by (5.86). With some efforts the expression for $\det(A_1)$ can be brought into the form

$$\begin{aligned} \det(A_1) = & |a|^{-8} (1 + \beta_2^2 |f_2|^2)^{-1} (|w|^2 + (1 + \beta_1^4 |f_1|^2)^2)^{-2} [(|a|^4 [|w|^2 + (1 + \beta_1^4 |f_1|^2)^2] \\ & - \beta_1 \beta_2 [|a|^2 (1 + \beta_1^4 |f_1|^2) + |b|^2 + |aw + ib^* \beta_1^2 f_1|^2])^2 \\ & + |(a^*)^4 (|w|^2 + (1 + \beta_1^4 |f_1|^2)^2) \beta_2 f_2 \\ & + \beta_1 \beta_2 (b^2 w + (a^*)^2 \beta_1^4 f_1^2 w^* - 2ia^* b \beta_1^2 f_1 (1 + \beta_1^4 |f_1|^2))|^2], \end{aligned} \quad (5.95)$$

where

$$\beta_i = 2\Im(p_i), \quad a = p_1 - p_2^*, \quad b = p_1 - p_2, \quad w = \beta_1^3 h_1 - 2i\beta_1^2 f_1. \quad (5.96)$$

The regularity condition (5.89) turns out to be automatically satisfied. This is seen as follows. First we note that

$$|a|^4 = |b|^4 + \beta_1 \beta_2 (|a|^2 + |b|^2) > \beta_1 \beta_2 (|a|^2 + |b|^2), \quad (5.97)$$

as a consequence of the first of the inequalities (5.93), and thus

$$|a|^4 (1 + \beta_1^4 |f_1|^2)^2 > \beta_1 \beta_2 (|a|^2 + |b|^2) (1 + \beta_1^4 |f_1|^2). \quad (5.98)$$

Using $|a|^2 > \beta_1 \beta_2$, this leads to

$$\begin{aligned} |a|^4 (|w|^2 + (1 + \beta_1^4 |f_1|^2)^2) & > \beta_1 \beta_2 (|a|^2 (1 + \beta_1^4 |f_1|^2) + |b|^2 + |aw|^2 + |ib^* \beta_1^2 f_1|^2) \\ & \geq \beta_1 \beta_2 (|a|^2 (1 + \beta_1^4 |f_1|^2) + |b|^2 + |aw + ib^* \beta_1^2 f_1|^2), \end{aligned} \quad (5.99)$$


which implies $\det(A_1) > 0$.

Figure 10 shows plots of \mathcal{E} at consecutive times, for a special choice of the data.

In the last example, the regularity of the superposition turned out to be a consequence of the ‘regular data’ we started with. But this example also demonstrates that it is quite difficult in general to evaluate the regularity condition (5.89). We note that also the cases treated in sections 5.1.1 and 5.2.1 may be regarded as special cases of ‘superpositions’ as formulated above. In particular, example 2 of section 5.1.1 provides us with another example where the superposition of regular data turned out to be regular again. It is unlikely that this is a special feature of our particular examples. But in order to tackle a general proof, we probably need different methods.

6. Conclusions

We summarize the relations between integrable systems and their hierarchies considered in this work in the following diagram:



Here pKP_Q and pdCM_Q stand, respectively, for the pKP and pdCM hierarchy in the matrix algebra $GL(M \times N, \mathbb{C})$ with product modified by a constant matrix Q (see (4.1)). pdCM_Q is related by pseudoduality (see (3.28)) to the hierarchy Ward_Q of Ward’s model with dependent variable in $GL(M \times N, \mathbb{C})$ (and product modified by Q). If $\text{rank}(Q) = 1$, solutions of pKP_Q are mapped to solutions of the scalar pKP hierarchy, an additional condition ensures that the resulting solution is real. Analogously, if $\text{rank}(Q) = m$ and an $su(m)$ condition holds, solutions of the pdCM_Q hierarchy are mapped to solutions of the $su(m)$ pdCM hierarchy (which is pseudodual to the hierarchy associated with Ward’s modified $SU(m)$ chiral model). Concerning the Miura transformation between the (matrix) pKP hierarchy and the modified KP (mKP) hierarchy, and its dispersionless limit (see the dashed arrows in the diagram), see [76]. The relations provided by the dispersionless scaling limits in the diagram have actually been anticipated in [76] (see the remark in section 4 therein).

In the present work, we demonstrated how the dispersionless scaling can be used to transfer a method of constructing exact solutions from the (matrix or ‘noncommutative’) pKP hierarchy to the pdCM hierarchy. Indeed, proposition 5.1 is an analogue of theorem 4.1 in [75] (which we recalled as theorem 4.1). We showed that large classes of exact solutions of the pdCM hierarchy can be obtained with its help. In particular, we presented examples of various multiple lump configurations of the $su(m)$ pdCM. The general result formulated in proposition 5.1 is a source of even more classes of exact solutions.

Our method to generate exact solutions of the $su(m)$ pdCM hierarchy is based on quite simple formulae and quickly produces interesting solutions (like lumps with ‘anomalous scattering’). But a more systematic treatment, in particular of multi-lump solutions, requires deeper methods (of matrix calculus), and further insights are needed as to how the *a priori* given plethora of parameters can efficiently be reduced. It would also be of interest to compare this method with an inverse scattering approach.

Solutions of the $su(m)$ pdCM hierarchy can also be obtained from solutions of Ward’s chiral model hierarchy by integrating (3.28) (or equivalently (3.43)). In any case, one should expect an analogous structure of localized solutions, and this indeed turns out to be the case in examples. A deviation in the corresponding plots is caused by the fact that the ‘energy’ expression for the pdCM differs from the energy of the Ward model by a term that causes an asymmetry in the x -direction, see the remark in section 3.2 and also the appendix¹⁰. Our method to generate solutions of the pdCM hierarchy seems to be quite different from the methods that were used to construct solutions of Ward’s model. In particular, in the latter model solutions with ‘anomalous scattering’ have been obtained by taking suitable limits of

¹⁰ We note that x and y have to be exchanged for comparing our formulae with those in the literature on the Ward model.

families of non-interacting lump solutions. In our approach, corresponding solutions of the pdCM hierarchy are directly given by matrix data involving Jordan blocks. Moreover, we have seen that even the simple multiple lump solutions of section 5.1.1 can exhibit an anomalous behaviour within some compact space region (see figure 3), whereas asymptotically (i.e. compared at large enough negative and positive times) no deflection is observed. We noted that this has a KP-I counterpart [85] and also analogues in some other systems [86, 87].

The fact that the dispersionless scaling limit of matrix pKP (respectively mKP) is a simple reduction of a potential version of the (four-dimensional) self-dual Yang–Mills equation raises the question whether there is a (four-dimensional) integrable system that has the full self-dual Yang–Mills equation as a dispersionless limit and that admits a reduction to matrix KP (respectively mKP).

Appendix A. The ‘anti-pdCM hierarchy’

Writing (3.42) as $Jd(J^{-1}\bar{d}J)J^{-1} = 0$, using the Leibniz rule and (3.36), we obtain the equivalent form

$$\bar{d}((dJ)J^{-1}) = 0 \tag{A.1}$$

of the hierarchy associated with Ward’s chiral model, where d and \bar{d} exchanged their roles. This is integrated by introducing a potential $\tilde{\varphi}$ such that

$$(dJ)J^{-1} = \bar{d}\tilde{\varphi}. \tag{A.2}$$

Rewriting the last equation in the form $dJ = (\bar{d}\tilde{\varphi})J$, we obtain the integrability condition

$$d\bar{d}\tilde{\varphi} = \bar{d}\tilde{\varphi} \wedge \bar{d}\tilde{\varphi}. \tag{A.3}$$

In components, this becomes

$$\tilde{\varphi}_{x_n x_{m+1}} - \tilde{\varphi}_{x_m x_{n+1}} = [\tilde{\varphi}_{x_{n+1}}, \tilde{\varphi}_{x_{m+1}}], \quad m, n = 1, 2, \dots, \tag{A.4}$$

which we refer to as the *anti-pdCM hierarchy*. For $m = 1$ and $n = 2$ we have

$$\tilde{\varphi}_{x_1 x_3} - \tilde{\varphi}_{x_2 x_2} = [\tilde{\varphi}_{x_2}, \tilde{\varphi}_{x_3}]. \tag{A.5}$$

In terms of the coordinates given by (3.19), it takes the form

$$\tilde{\varphi}_{tt} - \tilde{\varphi}_{xx} - \tilde{\varphi}_{yy} + [\tilde{\varphi}_t + \tilde{\varphi}_x, \tilde{\varphi}_y] = 0. \tag{A.6}$$

Since this equation is obtained from (3.20) by $x \mapsto -x$, so are its Lagrangian $\tilde{\mathcal{L}}$ and energy–momentum tensor \tilde{T}^μ_ν from those in section 3.1. For $\tilde{\varphi}$ in $su(m)$,

$$\tilde{\mathcal{E}} = \tilde{T}^0_0 + \tilde{T}^0_1 = -\frac{1}{2} \text{tr}((\tilde{\varphi}_t + \tilde{\varphi}_x)^2 + \tilde{\varphi}_y^2) \tag{A.7}$$

is then a *non-negative* conserved density.

Associated with any solution J of Ward’s chiral model hierarchy via (3.43) and (A.2), there are solutions φ and $\tilde{\varphi}$ of the pdCM hierarchy and the anti-pdCM hierarchy, respectively. Using $J_t J^{-1} - J_x J^{-1} = \tilde{\varphi}_y$ and $J_y J^{-1} = \tilde{\varphi}_t + \tilde{\varphi}_x$, which follow from (A.2), we find that

$$\tilde{\mathcal{E}} = \mathcal{E}_{\text{Ward}} + \text{tr}(J^{-1} J_t J^{-1} J_x). \tag{A.8}$$

Combining this with (3.33), leads to

$$\mathcal{E}_{\text{Ward}} = \frac{1}{2}(\mathcal{E} + \tilde{\mathcal{E}}). \tag{A.9}$$

The next result is an analogue of corollary 5.1 and leads to a class of solutions of the anti-pdCM hierarchy.

Proposition A1. *Let (P, K, \mathcal{X}') be data that determine via corollary 5.1 a solution Φ (given by (5.19)) of the pdCM hierarchy with $Q = [P, K]$. If P is invertible (as in all our examples in section 5), then Φ also solves*

$$d\bar{d}\Phi = \bar{d}\Phi \wedge \tilde{Q}\bar{d}\Phi \tag{A.10}$$

with $\tilde{Q} = P^{-1}QP^{-1}$.

Proof. As a consequence of (5.17), $\mathcal{X} := \mathcal{X}' - K$ solves

$$d\mathcal{X} = \bar{d}\mathcal{X}P^{-1},$$

and consequently $\Phi = \mathcal{X}^{-1}$ satisfies

$$d\Phi = (\bar{d}\Phi)W^{-1}, \quad W = \mathcal{X}P\mathcal{X}^{-1}.$$

Now we note that (5.18) and (5.20) imply $[P^{-1}, \mathcal{X}] = [K, P^{-1}] = P^{-1}QP^{-1}$. Hence

$$W^{-1} = P^{-1}(\mathcal{X} + QP^{-1})\mathcal{X}^{-1} = P^{-1} + P^{-1}QP^{-1}\Phi,$$

and we obtain

$$d\bar{d}\Phi = -\bar{d}d\Phi = (\bar{d}\Phi) \wedge P^{-1}QP^{-1}\bar{d}\Phi. \quad \square$$

If moreover the assumptions of proposition 5.4 are satisfied, then

$$P^{-1}QP^{-1} = -P^{-1}VV^\dagger TP^{-1} = -P^{-1}V(P^{-1}V)^\dagger T, \tag{A.11}$$

and (A.10) implies that

$$\tilde{\varphi} = -(P^{-1}V)^\dagger T\Phi P^{-1}V \tag{A.12}$$

solves the anti-Hermitian anti-pdCM hierarchy (A.3). The data (P, \mathcal{X}', T, V) therefore determine a solution (5.31) of the anti-Hermitian pdCM hierarchy and also a solution (A.12) of the anti-Hermitian anti-pdCM hierarchy.

Although elaboration of examples suggests that the pair $(\varphi, \tilde{\varphi})$ determined by the data (P, \mathcal{X}', T, V) indeed belongs to the same solution J of Ward's chiral model hierarchy (via (3.43) and (A.2)), we were not able so far to prove this.

References

- [1] Kodama Y 1988 A method for solving the dispersionless KP equation and its exact solutions *Phys. Lett. A* **129** 223–6
- [2] Kodama Y and Gibbons J 1989 A method for solving the dispersionless KP hierarchy and its exact solutions: II *Phys. Lett. A* **135** 167–70
- [3] Kupershmidt B A 1990 The quasiclassical limit of the modified KP hierarchy *J. Phys. A: Math. Gen.* **23** 871–86
- [4] Krichever I 1992 The dispersionless Lax equations and topological minimal models *Commun. Math. Phys.* **143** 415–29
- [5] Zakharov V E 1994 Dispersionless limit of integrable systems in $(2 + 1)$ -dimensions *Singular Limits of Dispersive Waves (NATO Adv. Sci. Inst. Ser. B Phys. vol 320)* ed N Ercolani, I Gabitov, D Levermore and D Serre (New York: Plenum) pp 165–75
- [6] Aoyama S and Kodama Y 1994 Topological conformal field theory with a rational W potential and the dispersionless KP hierarchy *Mod. Phys. Lett. A* **9** 2481–92
- [7] Takasaki K and Takebe T 1995 Integrable hierarchies and dispersionless limit *Rev. Math. Phys.* **7** 743–808
- [8] Takasaki K 1995 Symmetries and tau function of higher dimensional dispersionless integrable hierarchies *J. Math. Phys.* **36** 3574–607
- [9] Strachan I 1995 The Moyal bracket and the dispersionless limit of the KP hierarchy *J. Phys. A: Math. Gen.* **28** 1967–76
- [10] Strachan I A B 1997 A geometry for multidimensional integrable systems *J. Geom. Phys.* **21** 255–78

- [11] Chang J-H and Tu M-H 2000 On the Miura map between the dispersionless KP and dispersionless modified KP hierarchies *J. Math. Phys.* **41** 5391–406
- [12] Wiegmann P B and Zabrodin A 2000 Conformal maps and integrable hierarchies *Commun. Math. Phys.* **213** 523–38
- [13] Konopelchenko B, Martínez Alonso L and Ragnisco O 2001 The $\bar{\partial}$ -approach to the dispersionless KP hierarchy *J. Phys. A: Math. Gen.* **34** 10209–17
- [14] Dunajski M, Mason L J and Tod P 2001 Einstein–Weyl geometry, the dKP equation and twistor theory *J. Geom. Phys.* **37** 63–93
- [15] Dunajski M and Tod P 2002 Einstein–Weyl spaces and dispersionless Kadomtsev–Petviashvili equation from Painlevé I and II *Phys. Lett. A* **303** 253–64
- [16] Konopelchenko B G and Martínez Alonso L 2002 Dispersionless scalar integrable hierarchies, Whitham hierarchy, and the quasiclassical $\bar{\partial}$ -dressing method *J. Math. Phys.* **43** 3807–23
- [17] Konopelchenko B G and Martínez Alonso L 2002 Nonlinear dynamics on the plane and integrable hierarchies of infinitesimal deformations *Stud. Appl. Math.* **109** 313–36
- [18] Mañas M, Martínez Alonso L and Medina E 2002 Reductions and hodograph solutions of the dispersionless KP hierarchy *J. Phys. A: Math. Gen.* **35** 401–17
- [19] Mañas M, Martínez Alonso L and Medina E 2002 Reductions of the dispersionless KP hierarchy *Theor. Math. Phys.* **133** 1712–21
- [20] Guil F, Mañas M and Martínez Alonso L 2003 On twistor solutions of the dKP equation *J. Phys. A: Math. Gen.* **36** 6457–72
- [21] Ferapontov E V and Khusnutdinova K R 2004 Hydrodynamic reductions of multidimensional dispersionless PDEs: the test for integrability *J. Math. Phys.* **45** 2365–77
- [22] Manakov S V and Santini P M 2006 The Cauchy problem on the plane for the dispersionless Kadomtsev–Petviashvili equation *Preprint nlin.SI/0604023*
- [23] Konopelchenko B G and Magri F 2006 Coisotropic deformations of associative algebras and dispersionless integrable hierarchies *Preprint nlin.SI/0606069*
- [24] Zakharov V E and Kuznetsov E A 1986 Multi-scale expansions in the theory of systems integrable by the inverse scattering transform *Physica D* **18** 455–63
- [25] Leznov A N 1987 Equivalence of four-dimensional self-duality equations and the continuum analog of the principal chiral field problem *Theor. Math. Phys.* **73** 1233–7
- [26] Mason L J and Woodhouse N M J 1996 *Integrability, Self-Duality, and Twistor Theory* (Oxford: Clarendon)
- [27] Ward R S 1988 Soliton solutions in an integrable chiral model in $2 + 1$ dimensions *J. Math. Phys.* **29** 386–9
- [28] Ward R S 1988 Integrability of the chiral equations with torsion term *Nonlinearity* **1** 671–9
- [29] Ward R S 1989 Twistors in $2 + 1$ dimensions *J. Math. Phys.* **30** 2246–51
- [30] Ward R S 1990 Classical solutions of the chiral model, unitons, and holomorphic vector bundles *Commun. Math. Phys.* **128** 319–32
- [31] Ward R 1994 Sigma models in $2 + 1$ dimensions *Harmonic Maps and Integrable Systems (Aspects of Mathematics vol E23)* ed A P Fordy and J C Wood (Vieweg: Braunschweig) pp 193–202
- [32] Ward R S 1995 Nontrivial scattering of localized solitons in a $(2 + 1)$ -dimensional integrable system *Phys. Lett. A* **208** 203–8
- [33] Leese R A 1989 Extended wave solutions in an integrable chiral model in $2 + 1$ dimensions *J. Math. Phys.* **30** 2072–7
- [34] Villarroel J 1989 Scattering data for the self-duality equations *Inverse Problems* **5** 1157–62
- [35] Sutcliffe P M 1992 Non-trivial soliton scattering in an integrable chiral model in $(2 + 1)$ -dimensions *J. Math. Phys.* **33** 2269–78
- [36] Sutcliffe P M 1993 Yang–Mills–Higgs solitons in $2 + 1$ dimensions *Phys. Rev. D* **47** 5470–76
- [37] Piette B, Sutcliffe P M and Zakrzewski W J 1992 Soliton antisoliton scattering in $(2 + 1)$ dimensions *Int. J. Mod. Phys. C* **3** 637–60
- [38] Ioannidou T and Ward R S 1995 Conserved quantities for integrable chiral equations in $2 + 1$ dimensions *Phys. Lett. A* **208** 209–13
- [39] Ioannidou T 1996 Soliton solutions and nontrivial scattering in an integrable chiral model in $(2 + 1)$ dimensions *J. Math. Phys.* **37** 3422–41
- [40] Ioannidou T 2000 Soliton solutions of the modified chiral model in $(2 + 1)$ dimensions *Solitons: Properties, Dynamics, Interactions, Applications (CRM Series in Mathematical Physics)* ed R MacKenzie, M B Paranjape and W J Zakrzewski (Berlin: Springer) pp 75–9
- [41] Ioannidou T and Zakrzewski W 1998 A note on Ward’s chiral model *Phys. Lett. A* **242** 233–8
- [42] Ioannidou T and Zakrzewski W 1998 Lagrangian formulation of the general modified chiral model *Phys. Lett. A* **249** 303–6

- [43] Ioannidou T and Zakrzewski W 1998 Solutions of the modified chiral model in (2+1) dimensions *J. Math. Phys.* **39** 2693–701
- [44] Ioannidou T and Manton N S 2005 The energy of scattering solitons in the Ward model *Proc. R. Soc. Lond. A* **461** 1965–73
- [45] Anand C 1997 Ward's solitons *Geom. Top.* **1** 9–20
- [46] Anand C K 1998 Ward's solitons II: exact solutions *Can. J. Math.* **50** 1119–37
- [47] Zait R A, Mourad M F and Abdel-Wahab N H 2005 Nonlocal conserved currents for the modified chiral model in (2 + 1) dimensions *Can. J. Math.* **83** 1161–7
- [48] Dai B and Terng C-L 2005 Periodic and homoclinic solutions of the modified 2 + 1 chiral model *J. Math. Phys.* **46** 062706-1–14
- [49] Dai B, Terng C-L and Uhlenbeck K 2006 On the space-time monopole equation *Preprint math/0602607*
- [50] Dai B and Terng C-L 2007 Bäcklund transformations, Ward solitons, and unitons *J. Diff. Geom.* **75** 57–108
- [51] Dunajski M and Manton S 2005 Reduced dynamics of Ward solitons *Nonlinearity* **18** 1677–89
- [52] Dunajski M and Plansangkate P 2007 Topology and energy of time-dependent unitons *Proc. R. Soc. A* **463** 945–59
- [53] Ward R S 1999 Two integrable systems related to hyperbolic monopoles *Asian J. Math.* **3** 325–33
- [54] Kotecha V and Ward R S 2001 Integrable Yang–Mills–Higgs equations in 3-dimensional de Sitter space-time *J. Math. Phys.* **42** 1018–25
- [55] Ji X-F and Zhou Z-X 2005 Asymptotic behaviour of solitons with a double spectral parameter for the Bogomolny equation in (2 + 1)-dimensional anti de Sitter space *Chin. Phys. Lett.* **22** 1570–2
- [56] Lechtenfeld O and Popov A D 2001 Scattering of noncommutative solitons in 2 + 1 dimensions *Phys. Lett. B* **523** 178–84
- [57] Lechtenfeld O and Popov A D 2001 Noncommutative multi-solitons in 2 + 1 dimensions *J. High Energy Phys.* **JHEP11(2001)040**
- [58] Bieling S 2002 Interaction of noncommutative plane waves in 2 + 1 dimensions *J. Phys. A: Math. Gen.* **35** 6281–91
- [59] Wolf M 2002 Soliton-antisoliton scattering configurations in a noncommutative sigma model in 2+1 dimensions *J. High Energy Phys.* **JHEP06(2002)055**
- [60] Ihl M and Uhlmann S 2003 Noncommutative extended waves and soliton-like configurations in $N = 2$ string theory *Int. J. Mod. Phys. A* **18** 4889–932
- [61] Chu C-S and Lechtenfeld O 2005 Time-space noncommutative Abelian solitons *Phys. Lett. B* **625** 145–55
- [62] Chu C-S and Lechtenfeld O 2006 Emergence of time from dimensional reduction in noncommutative geometry *Mod. Phys. Lett. A* **21** 639–47
- [63] Klawunn M, Lechtenfeld O and Petersen S 2006 Moduli-space dynamics of noncommutative Abelian sigma-model solitons *J. High Energy Phys.* **JHEP06(2006)028**
- [64] Leznov A N and Mukhtarov M A 1987 Deformation of algebras and solution of self-duality equation *J. Math. Phys.* **28** 2574–8
- [65] Leznov A N and Saveliev M V 1989 Exactly and completely integrable nonlinear dynamical systems *Acta Appl. Math.* **16** 1–74
- [66] Papachristou C J 1989 Nonlocal currents for the self-dual Yang–Mills equation *Phys. Lett. A* **138** 493–6
- [67] Papachristou C J 1991 Recursion operator and current algebras for the potential $SL(N, \mathbb{C})$ self-dual Yang–Mills equation *Phys. Lett. A* **154** 29–34
- [68] Fokas A S and Ioannidou T 2001 The inverse problem for the Ward equation and for the 2 + 1 chiral model *Comm. Appl. Anal.* **5** 235–46
- [69] Fradkin E S and Tseytlin A A 1985 Quantum equivalence of dual field theories *Ann. Phys.* **162** 31–48
- [70] Curtright T and Zachos C 1994 Currents, charges, and canonical structure of pseudodual chiral models *Phys. Rev. D* **49** 5408–21
- [71] Curtright T and Zachos C 1995 The paradigm of pseudodual chiral models *PASCOS '94* ed K C Wali (Singapore: World Scientific) pp 381–90
- [72] Zakharov V E and Mikhailov A V 1978 Relativistically invariant two-dimensional models of field theory which are integrable by means of the inverse scattering problem method *Sov. Phys.—JETP* **47** 1017–27
- [73] Nappi C 1980 Some properties of an analog of the chiral model *Phys. Rev. D* **21** 418–20
- [74] Fridling B E and Jevicki A 1984 Dual representations and ultraviolet divergences in nonlinear σ -models *Phys. Lett. B* **134** 70–4
- [75] Dimakis A and Müller-Hoissen F 2007 Burgers and KP hierarchies: a functional representation approach *Theor. Math. Phys.* **152** 933–47
- [76] Dimakis A and Müller-Hoissen F 2006 Functional representations of integrable hierarchies *J. Phys. A: Math. Gen.* **39** 9169–86

- [77] Pohlmeyer K 1980 On the Lagrangian theory of anti-self-dual fields in four-dimensional Euclidean space *Commun. Math. Phys.* **72** 37–47
- [78] Manakov S V and Zakharov V E 1981 Three-dimensional model of relativistic-invariant field theory, integrable by the inverse scattering transform *Lett. Math. Phys.* **5** 247–53
- [79] Frölicher A and Nijenhuis A 1956 Theory of vector-valued differential forms: I. Derivations in the graded ring of differential forms *Proc. K. Ned. Acad. Wet. A* **59** 338–59
- [80] Crampin M, Sarlet W and Thompson G 2000 Bi-differential calculi, bi-Hamiltonian systems and conformal Killing tensors *J. Phys. A: Math. Gen.* **33** 8755–70
- [81] Dimakis A and Müller-Hoissen F 2000 Bi-differential calculi and integrable models *J. Phys. A: Math. Gen.* **33** 957–74
- [82] Takasaki K 1990 Hierarchy structure in integrable systems of gauge fields and underlying Lie algebras *Commun. Math. Phys.* **127** 225–38
- [83] Dimakis A and Müller-Hoissen F 2001 Bicomplexes and Bäcklund transformations *J. Phys. A: Math. Gen.* **34** 9163–94
- [84] Ogielski A T, Prasad M K, Sinha A and Chau Wang L-L 1980 Bäcklund transformations and local conservation laws for principal chiral fields *Phys. Lett. B* **91** 387–91
- [85] Lu Z, Tian E M and Grimshaw R 2004 Interaction of two lump solitons described by the Kadomtsev–Petviashvili I equation *Wave Motion* **40** 123–35
- [86] McWilliams J C and Zabusky N J 1982 Interactions of isolated vortices: I. Modons colliding with modons *Geophys. Astrophys. Fluid Dyn.* **19** 207–27
- [87] Gibbons G W and Manton N S 1986 Classical and quantum dynamics of BPS monopoles *Nucl. Phys. B* **274** 183–224
- [88] Gorshkov K A, Pelinovskii D E and Stepanyants Yu A 1993 Normal and anomalous scattering, formation and decay of bound states of two-dimensional solitons described by the Kadomtsev–Petviashvili equation *JETP* **77** 237–45
- [89] Pelinovsky D 1994 Rational solutions of the Kadomtsev–Petviashvili hierarchy and the dynamics of their poles: I. new form of a general rational solution *J. Math. Phys.* **35** 5820–30
- [90] Infeld E, Senatorski A and Skorupski A A 1995 Numerical simulations of Kadomtsev–Petviashvili soliton interactions *Phys. Rev. E* **51** 3183–91
- [91] Ablowitz M J and Villarroel J 1997 Solutions of the time dependent Schrödinger and the Kadomtsev–Petviashvili equations *Phys. Rev. Lett.* **78** 570–3
- [92] Villarroel J and Ablowitz M J 1999 On the discrete spectrum of the nonstationary Schrödinger equation and multipole lumps of the Kadomtsev–Petviashvili I equation *Commun. Math. Phys.* **207** 1–47
- [93] Ablowitz M J, Chakravarty S, Trubatch A D and Villarroel J 2000 A novel class of solutions of the non-stationary Schrödinger and the Kadomtsev–Petviashvili I equations *Phys. Lett. A* **267** 132–46
- [94] Rosenzweig C and Srivastava A M 1991 Towards a qualitative understanding of the scattering of topological defects *Phys. Rev. D* **43** 4029–41
- [95] Kudryavtsev A, Piette B and Zakrzewski W J 1993 π/N scattering in 2+1 dimensions *Phys. Lett. A* **180** 119–23
- [96] MacKenzie R 1995 Remarks on gauge vortex scattering *Phys. Lett. B* **352** 96–8
- [97] Nguyen Duc J-M and Sommeria J 1988 Experimental characterization of steady two-dimensional vortex couples *J. Fluid Mech.* **192** 175–92
- [98] van Heijst G J F and Flór J B 1989 Dipole formation and collisions in a stratified fluid *Nature* **340** 212–5
- [99] Voropayev S I and Afanasyev Ya D 1992 Two-dimensional vortex-dipole interactions in a stratified fluid *J. Fluid Mech.* **236** 665–89
- [100] Ward R S 1985 Slowly-moving lumps in the CP^1 model in $(2+1)$ dimensions *Phys. Lett. B* **158** 424–8
- [101] Leese R 1990 Low-energy scattering of solitons in the CP^1 model *Nucl. Phys. B* **344** 33–72
- [102] Leese R A 1991 Q -lumps and their interactions *Nucl. Phys. B* **366** 283–311
- [103] Zakrzewski W J 1991 Soliton-like scattering in the $O(3)$ σ -model in $(2+1)$ dimensions *Nonlinearity* **4** 429–75
- [104] Cova R J and Zakrzewski W J 1997 Soliton scattering in the $O(3)$ model on a torus *Nonlinearity* **10** 1305–17
- [105] Speight J M 1998 Lump dynamics in the CP^1 model on the torus *Commun. Math. Phys.* **194** 513–39
- [106] Manton N and Sutcliffe P 2004 *Topological Solitons* (Cambridge: Cambridge University Press)
- [107] Manton N S 1987 Is the $B = 2$ Skyrmeion axially symmetric? *Phys. Lett. B* **192** 177–9
- [108] Leese R A, Peyrard M and Zakrzewski W J 1990 Soliton scatterings in some relativistic models in $(2+1)$ dimensions *Nonlinearity* **3** 773–807
- [109] Sutcliffe P M 1991 The interaction of Skyrme-like lumps in $(2+1)$ dimensions *Nonlinearity* **4** 1109–21
- [110] Moriarty K J M, Myers E and Rebbi C 1988 Dynamical interactions of cosmic strings and flux vortices in superconductors *Phys. Lett. B* **207** 411–8
- [111] Ruback P J 1988 Vortex string motion in the Abelian Higgs model *Nucl. Phys. B* **296** 669–78

- [112] Shellard E P S and Ruback P J 1988 Vortex scattering in two dimensions *Phys. Lett. B* **209** 262–70
- [113] Myers E, Rebbi C and Strilka R 1992 Study of the interaction and scattering of vortices in the Abelian Higgs (or Ginzburg–Landau) model *Phys. Rev. D* **45** 1355–64
- [114] Strachan I 1992 Low-velocity scattering of vortices in a modified Abelian Higgs model *J. Math. Phys.* **33** 102–10
- [115] Samols T M 1992 Vortex scattering *Commun. Math. Phys.* **145** 149–79
- [116] Manton N S 1991 Vortices and anyons *Phys. Rev. Lett.* **67** 1462–3
- [117] Burzlaff J and McCarthy P 1991 A study of a 90° vortex–vortex scattering process *J. Math. Phys.* **32** 3376–80
- [118] Abdelwahid F and Burzlaff J 1994 Existence theorems for 90° vortex–vortex scattering *J. Math. Phys.* **35** 4651–60
- [119] Arthur K and Burzlaff J 1996 Existence theorems for π/n vortex scattering *Lett. Math. Phys.* **36** 311–8
- [120] Atiyah M F and Hitchin N J 1985 Low energy scattering of non-Abelian monopoles *Phys. Lett. A* **107** 21–5
- [121] Atiyah M F, Hitchin N J, Stuart J T and Tabor M 1985 Low-energy scattering of non-Abelian magnetic monopoles *Phil. Trans. R. Soc. A* **315** 459–69
- [122] Dancer A and Leese R 1993 Dynamics of $SU(3)$ monopoles *Proc. R. Soc. Lond. A* **440** 421–30
- [123] Mañas M and Santini P M 1997 Solutions of the Davey–Stewartson II equation with arbitrary rational localization and nontrivial interaction *Phys. Lett. A* **227** 325–34
- [124] Villarroel J and Ablowitz M J 2003 On the discrete spectrum of systems in the plane and the Davey–Stewartson II equation *SIAM J. Math. Anal.* **34** 1253–78
- [125] Ablowitz M J, Manakov S V and Schultz C L 1990 On the boundary conditions of the Davey–Stewartson equation *Phys. Lett. A* **148** 50–2