

The path integral formulation of fractional Brownian motion for the general Hurst exponent

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys. A: Math. Theor. 41 282002

(<http://iopscience.iop.org/1751-8121/41/28/282002>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.149

The article was downloaded on 03/06/2010 at 06:58

Please note that [terms and conditions apply](#).

FAST TRACK COMMUNICATION

The path integral formulation of fractional Brownian motion for the general Hurst exponent

I Calvo¹ and R Sánchez²¹ Laboratorio Nacional de Fusión, Asociación EURATOM-CIEMAT, 28040 Madrid, Spain² Fusion Energy Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USAE-mail: ivan.calvo@ciemat.es and sanchezferlr@ornl.gov

Received 9 May 2008, in final form 2 June 2008

Published 19 June 2008

Online at stacks.iop.org/JPhysA/41/282002**Abstract**

In 1995, Sebastian (1995 *J. Phys. A: Math. Gen.* **28** 4305) gave a path integral computation of the propagator of subdiffusive fractional Brownian motion (fBm), i.e. fBm with a Hurst or self-similarity exponent $H \in (0, 1/2)$. The extension of Sebastian's calculation to superdiffusion, $H \in (1/2, 1]$, becomes however quite involved due to the appearance of additional boundary conditions on fractional derivatives of the path. In this communication, we address the construction of the path integral representation in a different fashion, which allows us to treat both subdiffusion and superdiffusion on an equal footing. The derivation of the propagator of fBm for the general Hurst exponent is then performed in a neat and unified way.

PACS numbers: 05.40.-a, 02.50.Ey, 05.10.Gg

1. Introduction

The Langevin representation for Brownian motion has been known for a long time [1, 2]. In one dimension, it can be formally written as

$$x(t) = x_0 + \int_0^t \xi(t') dt', \quad (1)$$

where $\xi(t)$ is a Gaussian, uncorrelated noise function

$$\langle \xi(t) \xi(t') \rangle = \delta(t - t'). \quad (2)$$

The propagator of Brownian motion, which gives the probability of finding the Brownian walker at some location x at time $t > 0$ after having started at x_0 at time 0, is a Gaussian whose width grows as $t^{1/2}$:

$$G(x, t|x_0, 0) = \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{(x - x_0)^2}{2t}\right). \quad (3)$$

However, many interesting processes (see [3] for a review) ranging from hydrology [4] to finance [5] or tracer transport by turbulent flows [6] exhibit anomalous diffusion, $\langle (x - x_0)^2 \rangle \propto t^{2H}$, with $H \neq 1/2$. In many situations the anomalous behaviour is due to spatio-temporal correlations, which translate into long-term time correlations in Lagrangian quantities. A generalization of Brownian motion which was proposed to include such correlations is known as fractional Brownian motion (fBm) [7]. Although slightly different representations can be found in the literature, the more convenient one for our purposes is

$$x(t) = x_0 + \frac{1}{\Gamma(H + \frac{1}{2})} \int_0^t (t - t')^{H - \frac{1}{2}} \xi(t') dt', \quad (4)$$

where $\Gamma(\cdot)$ is the Gamma function and $\xi(t)$ is still Gaussian and uncorrelated³. The exponent H corresponds to both the self-similarity exponent of the process [7] and the Hurst exponent [10] of the incremental process $dx(t) := x(t + dt) - x(t)$. On the one hand, the average motion is invariant under the transformation $(x, t) \rightarrow (\mu^H x, \mu t)$. On the other hand, the increments $dx(t)$ are uncorrelated for $H = 1/2$ and anticorrelated (correlated) for $H \in (0, 1/2)$ ($H \in (1/2, 1)$). Consequently, (4) can be used to model diffusive ($H = 1/2$), subdiffusive ($H \in (0, 1/2)$) and superdiffusive ($H \in (1/2, 1)$) transport processes.

In a beautiful paper [11], Sebastian showed that it is very advantageous to rewrite fBm in terms of fractional differential operators,

$$x(t) = x_0 + {}_0D_t^{-(H+1/2)} \xi, \quad (5)$$

where ${}_0D_t^{-\alpha}$ is the Riemann–Liouville fractional integral of order α [12, 13]. Indeed, by combining path integral methods [14] with fractional calculus, he computed the exact propagator of (4) for $H \leq 1/2$. However, the case $H > 1/2$ is not addressed in [11] because the problem becomes quite more complicated from the technical point of view. The difficulties may be explained as follows. Sebastian constructs the path integral by inverting (5) in terms of Riemann–Liouville fractional differential operators. This is easy for $H \leq 1/2$ because such inversion is well defined simply imposing the boundary conditions $x(0) = x_0$ and $x(T) = x_T$. However, if $H > 1/2$ additional boundary conditions involving fractional derivatives at $t = 0$ are required. Not only is the physical meaning of these ‘fractional initial conditions’ unclear, but also the computation along the lines of [11] results too cumbersome.

In this communication, we formulate the path integral for fBm using instead $\xi(t)$ as the integration variable. This choice avoids the need to invert (5), and thus the need for the additional fractional initial conditions. In our approach the difficulty resides in how to manage the boundary conditions of $x(t)$, which translate into a non-local constraint on the admissible realizations of the noise. We will see below that the usage of Lagrange multipliers allows us to do this in an easy and transparent way, thus providing a general path integral derivation of the well-known propagator of fBm [15].

2. Path integral calculation of the propagator

Given a Langevin equation describing the motion of a particle, the propagator $G(x_T, T | x_0, 0)$ is defined as the probability of finding the particle at $x = x_T$ at time $t = T$; if initially, $t = 0$, it was located at $x = x_0$. Hence, the propagator can be viewed as the motion of the particle

³ In [7, 8] the fractional Brownian motion is defined by (4) except that the lower limit of the integral is taken at $-\infty$, and the issue of boundary conditions is not present. Our definition is better suited for the class of applications we are interested in (e.g. analysis of a series of Lagrangian velocities in turbulent plasmas). For such physical problems, the stochastic differential equation is naturally formulated in a domain $t \in (t_0, \infty)$, with finite t_0 . An equation analogous to (4) was used in [9], where fBm and fractional time processes are compared.

averaged over all realizations of the noise compatible with the boundary conditions $x(0) = x_0$ and $x(T) = x_T$. In the path integral formalism this is formulated as follows. The essential object is the probability measure $\mathcal{P}(\xi(t))\mathcal{D}\xi(t)$ on the space of maps $\xi(t) : [0, T] \rightarrow \mathbb{R}$. Since $\xi(t)$ is uncorrelated in time and distributed as a Gaussian for each t , this is naturally defined as

$$\mathcal{P}(\xi(t))\mathcal{D}\xi(t) = \exp\left(-\frac{1}{2} \int_0^T \xi(t)^2 dt\right) \mathcal{D}\xi(t). \tag{6}$$

We want to compute the propagator of (5). First, note that the boundary conditions $x(0) = x_0, x(T) = x_T$ translate into the following constraint on $\xi(t)$:

$${}_0D_T^{-(H+1/2)}\xi = x_T - x_0. \tag{7}$$

Therefore, the propagator can be written as the expectation value

$$G(x_T, T|x_0, 0) = \int \delta({}_0D_T^{-(H+1/2)}\xi - (x_T - x_0)) \exp\left(-\frac{1}{2} \int_0^T \xi(t)^2 dt\right) \mathcal{D}\xi(t), \tag{8}$$

where the Dirac delta function ensures that we only integrate over maps $\xi(t)$ satisfying (7).

We proceed to compute the path integral on the right-hand side of (8). Define the action

$$S[\xi(t)] = \frac{1}{2} \int_0^T \xi(t)^2 dt, \tag{9}$$

and consider infinitesimal variations on the space of maps $\xi(t)$ satisfying (7). We denote by $\bar{\xi}(t)$ the map which makes the action stationary under such variations. Observe that, in particular, $\bar{\xi}(t)$ verifies (7). Now, we perform the following change of variables in (8):

$$\xi(t) = \bar{\xi}(t) + \eta(t). \tag{10}$$

The constraint (7) implies that

$${}_0D_T^{-(H+1/2)}\eta = 0. \tag{11}$$

Using that the action is quadratic in $\xi(t)$ and that $\bar{\xi}(t)$ makes it stationary,

$$G(x_T, T|x_0, 0) = e^{-S[\bar{\xi}(t)]} \int \delta({}_0D_T^{-(H+1/2)}\eta) e^{-S[\eta(t)]} \mathcal{D}\eta(t), \tag{12}$$

whence we deduce that the propagator is of the form

$$G(x_T, T|x_0, 0) = f(T) e^{-S[\bar{\xi}(t)]}. \tag{13}$$

The function $f(T)$ will be determined at the end of the calculation by imposing the normalization condition

$$\int_{-\infty}^{\infty} G(x_T, T|x_0, 0) dx_T = 1, \quad \forall T. \tag{14}$$

It only remains to compute the stationary points of $S[\xi(t)]$ subject to the constraint (7). It is difficult to do it directly, for the condition (7) is hard to implement. The technique of Lagrange multipliers saves the day, though. Our problem is equivalent to finding the stationary points of

$$\tilde{S}[\xi(t), \lambda] = \frac{1}{2} \int_0^T \xi(t)^2 dt + \lambda({}_0D_T^{-(H+1/2)}\xi - (x_T - x_0)), \tag{15}$$

under arbitrary infinitesimal variations of $\xi(t)$ and the Lagrange multiplier $\lambda \in \mathbb{R}$.

From variations with respect to λ ,

$$\delta_\lambda \tilde{S}[\xi(t), \lambda] = \delta\lambda({}_0D_T^{-(H+1/2)}\xi - (x_T - x_0)), \tag{16}$$

we recover, of course, the constraint (7)

$${}_0D_T^{-(H+1/2)}\bar{\xi} = x_T - x_0. \tag{17}$$

Let us perform now variations in $\xi(t)$:

$$\delta_{\xi}\tilde{S}[\xi(t), \lambda] = \int_0^T \xi(t)\delta\xi(t) dt + \lambda_0 D_T^{-(H+1/2)}\delta\xi. \tag{18}$$

Writing explicitly the fractional integral

$$\delta_{\xi}\tilde{S}[\xi(t), \lambda] = \int_0^T \xi(t)\delta\xi(t) dt + \frac{\lambda}{\Gamma(H+1/2)} \int_0^T (T-t)^{H-1/2}\delta\xi(t) dt \tag{19}$$

$$= \int_0^T \left(\xi(t) + \frac{\lambda}{\Gamma(H+1/2)}(T-t)^{H-1/2} \right) \delta\xi(t) dt. \tag{20}$$

Since $\delta_{\xi}\tilde{S}[\xi(t), \lambda]$ must vanish for arbitrary $\delta\xi(t)$, we deduce

$$\bar{\xi}(t) + \frac{\lambda}{\Gamma(H+1/2)}(T-t)^{H-1/2} = 0. \tag{21}$$

The Lagrange multiplier is determined by applying ${}_0D_T^{-(H+1/2)}$ to equation (21) and using (17):

$$x_T - x_0 + \frac{\lambda}{\Gamma(H+1/2)^2} \int_0^T (T-t)^{2H-1} dt = 0. \tag{22}$$

Immediately, we find

$$x_T - x_0 + \frac{\lambda}{\Gamma^2(H+1/2)} \frac{T^{2H}}{2H} = 0 \tag{23}$$

and

$$\lambda = -2H\Gamma^2(H+1/2) \frac{x_T - x_0}{T^{2H}}. \tag{24}$$

Inserting this in (21)

$$\bar{\xi}(t) = 2H\Gamma(H+1/2) \frac{x_T - x_0}{T^{2H}}(T-t)^{H-1/2}. \tag{25}$$

Going back to (13) we straightforwardly obtain

$$G(x_T, T|x_0, 0) = f(T) \exp\left(-H\Gamma^2(H+1/2) \frac{(x_T - x_0)^2}{T^{2H}}\right). \tag{26}$$

And $f(T)$ is obtained from the normalization property, (14),

$$f(T) = \sqrt{\frac{H}{\pi}} \frac{\Gamma(H+1/2)}{T^H}. \tag{27}$$

Hence, finally,

$$G(x_T, T|x_0, 0) = \sqrt{\frac{H}{\pi}} \frac{\Gamma(H+1/2)}{T^H} \exp\left(-H\Gamma^2(H+1/2) \frac{(x_T - x_0)^2}{T^{2H}}\right). \tag{28}$$

2.1. The kinetic equation

Once the propagator is known, we can easily derive the kinetic equation associated with fBm. The Fourier transform of $G(x, t|x_0, 0)$ with respect to x is

$$\hat{G}(k, t) = \exp\left(-\frac{t^{2H}}{4H\Gamma^2(H+1/2)}k^2\right). \quad (29)$$

Differentiating with respect to t

$$\partial_t \hat{G}(k, t) = -\frac{t^{2H-1}}{2\Gamma^2(H+1/2)}k^2 \hat{G}(k, t). \quad (30)$$

Finally, we Fourier invert this expression to get the kinetic equation

$$\partial_t G(x, t) = \frac{t^{2H-1}}{2\Gamma^2(H+1/2)} \partial_x^2 G(x, t), \quad (31)$$

which is a diffusion equation with time-dependent diffusivity, as originally deduced in [15] from arguments based on the fluctuation–dissipation theorem. The invariance of (31) under the transformation $(x, t) \rightarrow (\mu^H x, \mu t)$ is manifest. Obviously, when $H = 1/2$ we retrieve the standard diffusion equation associated with Brownian motion

$$\partial_t G(x, t) = \frac{1}{2} \partial_x^2 G(x, t). \quad (32)$$

3. Conclusions

In this communication we have computed by path integral methods the well-known propagator of fBm for general Hurst exponent, extending the construction of [11], which was restricted to the subdiffusive case. The key technical point in our approach is to use the noise as path integral variable instead of the coordinate of the particle trajectory.

Acknowledgments

I C acknowledges the hospitality of Oak Ridge National Laboratory, where this work was carried out. Part of this research was sponsored by the Laboratory Research and Development Program of Oak Ridge National Laboratory, managed by UT-Battelle, LLC, for the US Department of Energy under contract number DE-AC05-00OR22725.

References

- [1] Langevin P 1908 *C. R. Acad. Sci., Paris* **146** 530–3
- [2] Uhlenbeck G E and Ornstein L S 1930 *Phys. Rev.* **36** 823–41
- [3] Metzler R and Klafter J 2004 *J. Phys. A: Math. Gen.* **37** R161–208
- [4] Molz F J, Liu H H and Szulga J 1997 *Water Resour. Res.* **33** 2273
- [5] Elliott R J and van der Hoek J 2001 Fractional Brownian motion and financial modeling *Trends in Mathematics* (Basel: Birkhäuser)
- [6] Sánchez R, Carreras B A, Newman D E, Lynch V E and van Milligen B Ph 2006 *Phys. Rev. E* **74** 016305
- [7] Mandelbrot B B and van Ness J W 1968 *SIAM Rev.* **10** 422
- [8] Samorodnitsky G and Taqqu M S 1994 *Stable Non-Gaussian Processes* (New York: Chapman and Hall)
- [9] Lutz E 2001 *Phys. Rev. E* **64** 051106
- [10] Hurst H E 1951 *Trans. Am. Soc. Civ. Eng.* **116** 770
- [11] Sebastian K L 1995 *J. Phys. A: Math. Gen.* **28** 4305
- [12] Oldham K and Spanier J 1974 *The Fractional Calculus* (New York: Academic)
- [13] Podlubny I 1998 *Fractional Differential Equations* (New York: Academic)
- [14] Feynman R P and Hibbs A R 1965 *Quantum Mechanics and Path Integrals* (New York: McGraw-Hill)
- [15] Wang K G and Lung C W 1990 *Phys. Lett. A* **151** 119