

q -difference equations of KdV type and Chazy-type second-degree difference equations

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys. A: Math. Theor. 41 332005

(<http://iopscience.iop.org/1751-8121/41/33/332005>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.150

The article was downloaded on 03/06/2010 at 07:06

Please note that [terms and conditions apply](#).

FAST TRACK COMMUNICATION

q-difference equations of KdV type and Chazy-type second-degree difference equations

Chris M Field¹, Nalini Joshi² and Frank W Nijhoff³¹ Korteweg-de Vries Institute for Mathematics, Universiteit van Amsterdam,
Plantage Muidergracht 24, 1018 TV Amsterdam, The Netherlands² School of Mathematics and Statistics F07, University of Sydney, NSW 2006, Australia³ School of Mathematics, University of Leeds, Leeds LS2 9JT, UKE-mail: cfield@science.uva.nl, nalini@maths.usyd.edu.au and nijhoff@maths.leeds.ac.uk

Received 16 May 2008, in final form 25 June 2008

Published 18 July 2008

Online at stacks.iop.org/JPhysA/41/332005**Abstract**

By imposing special compatible similarity constraints on a class of integrable partial *q*-difference equations of KdV-type we derive a hierarchy of second-degree ordinary *q*-difference equations. The lowest (non-trivial) member of this hierarchy is a second-order second-degree equation which can be considered as an analogue of equations in the class studied by Chazy. This second-order second-degree equation follows from a system in terms of two variables from which also follows an associated third-order first-degree equation. We present the isomonodromic deformation problem for the two-variable system and discuss the relation between the hierarchy of second-degree ordinary *q*-difference equations and other equations of Painlevé type.

PACS numbers: 02.30.Ik, 04.60.Nc, 05.45.-a

Mathematics Subject Classification: 34M55, 37J35, 37K10, 37K60, 39A13

1. Introduction

The construction and study of discrete Painlevé equations has been a topic of research interest for almost two decades [13, 21, 28, 35]. Reviews of the subject may be found in [14, 16]. The subject has culminated in the classification by Sakai of discrete as well as continuous Painlevé equations based on the algebraic geometry of the corresponding rational surfaces associated with the spaces of initial conditions [37]. As a byproduct of the latter treatment, a ‘mistress’ discrete Painlevé equation with elliptic dependence on the independent variable was discovered.

In the history of the Painlevé program, after the classification results for second-order first-degree equations, Painlevé’s students, Chazy and Garnier [6, 7, 15], investigated the classification of second-order second-degree equations and third-order first-degree equations.

The classification of the second-degree class was completed by Cosgrove in recent years [8, 9]. A partial classification for the third-order case was also obtained by the aforementioned authors. The work of Bureau [3, 4] is also important in this respect. No classification results exist for the analogous discrete case and hardly any examples of second-order second-degree difference equations exist to date, with the notable exception of an (additive difference) equation given by Estévez and Clarkson [10].

A key result of this communication is a second-order second-degree equation, which can be considered as a q -analogue of an equation in the Chazy–Cosgrove class. This new equation contains four free parameters, which suggests that it could be a q -difference analogue of the second-order second-degree differential equation that is a counterpart of the sixth Painlevé equation. There are several forms of a second-order second-degree equation related to the sixth Painlevé equation that have appeared in the literature, notably one derived by Fokas and Ablowitz [12] and another appearing in the work of Okamoto [32]. Difference analogues of the Fokas–Ablowitz equation have been provided by Grammaticos and Ramani [34], but these difference equations were of first degree.

The second-order second-degree q -difference equation follows from a system in terms of two variables, from which a third-order first-degree equation also follows. For the system in terms of two variables we establish integrability through a Lax pair in the form of an isomonodromic deformation system. Furthermore, we show that the equation arises as a similarity reduction from an integrable partial q -difference equation. Higher-order members of this hierarchy also exist. We show explicitly through the same procedure the construction of higher-order second-degree equations, which form a hierarchy associated with the new equation.

2. q -difference similarity reduction

Lattice equations of KdV-type were introduced and studied over the last three decades [19, 29], see [27] for a review. These lattice equations can be formulated as partial difference equations on a lattice with step sizes that enter as parameters of the equation. Conventionally we think of these parameters as fixed constants. However, in agreement with the integrability of these equations, there exists the freedom to take the parameters as functions of the local lattice coordinate in each corresponding direction. In this communication we consider the case when the parameters depend exponentially with base q on the lattice coordinates.

We work in a space \mathcal{F} of functions f of arbitrarily many variables a_i ($i = 1, \dots, M$ for any M) on which we define the q -shift operations

$${}_qT_i f(a_1, \dots, a_N) := f(a_1, \dots, q a_i, \dots, a_N).$$

For $u, v, z \in \mathcal{F}$, we consider the following systems of nonlinear partial q -difference equations:

$$(u - {}_qT_i {}_qT_j u)({}_qT_j u - {}_qT_i u) = (a_i^2 - a_j^2)q^2, \quad (2.1)$$

$$a_j({}_qT_j v) {}_qT_i {}_qT_j v + a_i({}_qT_j v)v = a_i({}_qT_i v) {}_qT_j {}_qT_i v + a_j({}_qT_i v)v \quad (2.2)$$

and

$$a_i^2(z - {}_qT_i z)({}_qT_j z - {}_qT_i {}_qT_j z) = a_j^2(z - {}_qT_j z)({}_qT_i z - {}_qT_i {}_qT_j z), \quad (2.3)$$

where $i, j = 1, \dots, M$. Each of these systems (2.1)–(2.3) represents a multi-dimensionally consistent family of partial difference equations, in the sense of [1, 31], which implies that they constitute holonomic systems of nonlinear partial q -difference equations. Another way to formulate this property is through an underlying linear system which takes the form

$${}_qT_i^{-1} \phi = M_i(k)\phi, \quad (2.4)$$

where $\phi = \phi(k; \{a_j\})$ is a two-component vector-valued function and by consistency, ${}_q T_i^{-1} {}_q T_j^{-1} \phi = {}_q T_j^{-1} {}_q T_i^{-1} \phi$ leads to the set of Lax equations (for each pair of indices i, j)

$$({}_q T_i^{-1} M_j) M_i = ({}_q T_j^{-1} M_i) M_j. \tag{2.5}$$

We will consider three different cases, associated respectively with equations (2.1)–(2.3). To avoid proliferation of symbols we use the same symbol $M_i(k)$ for each of the respective Lax matrices. For specific choices of the matrices M_i the Lax equations (2.5) lead to the nonlinear equations given above. In the case of the q -lattice KdV (2.1), the Lax matrices M_i are given by

$$M_i(k; \{a_j\}) = \frac{1}{a_i - k} \begin{pmatrix} a_i - {}_q T_i^{-1} u, & 1 \\ k^2 - a_i^2 + (a_i + u)(a_i - {}_q T_i^{-1} u), & a_i + u \end{pmatrix}. \tag{2.6}$$

In the case of the q -lattice mKdV (2.2), the Lax matrices M_i are given by

$$M_i(k; \{a_j\}) = \frac{1}{a_i - k} \begin{pmatrix} a_i ({}_q T_i^{-1} v) / v, & k^2 / v \\ {}_q T_i^{-1} v, & a_i \end{pmatrix}. \tag{2.7}$$

Finally, in the case of the q -lattice SKdV (2.3), the Lax matrices M_i are given by

$$M_i(k; \{a_j\}) = \frac{a_i}{a_i - k} \begin{pmatrix} 1, & (k^2 / a_i^2) (z - {}_q T_i^{-1} z)^{-1} \\ z - {}_q T_i^{-1} z, & 1 \end{pmatrix}. \tag{2.8}$$

These Lax matrices are straightforward generalizations of those with constant lattice parameters given in e.g. [30].

We mention that the solutions of the equations (2.1)–(2.3) are related through discrete Miura-type relations, namely

$$a_i (z - {}_q T_i^{-1} z) = v ({}_q T_i^{-1} v), \tag{2.9a}$$

$$s = (a_i - {}_q T_i^{-1} u) v - a_i {}_q T_i^{-1} v, \tag{2.9b}$$

$${}_q T_i^{-1} s = a_i v - (a_i + u) {}_q T_i^{-1} v, \tag{2.9c}$$

where $s \in \mathcal{F}$ is an auxiliary dependent variable. From these relations, the partial q -difference equations (2.1)–(2.3) can be derived by eliminating s .

Similarity reductions of lattice equations have been considered in [25, 26, 30, 28, 31] where it was shown that scaling invariance of the solution can be implemented through additional compatible constraints on the lattice equations. In the present case of (2.1)–(2.3) these constraints adopt the following form [11]

$$u(\{q^{-N} a_i\}) = q^{-N} \frac{1 - \lambda(q^N - 1)(-1)^{\sum_i {}^q \log a_i}}{1 + \lambda(q^N - 1)(-1)^{\sum_i {}^q \log a_i}} u(\{a_j\}), \tag{2.10a}$$

$$v(\{q^{-N} a_i\}) = \frac{1 - \lambda(q^N - 1)(-1)^{\sum_i {}^q \log a_i}}{1 + \mu(q^N - 1)} v(\{a_j\}), \tag{2.10b}$$

$$z(\{q^{-N} a_i\}) = q^N \frac{1 - \mu(q^N - 1)}{1 + \mu(q^N - 1)} z(\{a_j\}), \tag{2.10c}$$

where λ and μ are constant parameters of the reduction and $N \in \mathbb{N}$ represents a ‘periodicity freedom’. The notation ${}^q \log x$ refers to the logarithm of x with base q .

In order to compute the corresponding isomonodromic deformation problems associated with the similarity reductions we have the following constraints on the vector function of the Lax pairs. In the case of (2.1) we have

$$\begin{aligned} \phi(q^N k; \{a_j\}) &= \begin{pmatrix} (1 + \lambda(q^N - 1)(-1)^{\sum_i q^{\log a_i}}, & 0 \\ -2\lambda q^{\frac{q^N - 1}{q - 1}} (\sum_i a_i) (-1)^{\sum_i q^{\log a_i}}, & q^N (1 - \lambda(q^N - 1)(-1)^{\sum_i q^{\log a_i}}) \end{pmatrix} \phi(k; \{q^{-N} a_j\}). \end{aligned} \tag{2.11}$$

In the case of (2.2)

$$\begin{aligned} \phi(q^N k; \{a_j\}) &= \begin{pmatrix} (1 + \lambda(q^N - 1)(-1)^{\sum_i q^{\log a_i}} & 0 \\ 0 & q^{-N} (1 + \mu(q^N - 1)) \end{pmatrix} \phi(k; \{q^{-N} a_j\}). \end{aligned} \tag{2.12}$$

In the case of (2.3)

$$\begin{aligned} \phi(q^N k; \{a_j\}) &= \begin{pmatrix} (1 - \mu(q^N - 1)) & 0 \\ 0 & q^{-N} (1 + \mu(q^N - 1)) \end{pmatrix} \phi(k; \{q^{-N} a_j\}). \end{aligned} \tag{2.13}$$

The similarity constraints (2.11)–(2.13), in conjunction with the discrete linear equations (2.6)–(2.8) can be used to derive corresponding q -isomonodromic deformation problems. That is (2.11)–(2.13) lead to q -difference equations in the spectral variable k , hence together with the lattice equation Lax pairs we obtain q -isomonodromic deformation problems for the corresponding reductions.

Remarks:

- (i) The similarity constraints above were obtained through an approach based on Jackson-type integrals, the details of which will be presented elsewhere [11]. By construction, these constraints are compatible with the underlying lattice equations, which can be checked *a posteriori* by an explicit calculation, presented in the appendix.
- (ii) In this approach, the dynamics in terms of the variables a_i appear through appropriately chosen q -analogues of exponential functions, whereas the relevant Jackson integrals exhibit an invariance through scaling by factors q^N .
- (iii) The parameters λ and μ arise in this setting through boundary contributions in a manner analogous to the derivation in [30].

In the remainder of this communication our aim is to implement the similarity constraint to obtain explicit reductions to ordinary q -difference equations. For simplicity we consider only the reduction of the q -mKdV equation (2.2), leaving considerations of the q -KdV and q -SKdV to a future publication [11]. There are two possible scenarios to derive similarity reductions of the lattice equations using the constraint (2.10b).

‘Periodic’ similarity reduction. By fixing $M = 2$ and allowing N to vary, we select two lattice directions, say the variables a_1 and a_2 , and consider similarity reductions with different values of N . This is a q -variant of the periodic staircase-type reduction of partial difference equations on the two-dimensional lattice. For instance, with $N = 2$ the reduction is a second-order first-degree q -Painlevé equation. Increasing N leads to q -difference Painlevé type-equations of increasing order. However, we will not pursue this route here but leave it to a subsequent publication [11]. These reductions are reminiscent of the work [17, 18, 36].

Multi-variable similarity reduction. The similarity constraints provide the mechanism to couple together two or more lattice directions. By considering the case $N = 1$ we implement

the similarity constraints on an extended lattice of three or more dimensions in order to obtain coupled ordinary q -difference equations, in a way that is reminiscent of the approach of [31]. This is considered in the next section.

We have not considered the more general case of arbitrary $M, N \in \mathbb{N}$, which we will postpone to a future publication [11].

3. Multi-variable similarity reduction

In this section we consider explicitly the $M = 1, 2$ and 3 cases for $N = 1$.

For simplicity we shall in what follows denote the coefficient in (2.10b) as γ , i.e.

$$\gamma = \frac{1 - \lambda(q - 1)(-1)^{\sum_i q \log a_i}}{1 + \mu(q - 1)} \Rightarrow v(\{q^{-1}a_i\}) = \gamma v(\{a_i\}), \quad (3.1)$$

where γ alternates between two values, i.e., ${}_q T_i^2 \gamma = \gamma$.

In contrast to the usual difference case which was explored in [31] where in the case of two variables we obtain a nontrivial OΔE as a reduction, in the q -difference case we have to consider at least three independent variables to obtain a nontrivial system of OΔEs as a reduction.

In [31] the compatibility between the similarity constraint and the lattice system was established and led to a system of higher order difference equations in the reduction, namely equations which were on the level of the first Garnier system. In contrast to the $q = 1$ work, the 3D similarity constraint here is somewhat simpler and leads to a second-order equation (which is of second-degree, and is a principal result of this communication).

3.1. Two-variable case

Let us now select among the collection of variables $\{a_j\}$ two specific ones which for simplicity we will call a and b . Denote the q -shifts in these variables by an over-tilde $\tilde{}$ and an over-hat $\hat{}$ respectively. Equation (2.2) may now be written as

$$b\hat{v}\hat{v} + a\hat{v}v = a\tilde{v}\tilde{v} + b\tilde{v}\tilde{v}, \quad (3.2)$$

where the over-tilde $\tilde{}$ refers to the q -translation $a \mapsto qa$ and the over-hat $\hat{}$ refers to the q -translation $b \mapsto qb$ (so if $v \equiv v(a, b)$, $\tilde{v} \equiv v(qa, b)$, $\hat{v} \equiv v(a, bq)$, ...).

Equation (3.1) gives the constraint $v = \gamma\hat{v}$ to impose on (3.2) (where $\hat{\tilde{v}} = \tilde{\hat{v}} = \gamma$). This leads to the linear first-order (in that it is a two-point) ordinary difference equation

$$\hat{v} = C\tilde{v}, \quad (3.3)$$

where $C = \tilde{\gamma}(a\gamma + b)/(a + b\gamma)$. In the appendix the consistency between the lattice equation (3.2) and the constraint (3.3) is shown by direct computation.

3.2. Three-variable case

Take three copies of the lattice mKdV equation with $a_1 = a, a_2 = b, a_3 = c$,

$$b\hat{v}\hat{v} + a\hat{v}v = a\tilde{v}\tilde{v} + b\tilde{v}\tilde{v}, \quad (3.4a)$$

$$c\hat{v}\hat{v} + a\hat{v}v = a\tilde{v}\tilde{v} + c\tilde{v}\tilde{v}, \quad (3.4b)$$

$$c\hat{v}\hat{v} + b\hat{v}v = b\tilde{v}\tilde{v} + c\tilde{v}\tilde{v}, \quad (3.4c)$$

where $\bar{c} = qc$, together with the constraint

$$v(q^{-1}a, q^{-1}b, q^{-1}c) = \gamma v(a, b, c). \tag{3.5}$$

(The similarity constraint is shown by a direct computation to be compatible with the multidimensionally consistent system of mKdV lattice equations in the appendix.) We now proceed to derive the reduced system which leads to a (higher-degree) ordinary q -difference equation in terms of one selected independent variable, say the variable a . The remaining variables b and c will play the role of parameters in the reduced equation. Thus, we can derive the following system of two coupled OΔEs for $v(a, b, c)$ and $w(a, b, c) \equiv v(a, b, q^{-1}c)$:

$$\gamma v = w \frac{a\tilde{\gamma}\tilde{v} - b\tilde{w}}{a\tilde{w} - b\tilde{\gamma}\tilde{v}}, \tag{3.6a}$$

$$\tilde{w} = v \frac{aw - c\tilde{v}}{a\tilde{v} - cw}, \tag{3.6b}$$

where $\tilde{\gamma} = \gamma$. We consider the system (3.6a) and (3.6b) to constitute a q -Painlevé system with four free parameters.

The system (3.6a) and (3.6b) can be reduced to a second-order second-degree ordinary difference equation as follows. Introduce the variables

$$X = \frac{v}{w}, \quad V = \frac{\tilde{v}}{v}, \quad W = \frac{\tilde{w}}{w}, \tag{3.7}$$

then from (3.6a) we obtain

$$\tilde{\gamma} \frac{\tilde{v}}{w} = \tilde{\gamma} V X W = \frac{a\gamma X + b}{b\gamma X + a}, \tag{3.8}$$

whereas from (3.6b) we get

$$W = \frac{VX}{\tilde{X}} = \frac{a + q^{-1}cXV}{q^{-1}c/X + aV}, \tag{3.9}$$

using also the definitions (3.7). Thus, we obtain a quadratic equation for V in terms of X and \tilde{X} and hence also we have W in terms of X and \tilde{X} . Inserting these into (3.8) we obtain a second-order algebraic equation for X . Alternatively, avoiding the emergence of square roots, the following second-order second-degree equation for X may be derived:

$$\begin{aligned} & \left[\tilde{\gamma}^2 \tilde{X} \tilde{X} - \left(\frac{a\gamma X + b}{b\gamma X + a} \right)^2 \right]^2 \\ &= \tilde{\gamma} \frac{c^2}{a^2} \frac{1}{X} \left(\frac{a\gamma X + b}{b\gamma X + a} \right) \left[\tilde{\gamma} \tilde{X} (1 - X\tilde{X}) + q^{-1} (1 - X\tilde{X}) \frac{a\gamma X + b}{b\gamma X + a} \right] \\ & \times \left[q^{-1} \tilde{\gamma} \tilde{X} (1 - X\tilde{X}) + (1 - X\tilde{X}) \frac{a\gamma X + b}{b\gamma X + a} \right]. \end{aligned} \tag{3.10}$$

We consider this second-degree equation to be one of the main results of this communication.

The following third-order (four-point) ordinary non-autonomous difference equation of first-degree also follows directly from the system (3.8) and (3.9),

$$\begin{aligned} \Delta \frac{qa\tilde{\gamma}\tilde{X} + b}{b\tilde{\gamma}\tilde{X} + qa} & \left[q^{-1}\gamma X(1 - \tilde{X}\tilde{X}) + (1 - X\tilde{X}) \frac{qa\tilde{\gamma}\tilde{X} + b}{b\tilde{\gamma}\tilde{X} + qa} \right] \\ &= q\tilde{\Delta} \frac{a\gamma X + b}{b\gamma X + a} \left[\tilde{\gamma} \tilde{X} (1 - X\tilde{X}) + q^{-1} (1 - X\tilde{X}) \frac{a\gamma X + b}{b\gamma X + a} \right], \end{aligned} \tag{3.11}$$

where Δ is given by

$$\Delta = \tilde{\gamma}^2 \tilde{X} \tilde{X} - \left(\frac{a\gamma X + b}{b\gamma X + a} \right)^2.$$

This equation may help to fix the ‘proper’ branch of the multivalued second-degree correspondence (3.10).

We now proceed to present the Lax pair for the q -Painlevé system (3.6a) and (3.6b) and the system (3.8) and (3.9). The Lax pair is formed by considering the compatibility of two paths on the lattice: along a ‘period’ then in the a direction and evolving in the a direction then along a ‘period’. Using (2.12) the evolution along a period is converted into a dilation of the spectral parameter, k , by q . The result is the following isomonodromic q -difference system for the vector $\phi(k; a)$ which using the results of section 2 yields

$$\phi(k; q^{-1}a) = M(k; a)\phi(k; a), \tag{3.12a}$$

$$\phi(qk; a) = L(k; a)\phi(k; a), \tag{3.12b}$$

where

$$M(k; a) = \frac{1}{a - k} \begin{pmatrix} a\underline{v}/v & k^2/v \\ \underline{v} & a \end{pmatrix}, \tag{3.13a}$$

and

$$L(k; a) = \frac{1}{a - k} \begin{pmatrix} a\gamma v/\tilde{v} & k^2/\tilde{v} \\ q^{-1}\gamma v & q^{-1}a \end{pmatrix} \begin{pmatrix} b\tilde{\gamma}\tilde{v}/w & k^2/w \\ \tilde{\gamma}\tilde{v} & b \end{pmatrix} \begin{pmatrix} cw/v & k^2/v \\ w & c \end{pmatrix} \tag{3.13b}$$

where we have suppressed the dependence on the variables b and c (which now play the role of parameters) and omitted the unnecessary prefactors $(b - k)^{-1}$ and $(c - k)^{-1}$, as well as an over factor $q^{-1}(1 + \mu(q - 1))$.

The consistency condition obtained from the two ways of expressing $\phi(qk; q^{-1}a)$ in terms of $\phi(k; a)$ is formed by the Lax equation

$$L(k; q^{-1}a)M(k; a) = M(qk; a)L(k; a). \tag{3.14}$$

A gauge transformation can be obtained expressing the Lax matrices in terms of the variables introduced in (3.7). Setting

$$\mathcal{M}(k; a) = \frac{1}{a - k} \begin{pmatrix} a/\underline{Y} & k^2 \\ 1 & a\underline{Y} \end{pmatrix}, \tag{3.15a}$$

$$\mathcal{L}(k; a) = \frac{1}{a - k} \begin{pmatrix} \tilde{\gamma}(ab\gamma X + k^2) & k^2(a\gamma X + b)/V \\ q^{-1}\tilde{\gamma}V(a + b\gamma X) & q^{-1}(ab + k^2\gamma X) \end{pmatrix} \begin{pmatrix} c/X & k^2 \\ 1/X & c \end{pmatrix}, \tag{3.15b}$$

the Lax equations (3.14) (replacing L and M by \mathcal{L} and \mathcal{M} respectively) yield a set of relations equivalent to the following two equations:

$$\tilde{\gamma}V\underline{Y}\tilde{X} = \frac{a\gamma X + b}{a + b\gamma X}, \tag{3.16}$$

$$aV^2 + q^{-1}c \left(\frac{1}{X} - \tilde{X} \right) V - a\frac{\tilde{X}}{X} = 0, \tag{3.17}$$

using also $\tilde{\gamma} = \underline{\gamma}$. This set follows directly from (3.8) and (3.9).

3.3. Four-variable case

Suppose we have four variables $a_i, i = 1, \dots, 4$. Select $a = a_1$ to be the independent variable after reduction. Introduce the dependent variables $w_{j-2} = {}_qT_j^{-1}v, j = 3, 4$. Then directly from the q -lattice mKdV equation (2.2) we have the set of equations

$$w_j = v \frac{aw_j - a_{j+2}\tilde{v}}{a\tilde{v} - a_{j+2}w_j}, \quad j = 1, 2, \tag{3.18}$$

where as before the tilde denotes a q -shift in the variable a . At the same time the multiply shifted object ${}_qT_3^{-1}{}_qT_4^{-1}\tilde{v}$ can be expressed in a unique way (due to the CAC property) in terms of \tilde{v} and ${}_qT_j^{-1}v = w_{j-2}, j = 3, 4$, by iterating the relevant copies of the q -lattice mKdV equation in the variables $a_j, j \neq 2$, leading to an expression of the form ${}_qT_3^{-1}{}_qT_4^{-1}\tilde{v} =: F(\tilde{v}, w_1, w_2)$, where F is easily obtained explicitly. Imposing the similarity constraint (3.1) we obtain $\tilde{\gamma}{}_qT_2v = F(\tilde{v}, w_1, w_2)$ and inserting this expression into the q -lattice mKdV (2.2) with $i = 1, j = 2$ we obtain

$$\left(a + \frac{a_2 a_3 w_2 - a_4 w_1}{\gamma a_3 w_1 - a_4 w_2} \right) (a_2 \tilde{\gamma}^{-1} F(\tilde{v}, w_1, w_2) - a\tilde{v}) = (a_2^2 - a^2) v \tilde{v}. \tag{3.19}$$

With the explicit form of $F(\tilde{v}, w_1, w_2)$ equation (3.19) reads

$$\begin{aligned} & (a_2^2 - a^2) \gamma \tilde{\gamma} \tilde{v} (a_3 w_1 - a_4 w_2) [a(a_3^2 - a_4^2)\tilde{v} + a_3(a_4^2 - a^2)w_1 + a_4(a^2 - a_3^2)w_2] \\ &= [(a_2 a_3 - \gamma a a_4)w_2 + (\gamma a a_3 - a_2 a_4)w_1] [a(a_3^2 - a_4^2)(a_2 w_1 w_2 - \tilde{\gamma} a \tilde{v} \tilde{v}) \\ &+ (a_2 a_4(a^2 - a_3^2)\tilde{v} - a a_3(a_4^2 - a^2)\tilde{\gamma} \tilde{v})w_1 \\ &+ (a_2 a_3(a_4^2 - a^2)\tilde{v} - \tilde{\gamma} a a_4(a^2 - a_3^2)\tilde{v})w_2], \end{aligned} \tag{3.20a}$$

and this is supplemented by the two equations

$$avw_1 + a_3w_1w_1 = a_3v\tilde{v} + a\tilde{v}w_1, \tag{3.20b}$$

$$avw_2 + a_4w_2w_2 = a_4v\tilde{v} + a\tilde{v}w_2, \tag{3.20c}$$

which is equivalent to a five-point (fourth-order) q -difference equation in terms of v alone, containing five free parameters: a_2, a_3, a_4, λ and μ (inside γ and $\tilde{\gamma}$). This would be an algebraic equation, so we proceed as follows in order to derive a higher-degree q -difference system. Introduce the variables

$$X_i = \frac{v}{w_i}, \quad W_i = \frac{\tilde{w}_i}{w_i}, \quad i = 1, 2, \tag{3.21}$$

while retaining the variable $V = \tilde{v}/v$ as before. By definition we have

$$\frac{V}{W_i} = \frac{\tilde{X}_i}{X_i}, \quad i = 1, 2, \tag{3.22}$$

and from (3.20b) and (3.20c) we obtain

$$W_i = \frac{qa + a_{i+2}VX_i}{qaV + a_{i+2}/X_i} = \frac{VX_i}{\tilde{X}_i}, \quad i = 1, 2, \tag{3.23}$$

whilst from (3.20a) we get

$$\begin{aligned}
 (a_2^2 - a^2)\gamma\tilde{\gamma}V & \left(\frac{a_3}{X_1} - \frac{a_4}{X_2} \right) \left(a \frac{a_3^2 - a_4^2}{\tilde{Y}} + a_3 \frac{a_4^2 - a^2}{X_1} + a_4 \frac{a^2 - a_3^2}{X_2} \right) \\
 & = \left(\frac{a_2 a_3 - \gamma a a_4}{X_2} + \frac{\gamma a a_3 - a_2 a_4}{X_1} \right) \left[a (a_3^2 - a_4^2) \left(\frac{a_2}{X_1 X_2} - \tilde{\gamma} a \frac{V}{\tilde{Y}} \right) \right. \\
 & \quad + \left(a_2 a_4 \frac{a^2 - a_3^2}{\tilde{Y}} - a a_3 (a_4^2 - a^2) \tilde{\gamma} V \right) \frac{1}{X_1} \\
 & \quad \left. + \left(a_2 a_3 \frac{a_4^2 - a^2}{\tilde{Y}} - \tilde{\gamma} a a_4 (a^2 - a_3^2) V \right) \frac{1}{X_2} \right]. \tag{3.24}
 \end{aligned}$$

From (3.23) we obtain the set of quadratic equations for V ,

$$qa \frac{X_i}{\tilde{X}_i} V^2 + a_{i+2} \left(\frac{1}{\tilde{X}_i} - X_i \right) V - qa = 0, \quad i = 1, 2, \tag{3.25}$$

from which by eliminating V we obtain

$$\begin{aligned}
 & [a_3(1 - X_1 \tilde{X}_1)X_2 - a_4(1 - X_2 \tilde{X}_2)X_1] [a_3(1 - X_1 \tilde{X}_1)\tilde{X}_2 - a_4(1 - X_2 \tilde{X}_2)\tilde{X}_1] \\
 & = q^2 a^2 (X_1 \tilde{X}_2 - X_2 \tilde{X}_1)^2. \tag{3.26}
 \end{aligned}$$

Furthermore, solving V from the quadratic system as

$$V = qa \frac{X_2 \tilde{X}_1 - X_1 \tilde{X}_2}{a_3(1 - X_1 \tilde{X}_1)X_2 - a_4(1 - X_2 \tilde{X}_2)X_1}, \tag{3.27}$$

and inserting this into (3.24) we obtain a second-order equation in both X_1, X_2 coupled to the equation (3.26) which is first order in both X_1, X_2 . It is this coupled system of two equations in X_1, X_2 which forms our higher order generalization of (3.10). The system of (3.24) and (3.26) with (3.27) constitutes a third-order system with five parameters.

The derivation of the Lax pair follows the same approach as that for the three-variable case (with an extra factor in \mathcal{L} due to the additional lattice direction). We omit details here, which we intend to publish in the future [11].

3.4. Beyond the four-variable case

It is straight forward to give the form of the full hierarchy, however due to lack of space we postpone this until a later publication [11].

4. Conclusion and discussion

In this communication we have presented the results of a scheme to derive partial q -difference equations of KdV type and consistent symmetries of the equations and demonstrated how it can be implemented. Lax matrices follow from this approach. A notable result is the derivation of the higher-degree equation (3.10), showing that the scheme presented here allows for the derivation of new results within the field of discrete integrable systems.

The first-, second- and third-order members of the $N = 1$ hierarchy have been presented. The scheme continues to give successively higher-order equations by considering successively higher dimensions of the original lattice equation. One may ask the natural question as to whether this gives an ‘interpolating’ hierarchy which, contrary to the usual cases, increases

the order and number of parameters of the equations by one in each step, rather than a two-step increase. A further natural question connected with this hierarchy is its relation to the q -Garnier systems of Sakai [38].

We will present full details of the scheme from which the lattice equations (2.1)–(2.3) and their associated constraints follow in a future publication [11]. There we will consider the most general case of symmetry reductions (arbitrary $N \in \mathbb{N}$) of all three lattice equations.

We also intend to return in a future publication to the question of limits and degeneracies of the equations presented in this paper. These include the $q \rightarrow 1$ continuum limit, the $q \rightarrow 1$ discrete limit and the $q \rightarrow 0$ crystal or ultradiscrete limit.

Acknowledgments

During the writing of this communication C M Field has been supported by the Australian Research Council Discovery Project Grant No. DP0664624 and by the Netherlands Organization for Scientific Research (NWO) in the VIDI-project ‘Symmetry and modularity in exactly solvable models’. F W Nijhoff would like to thank A Ramani for useful comments. N Joshi is also supported by the Australian Research Council Discovery Project Grant No. DP0664624. Part of the work presented here was performed whilst N Joshi was visiting the University of Leeds.

Appendix

In this appendix we present the result of explicit calculations showing the consistency of the lattice equations and constraints.

A.1. Two-variable consistency

We shall check the consistency between the lattice equation (3.2) and the constraint $v = \gamma \widehat{v}$ by direct computation. This computation is illustrated in figure A1. Assuming the values v_0, v_1 as indicated in figure A1 are given, we compute successively v_{12}, v_2 etc, where the subscripts refer to the shifts in the lattice variables a, b respectively, as is evident from figure A1. Points other than v_0 and v_1 are computed using either the lattice equation (indicated by \times) or by using the similarity constraint (indicated by \circ). The value $v_{-1,-2}$ is the first point which can be calculated in two different ways (hence indicated in the diagram by \otimes). Without making any particular assumptions on how γ depends on a and b , a straightforward calculation shows that the two ways of computing $v_{-1,-2}$ are indeed the same, for any choice of initial data v_0 and v_1 , provided that γ obeys the relation

$$\left(\frac{a+b\gamma}{b+a\gamma}\right)^{\widehat{\gamma}} \left(\frac{a+b\gamma}{b+a\gamma}\right)^{-1} = \frac{\widetilde{\gamma}}{\widehat{\gamma}}. \tag{A.1}$$

A particular solution of this relation is

$$\widehat{\gamma} = \gamma \quad \Leftrightarrow \quad \widehat{\gamma} = \widetilde{\gamma}, \tag{A.2}$$

and hence $\widetilde{\gamma} = \gamma$ implying that γ is an alternating ‘constant’ which is in accordance with the value given in (2.10b). The reduced equation in this case is (3.3), which can be readily integrated.

More generally, equation (A.1) can be resolved by setting

$$\frac{a+b\gamma}{b+a\gamma} = \frac{\widetilde{v}}{\widehat{v}}, \quad \gamma = \frac{\widehat{v}}{v}, \tag{A.3}$$




Figure A1. Consistency on the 2D lattice.




Figure A2. Consistency on the 3D lattice.

leading to the consequence that v has to solve the q -lattice mKdV (3.2). In principle we could take for v any solution of the reduced equation (3.3) and use this to parametrize the reduced equation for v via the relations (A.3). In any event, we see that the two-variable case does not lead to interesting nonlinear equations.

A.2. Three-variable consistency

In this case the consistency diagram is shown in figure A2. A similar notation as the previous case is used as is evident from figure A2. The initial data v_0, v_1 and v_2 are given, and the indicated values on the vertices are computed either by using one of the lattice equations (3.4a)–(3.4c) or the similarity constraint (3.5) over the diagonal. Thus, $v_{1,2}$ is obtained from (3.4a) yielding

$$v_{1,2} = v_0 \frac{av_2 - bv_1}{av_1 - bv_2},$$

whilst from the similarity constraint we obtain

$$v_{-1,-3} = \gamma_2 v_2, \quad v_{-3} = \gamma_{1,2} v_{1,2}, \quad v_{-2,-3} = \gamma_1 v_1,$$

assuming that γ shifts along the lattice, indicated by the indices, and finally the value of $v_{-1,-2,-3}$ can be computed in two different ways, leading to

$$v_{-1,-2,-3} = \gamma_0 v_0 = \frac{av_{-2,-3} - bv_{-1,-3}}{av_{-1,-3} - bv_{-2,-3}} v_{-3} = \frac{a\gamma_1 v_1 - b\gamma_2 v_2}{a\gamma_2 v_2 - b\gamma_1 v_1} \gamma_{1,2} v_0 \frac{av_2 - bv_1}{av_1 - bv_2},$$

leading quadratic identity in v_1 and v_2 . Assuming that the latter must hold identically, and thus setting all coefficients equal to zero, we obtain the following conditions on γ :

$$\gamma_{1,2,3} = \gamma_1 = \gamma_2 = \gamma_3,$$

from which we conclude that γ is an alternating ‘constant’, for instance

$$\gamma = \alpha\beta^{(-1)^{n+m+\dots}} \quad (\alpha, \beta \text{ constants}) \quad (\text{A.4})$$

and this leads to the conditions from which it is easily deduced that the form (3.1) of γ satisfies these conditions.

References

- [1] Bobenko A and Suris Y 2002 Integrable systems on quad-graphs *Int. Math. Res. Not.* **11** 573–611
- [2] Birkhoff G D 1913 The generalized Riemann problem for linear differential equations and the allied problems for linear difference and q -difference equations *Proc. Am. Acad. Sci.* **49** 521–68
- [3] Bureau F J 1972 Équations différentielles du second ordre en Y et du second degré en \check{Y} dont l’intégrale générale est à points critiques fixes *Ann. Mat. Pura Appl.* **91** 163–281
- [4] Bureau F J 1987 Sur des systèmes différentiels non linéaires du troisième ordre et les équations différentielles non linéaires associées *Acad. R. Belg. Bull. Cl. Sci.* **73** 335–53
- [5] Capel H W, Nijhoff F W and Papageorgiou V G 1991 Complete integrability of Lagrangian mappings and lattices of KdV type *Phys. Lett. A* **155** 377–87
- [6] Chazy J 1909 Sur les équations différentielles du second ordre à points critiques fixes *C. R. Acad. Sci., Paris* **148** 1381–4
- [7] Chazy J 1911 Sur les équations différentielles du troisième ordre et ordre supérieur dont l’intégrale générale a ses points critiques fixes *Acta Math.* **34** 317–85
- [8] Cosgrove C 2006 Chazy’s second-degree Painlevé equations *J. Phys. A: Math. Gen.* **39** 11955–71
- [9] Cosgrove C M and Scoufis G 1993 Painlevé classification of a class of differential equations of the second order and second degree *Stud. Appl. Math.* **88** 25–87
- [10] Estévez P G and Clarkson P A 2000 Discrete equations and the singular manifold method *SIDE III—Symmetries and Integrability of Difference Equations (CRM Proceedings and Lecture Notes vol 25)* ed D Levi and O Ragnisco (Providence, RI: American Mathematical Society) pp 139–46
- [11] Field C M, Joshi N and Nijhoff F W, in preparation
- [12] Fokas A S and Ablowitz M J 1982 A unified approach to transformations and elementary solutions of Painlevé equations *J. Math. Phys.* **23** 2033–42
- [13] Fokas A S, Its A R and Kitaev A V 1991 Discrete Painlevé equations and their appearance in quantum gravity *Commun. Math. Phys.* **142** 313–44
- [14] Grammaticos B, Nijhoff F W and Ramani A 1999 Discrete Painlevé equations *The Painlevé Property: One Century Later* ed R Conte (New York: Springer) pp 413–516
- [15] Garnier R 1912 Sur des équations différentielles du troisième ordre dont l’intégrale générale est uniforme et sur une classe d’équations nouvelles d’ordre supérieur *Ann. Écol. Norm. Sup.* **29** 1–126
- [16] Grammaticos B and Ramani A 2004 Discrete Painlevé equations: a review *Discrete Integrable Systems (Lect. Notes Phys. vol 644)* ed B Grammaticos, Y Kossmann-Schwarzbach and T Tamizhmani (New York: Springer) pp 245–321
- [17] Grammaticos B, Ramani A, Satsuma J, Willox R and Carstea A 2005 Reductions of integrable lattices *J. Nonlinear Math. Phys.* **12** 363–71
- [18] Hay M, Hietarinta J, Joshi N and Nijhoff F W 2007 A Lax pair for a lattice modified KdV equation, reductions to q -Painlevé equations and associated Lax pairs *J. Phys. A: Math. Theor.* **40** F61–F73
- [19] Hirota R 1977 Nonlinear partial difference equations: I. A difference analogue of the Korteweg–de Vries equation *J. Phys. Soc. Japan* **43** 1424–33
- [20] Hydon P and Rasin O 2007 Symmetries of integrable difference equations on the quad-graph *Stud. Appl. Math.* **119** 253–69
- [21] Jimbo M and Sakai H 1996 A q -analog of the sixth Painlevé equation *Lett. Math. Phys.* **38** 145–54
- [22] Kajiwara K, Noumi M and Yamada Y 2001 A study on the fourth q -Painlevé equation *J. Phys. A: Math. Gen.* **34** 8563–81
- [23] Kajiwara K, Noumi M and Yamada Y 2002 Discrete dynamical systems with $W(A_{m-1}^{(1)} \times A_{n-1}^{(1)})$ symmetry *Lett. Math. Phys.* **60** 211–9
- [24] Kajiwara K, Noumi M and Yamada Y 2002 q -Painlevé systems arising from q -KP hierarchy *Lett. Math. Phys.* **62** 259–68
- [25] Nijhoff F W 1996 On some ‘Schwarzian’ equations and their discrete analogues *Algebraic Aspects of Integrable Systems: In Memory of Irene Dorfman* ed A S Fokas and I M Gel’fand (Basle: Birkhäuser) pp 237–60
- [26] Nijhoff F W 1999 Discrete Painlevé equations and symmetry reduction on the lattice *Discrete Integrable Geometry and Physics* ed A I Bobenko and R Seiler (Oxford: Oxford University Press) pp 209–43

- [27] Nijhoff F W and Capel H W 1995 The discrete Korteweg–de Vries equation *Acta Appl. Math.* **39** 133–58
- [28] Nijhoff F W and Papageorgiou V G 1991 Similarity reductions of integrable lattices and discrete analogues of the Painlevé II equation *Phys. Lett. A* **153** 337–44
- [29] Nijhoff F W, Quispel G R W and Capel H W 1983 Direct linearization of nonlinear difference-difference equations *Phys. Lett. A* **97** 125–8
- [30] Nijhoff F W, Ramani A, Grammaticos B and Ohta Y 2001 On discrete Painlevé equations associated with the lattice KdV systems and the Painlevé VI equation *Stud. Appl. Math.* **106** 261–314
- [31] Nijhoff F W and Walker A J 2001 The discrete and continuous Painlevé VI hierarchy and the Garnier systems *Glasgow Math. J.* **43A** 109–23
- [32] Okamoto K 1987 Studies on the Painlevé equations: 1. 6th Painlevé equation PVI *Ann. Math.* **146** 337–81
- [33] Papageorgiou V G, Nijhoff F W, Grammaticos B and Ramani A 1992 Isomonodromic deformation problems for discrete analogues of Painlevé equations *Phys. Lett. A* **164** 57–64
- [34] Ramani A and Grammaticos B 2005 On the discrete form of the Fokas–Ablowitz equation *Chaos Solitons Fractals* **24** 1331–5
- [35] Ramani A, Grammaticos B and Hietarinta J 1991 Discrete versions of the Painlevé equations *Phys. Rev. Lett.* **67** 1829–32
- [36] Sahadevan R and Capel H W 2003 Complete integrability and confinement of nonautonomous modified Korteweg–de Vries and sine Gordon mappings *Physica A* **330** 373–90
- [37] Sakai H 2001 Rational surfaces associated with affine root systems and geometry of the Painlevé equations *Commun. Math. Phys.* **220** 165–229
- [38] Sakai H 2005 A q -analogue of the Garnier system *Funkcial. Ekvac.* **48** 273–97
- [39] Sakai H 2006 Lax form of the q -Painlevé equation associated with the $A_2^{(1)}$ surface *J. Phys. A: Math. Gen.* **39** 12203–10