

Lack of monotonicity in spin glass correlation functions

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys. A: Math. Theor. 41 385001

(<http://iopscience.iop.org/1751-8121/41/38/385001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.150

The article was downloaded on 03/06/2010 at 07:11

Please note that [terms and conditions apply](#).

Lack of monotonicity in spin glass correlation functions

P Contucci¹, F Unguendoli² and C Vernia²

¹ Dipartimento di Matematica, Università di Bologna, Piazza di Porta S Donato 5,
40127 Bologna, Italy

² Dipartimento di Matematica, Università di Modena e Reggio Emilia, via Campi 213/B,
41100 Modena, Italy

E-mail: contucci@dm.unibo.it, unguendoli@unimore.it and cecilia.vernia@unimore.it

Received 20 June 2008, in final form 22 June 2008

Published 22 August 2008

Online at stacks.iop.org/JPhysA/41/385001

Abstract

We study the response of a spin glass system with respect to the rescaling of its interaction random variables and investigate numerically the behaviour of the correlation functions with respect to the volume. While for a ferromagnet the local energy correlation functions increase monotonically with the scale and, by consequence, with respect to the volume of the system we find that in a general spin glass model those monotonicities are violated.

PACS numbers: 05.50.+q, 75.50.Lk

(Some figures in this article are in colour only in the electronic version)

When spins interact in a ferromagnetic system they tend to be aligned. That simple fact is reflected in structural properties of the statistical mechanics equilibrium state called Griffiths inequalities [1, 2] or more generally GKS [3] inequalities.

The first inequality states that the pressure of a spin system (minus the free energy times beta) does increase with the strength of each interaction among spins. The second says that the correlation among any set of spins increases with respect to the strength of the interaction of any other set of them.

Since the strength of the interaction allows us to switch on and off new parts of the system, the mentioned monotonicity properties can be easily turned into new ones like monotonicity with respect to the volume or with respect to the system dimensionality. All those properties are at the origins of the fruitful applications of those inequalities to prove rigorous results in statistical mechanics, like the existence of the thermodynamic limit for pressure and correlations, bounds for critical temperatures and exponents [4] and their mutual relations for different systems.

While the physical meaning of the Griffiths inequalities was clear and well understood much earlier than rigorously proved, for disordered systems like spin glasses there are no

a priori evident monotonicity properties due to the lack of ferromagnetism and the presence of competition (frustrated loops [5]).

The first case to be understood has been the Gaussian interaction. The use of the partial integration led to the first general proof of existence and monotonicity of thermodynamic limit for the pressure of a d -dimensional spin glass model with general potential [6, 7]. Defining the potential

$$U_\Lambda(J, \sigma) = \sum_{X \subset \Lambda} J_X \sigma_X, \tag{1}$$

where the coefficients J_X are a Gaussian family distributed as

$$\text{Av}(J_X) = 0, \quad \text{Av}(J_X J_Y) = \Delta_X^2 \delta_{X,Y} \tag{2}$$

with its associated random Gibbs–Boltzmann state ω and the quenched measure as

$$\langle - \rangle = \text{Av}[\omega(-)] \tag{3}$$

a straightforward computation [6, 7] gives

$$\langle J_X \sigma_X \rangle = \Delta_X^2 \text{Av}(1 - \omega_X^2), \tag{4}$$

which turns out to be positive by inspection. Although it was clear that the former positivity comes from convexity, it was recognized only few years later that the same result holds in full generality for random interaction J_X with zero average and not only for centred Gaussian variables: in [8] it is proved that (4) can be simply derived from thermodynamic convexity.

A natural perspective to look at the former inequalities is to consider the deformation of the general centred random variable J_X as $\lambda_X J_X$ with $\lambda_X > 0$. The quenched pressure P as a function of the set of lambda's has first and second derivatives

$$\frac{\partial P}{\partial \lambda_X} = \langle J_X \sigma_X \rangle \geq 0 \tag{5}$$

$$\frac{\partial^2 P}{\partial \lambda_X \partial \lambda_Y} = \frac{\partial \langle J_X \sigma_X \rangle}{\partial \lambda_Y} = \text{Av}[J_X J_Y (\omega_{XY} - \omega_X \omega_Y)]. \tag{6}$$

The two quantities (5) and (6) have been extensively studied in $d = 1$ with nearest-neighbour interaction and periodic (or free) boundary conditions in [9]: the sign of (5) also remains positive by shifting on positive values the J averages; the value of (6) turns out to be non-positive for zero mean and positive variance interactions and changes its sign when crossing a line in the mean-variance plane toward the ferromagnetic regime of zero variance and positive mean.

This paper deals with the study of the sign of (6) for higher dimensions or different topologies for the case with zero average interaction. Our findings can be summarized as follows: (6) does not have a definite sign. An explicit counterexample is found, for instance, in the case of a nearest-neighbour spin chain with an extra interaction connecting two distant spins. In principle, a specific topology could not affect the monotonicity in the volume. For this reason we test numerically the nearest-neighbour correlation function for two- and three-dimensional systems of increasing size and find an oscillating behaviour.

Let us consider a closed chain of six spins with nearest-neighbour interaction with one added interaction between the spins two and five. The dependence of the partition function on the couplings $J_{1,2}$ and $J_{2,3}$ is:

$$\begin{aligned} Z = \sum_{\sigma} \exp \left(\beta \sum J_{i,j} \sigma_i \sigma_j \right) &= a \cosh(\beta J_{1,2}) \cosh(\beta J_{2,3}) \\ &+ b \sinh(\beta J_{1,2}) \cosh(\beta J_{2,3}) + c \cosh(\beta J_{1,2}) \sinh(\beta J_{2,3}) \\ &+ d \sinh(\beta J_{1,2}) \sinh(\beta J_{2,3}), \end{aligned}$$

where the four coefficients are

$$\begin{aligned}
 a &= \left(\prod_{(i,j) \neq (1,2), (2,3)} \cosh(\beta J_{i,j}) \right) \cdot \sum_{\partial B = \emptyset} \left(\prod_{(i,j) \in B} \tanh(\beta J_{i,j}) \right), \\
 b &= \left(\prod_{(i,j) \neq (1,2), (2,3)} \cosh(\beta J_{i,j}) \right) \cdot \sum_{\partial B = (1,2)} \left(\prod_{(i,j) \in B} \tanh(\beta J_{i,j}) \right), \\
 c &= \left(\prod_{(i,j) \neq (1,2), (2,3)} \cosh(\beta J_{i,j}) \right) \cdot \sum_{\partial B = (2,3)} \left(\prod_{(i,j) \in B} \tanh(\beta J_{i,j}) \right) \\
 d &= \left(\prod_{(i,j) \neq (1,2), (2,3)} \cosh(\beta J_{i,j}) \right) \cdot \sum_{\partial B = (1,2) \cup (2,3)} \left(\prod_{(i,j) \in B} \tanh(\beta J_{i,j}) \right).
 \end{aligned}$$

The truncated correlation function is now

$$\omega_{12,23} - \omega_{12}\omega_{23} = 16 \frac{ad - bc}{Z^2}.$$

By gauge invariance the Bernoulli random model can be reduced to one in which the randomness is concentrated on the two couplings $J_{1,2} = \pm 1, J_{2,3} = \pm 1$ with probability $1/2$, and the remaining others $J_{i,j} = 1$.

Let us introduce the notation: $C := \cosh(\beta), S := \sinh(\beta), T := \tanh(\beta)$ and $T_{12} := \tanh(\beta J_{1,2}), T_{23} := \tanh(\beta J_{2,3})$ (and similarly for sinh and cosh).

We will indicate with $Z(+, +), Z(+, -), Z(-, +)$ and $Z(-, -)$ the partition functions computed with fixed values of $J_{1,2}$ and $J_{2,3}$.

An explicit computation gives

$$a = C^5; \quad b = c = C^5 T^3; \quad d = C^5 T^4 \quad \Rightarrow \quad ad - bc = C^{10} T^4 (1 - T^2)$$

and

$$\begin{aligned}
 Z &= C^5 C_1 C_2 [1 + T_{12} T^3 + T_{23} T^3 + T_{12} T_{23} T^4] \\
 \Rightarrow Z(++) &= C^7 [1 + 2T^4 + T^6] \\
 Z(+-) = Z(-+) &= C^7 [1 - T^6] \\
 Z(--) &= C^7 [1 - 2T^4 + T^6].
 \end{aligned}$$

Finally

$$\begin{aligned}
 \text{Av}[J_{1,2} J_{2,3} (\omega_{12,23} - \omega_{12}\omega_{23})] &= 16 C^{10} T^4 (1 - T^2) \text{Av} \left[\frac{J_{1,2} J_{2,3}}{Z^2} \right] \\
 &= 16 C^{10} T^4 (1 - T^2) \left\{ \frac{1}{Z^2(++)} + \frac{1}{Z^2(--)} - \frac{2}{Z^2(+-)} \right\} \\
 &= \frac{16 T^4 (1 - T^2)}{C^4} \left\{ \frac{1}{(1 + 2T^4 + T^6)^2} + \frac{1}{(1 - 2T^4 + T^6)^2} - \frac{2}{(1 - T^6)^2} \right\} \\
 &= \frac{16 T^4 (1 - T^2)}{C^4} \frac{8 T^6 [-1 + 3T^2 - 2T^6 + 2T^8 - 4T^{10} - T^{12} + 3T^{14}]}{(1 + 2T^4 + T^6)^2 \cdot (1 - 2T^4 + T^6)^2 \cdot (1 - T^6)^2}.
 \end{aligned}$$

In order to compute the sign, we notice that the square parenthesis term is

$$\begin{aligned}
 &[-1 + 3T^2 - 2T^6 + 2T^8 - 4T^{10} - T^{12} + 3T^{14}] \\
 &= \left[\frac{C^2 (S^2 - 1) (C^8 - S^8) + 2S^4 (C^8 - S^8) + S^8 (C^2 + S^2)}{C^{14}} \right].
 \end{aligned}$$




Figure 1. $Av[J_b \omega_b]$ as a function of L for the two-dimensional lattice with Gaussian and Bernoulli coupling, for two different temperature values $t = 0.5$ and $t = 1.5$ each.

The only term with possible sign change is $(S^2 - 1)$. For small β the leading term is then $-\cosh^{10}(\beta)$, which gives a negative contribution, while for large β everything is positive.

A plot of the function $Av[J_{1,2} J_{2,3} (\omega_{12,23} - \omega_{12} \omega_{23})]$ shows a change of sign around $\beta = 0.695$.

The numerical test is performed for d -dimensional cubic lattices Λ of volume $N = L^d$, with $d = 2, 3$. We analyse two cases of quenched disorder: the Bernoulli couplings with $J_{i,j} = \pm 1$ and the Gaussian couplings with zero mean and unit variance.

Given a spin configuration σ for a system of linear size L , we consider the observable

$$Av[J_b \omega_b], \tag{7}$$

where $b = (i, j)$ with $i, j \in \Lambda, |i - j| = 1$, ω_b is the thermal average of the quantity $\sigma_i \sigma_j$ and $Av[\cdot]$ is the average over the quenched disorder.

With a parallel-tempering algorithm [10] we investigate the correlation (7) for lattice sizes ranging from $L = 3$ to $L = 24$ in the case $d = 2$ and from $L = 3$ to $L = 10$ in the case $d = 3$. For each size we consider at least 2048 disorder realizations and, in order to thermalize the large sizes, we choose up to 37 temperature values in the range $0.5 \leq t \leq 2.3$, in which the critical temperatures of the three-dimensional models ($t_c \simeq 0.95$ for the Gaussian model [11] and $t_c \simeq 1.15$ for the Bernoulli model [12]) are contained. The thermalization in the parallel tempering procedure is tested by checking the symmetry of the probability distribution for the standard overlap q under the transformation $q \rightarrow -q$. Moreover, for the Gaussian coupling case another thermalization test is available: the internal energy can be calculated both as the temporal mean of the Hamiltonian or, using integration by parts, as the expectation of a simple function of the link overlap [13]. We checked that with our thermalization steps both




Figure 2. $Av[J_b \omega_b]$ as a function of L for the three-dimensional lattice with Gaussian and Bernoulli coupling and for two different temperature values $t = 0.5$, and $t = 1.5$ each.

Table 1. Parameters of the simulations: linear system size, number of sweeps used for thermalization, number of sweeps for measurement of the observable, number of disorder realizations, number of temperature values allowed in the parallel tempering procedure, temperature increment, minimum and maximum temperature values.

| L | Therm | Equil | Nreal | n_r | δt | t_{\min} | t_{\max} |
|--|--------|--------|-------|-------|------------|------------|------------|
| Two-dimensional lattice $L^d, d = 2$ | | | | | | | |
| 3–12 | 50 000 | 50 000 | 4096 | 19 | 0.1 | 0.5 | 2.3 |
| 16 | 50 000 | 50 000 | 2048 | 19 | 0.05 | 0.5 | 2.3 |
| 24 | 50 000 | 50 000 | 2600 | 19 | 0.05 | 0.5 | 2.3 |
| Three-dimensional lattice $L^d, d = 3$ | | | | | | | |
| 3–6 | 50 000 | 50 000 | 2048 | 19 | 0.1 | 0.5 | 2.3 |
| 8 | 50 000 | 50 000 | 2680 | 19 | 0.1 | 0.5 | 2.3 |
| 10 | 70 000 | 70 000 | 2048 | 37 | 0.05 | 0.5 | 2.3 |

measurements converge to the same value. All the parameters used in the simulations are reported in table 1.

The numerical results are displayed in figures 1 and 2, where the correlation (7) is represented as a function of the linear system size L for different temperatures both for the two- and three-dimensional system. We find that the correlation oscillates with respect to L , independently of the couplings (Bernoulli and Gaussian), of the temperatures and the dimension d ($d = 2, 3$).

The results presented here show that the monotonicity properties typical of ferromagnetic systems are clearly violated for spin glass models. Further effort is necessary to establish if the quantity (6) can keep a definite sign for specific lattice geometries or may depend on the relative position of the two sets X and Y .

Acknowledgments

PC thanks Hal Tasaki for suggesting the topology used in analytical example. Cristian Giardiná, Sandro Graffi, Frank Den Hollander and Hidetoshi Nishimori are acknowledged for many useful discussions. FU and CV acknowledge GNFM-INdAM for financial support.

References

- [1] Griffiths R B 1967 Correlation in ising ferromagnets *J. Math. Phys.* **8** 478–83
- [2] Griffiths R B 1964 A proof that the free energy of a spin system is extensive *J. Math. Phys.* **5** 1215–22
- [3] Kelly D G and Sherman S 1968 General Griffiths' inequalities on correlations in ising ferromagnets *J. Math. Phys.* **9** 466–84
- [4] Simon B 1993 *The Statistical Mechanics of Lattice Gases* (Princeton, NJ: Princeton University Press)
- [5] Mezard M, Parisi G and Virasoro M A 1987 *Spin Glass Theory and Beyond* (Singapore: World Scientific)
- [6] Contucci P and Graffi S 2004 Monotonicity and thermodynamic limit *J. Stat. Phys.* **115** 581–9
- [7] Contucci P and Graffi S 2004 On the surface pressure for the Edwards–Anderson Model *Comm. Math. Phys. Stat. Phys.* **248** 207–20
- [8] Contucci P and Lebowitz J 2007 Correlation inequalities for spin glasses *Annales Henri Poincaré* **8** 1461–7
- [9] Contucci P and Unguendoli F 2008 Correlation inequalities for spin glass in one dimension *Rend. Lincei Mat. Appl.*, at press
- [10] Hukushima K and Nemoto K 1996 Exchange Monte Carlo method and application to spin glass simulations *J. Phys. Soc. Jpn.* **65** 1604–8
- [11] Marinari E, Parisi G and Ruiz-Lorenzo J J 1998 Phase structure of the three-dimensional Edwards-Anderson spin glass *Phys. Rev. B* **58** 14852–63
- [12] Ballesteros H G, Cruz A, Fernandez L A, Martin-Mayor V, Pech J, Ruiz-Lorenzo J J, Tarancon A, Tellez P, Ullod C L and Ungil C 2000 Critical behavior of the three-dimensional ising spin glass *Phys. Rev. B* **62** 14237–45
- [13] Contucci P 2003 Replica equivalence in the Edwards-Anderson model *J. Phys. A: Math. Gen.* **36** 10961–6