

A boundary field induced first-order transition in the 2D Ising model: numerical study

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys. A: Math. Theor. 41 395001

(<http://iopscience.iop.org/1751-8121/41/39/395001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.150

The article was downloaded on 03/06/2010 at 07:12

Please note that [terms and conditions apply](#).

A boundary field induced first-order transition in the 2D Ising model: numerical study

Elmar Bittner and Wolfhard Janke

Institut für Theoretische Physik and Centre for Theoretical Sciences (NTZ), Universität Leipzig,
Postfach 100 920, D-04009 Leipzig, Germany

E-mail: elmar.bittner@itp.uni-leipzig.de and wolfhard.janke@itp.uni-leipzig.de

Received 4 June 2008, in final form 2 August 2008

Published 4 September 2008

Online at stacks.iop.org/JPhysA/41/395001

Abstract

In a recent paper, Clusel and Fortin (2006 *J. Phys. A: Math. Gen.* **39** 995) presented an analytical study of a first-order transition induced by an inhomogeneous boundary magnetic field in the two-dimensional Ising model. They identified the transition that separates the regime where the interface is localized near the boundary from that where it propagates inside the bulk. Inspired by these results, we measured the interface tension by using multimagnetic simulations combined with parallel tempering to determine the phase transition and the location of the interface. Our results are in very good agreement with the theoretical predictions. Furthermore, we studied the spin–spin correlation function for which no analytical results are available.

PACS numbers: 05.70.Fh, 02.70.Uu, 75.10.Hk

(Some figures in this article are in colour only in the electronic version)

1. Introduction

Wetting transitions are phase transitions in the surface layer of bulk systems which are induced by symmetry-breaking surface fields [1, 2]. The Ising model with a boundary magnetic field is a simple model for such a wetting problem, because Ising ferromagnets have the same critical behaviour as the analogous case of gas–fluid transitions, as has been pointed out by Nakanishi and Fisher [3]. The use of the Ising model with short-range interactions for wetting studies has not only the advantage that one can use all the advanced simulation techniques which have been developed in the past few years. Especially in two dimensions (2D), there are also a lot of theoretical results available for comparison.

The Ising model with a uniform boundary magnetic field on one side of a square lattice has been completely solved by McCoy and Wu [4], whereas the Ising model with a uniform bulk field can only be solved at the critical temperature [5]. For situations with fixed boundary

spins or equivalently infinite boundary magnetic fields [6], or finite boundary magnetic fields [7] some exact results have also been found. In a recent paper, Clusel and Fortin [8] presented an alternative method to that developed by McCoy and Wu for obtaining some exact results for the 2D Ising model with a general boundary magnetic field and for finite-size systems. Their method is based on the fermion representation of the Ising model using a Grassmann algebra. They applied this method for studying the first-order transition induced by an inhomogeneous boundary magnetic field in the 2D Ising model [9]. To be more precise, the boundary magnetic field acts on the $x = 1$ column of spins, being positive in the lower and negative in the upper half. By taking the thermodynamic limit exactly for a given geometry of the lattice, they obtained a simple equation for the transition line and also a threshold for the aspect ratio $\zeta = L_x/L_y = 1/4$, where this line moves into the complex plane. This vanishing of the transition line indicates the crossover from 1D behaviour for $L_x \ll L_y$ to 2D behaviour at large ζ , which is reflected in the behaviour of the boundary spin–spin correlation function.

The aim of this work is to check some of the predictions by carrying out Monte Carlo simulations of this model and to extend the results to parameter ranges and for observables where analytic solutions cannot be obtained. The rest of this paper is organized as follows. In section 2 we give the definition of the model and briefly summarize the theoretical predictions. A description of the employed simulation techniques and the results of our Monte Carlo simulations are presented in section 3, and concluding remarks can be found in section 4.

2. Model and theoretical predictions

We consider a 2D Ising model with a non-homogeneous magnetic field h_y located on one boundary of the system. The Hamiltonian is given by

$$\mathcal{H} = -J \sum_{x,y=1}^{L_x,L_y} (\sigma_{xy}\sigma_{x+1y} + \sigma_{xy}\sigma_{xy+1}) - \sum_{y=1}^{L_y} h_y \sigma_{1y}, \quad (1)$$

with free boundaries in the x -direction and periodic boundary conditions in the y -direction. To compare our results with the theoretical predictions of Clusel and Fortin [9], we consider the same profile of the boundary magnetic field acting on the $x = 1$ column of spins: $h_y = H$ for $y = 1, \dots, L_y/2$ and $h_y = -H$ for $y = L_y/2 + 1, \dots, L_y$, with $H \geq 0$.

In the limit of zero temperature, by using simple energetic arguments, Clusel and Fortin [9] showed that for small H all spins are aligned in one direction as in the bulk case for $H = 0$, see also figure 1. With increasing H , however, depending on the aspect ratio $\zeta = L_x/L_y$ two different interfaces can be formed. If $\zeta > \zeta_s = 1/4$ the interface is localized near the boundary, whereas for $\zeta < \zeta_s$ the interface propagates inside the bulk. The critical ratio ζ_s marks the crossover from 1D behaviour for $L_x \ll L_y$ towards 2D behaviour at large ζ . For $\zeta < \zeta_s$ and non-zero temperatures $T > 0$, with the abbreviations $t = \tanh(J/k_B T)$ and $u = \tanh(H/k_B T)$, the equation for the first-order transition line in the (t, u) -plane turns out to be a quadratic equation in u^2 [9]:

$$2t(1 + v(4\zeta))u^4 + (1 + t^2)(1 - 2tv(4\zeta) - t^2)u^2 + 2(v(4\zeta) - 1)t^3 = 0, \\ v(4\zeta) = \cosh \left[4\zeta \ln \left(\frac{1-t}{t(1+t)} \right) \right]. \quad (2)$$

In figure 2 we show the phase diagram for a system with aspect ratio $\zeta = 0.2$ (and $J = k_B = 1$). In the low-temperature regime we can approximate the above expression by comparing the energy of the interface with the energy induced by the magnetic field. This leads to $H \approx 2\sigma(T)\zeta/m_0(T)$, where $\sigma(T)$ and $m_0(T)$ are the known interface tension and




Figure 1. The phase diagram at zero temperature as a function of the aspect ratio $\zeta = L_x/L_y$ and the boundary magnetic field H .




Figure 2. The phase diagram for a system with $\zeta = 0.2$. The thick line shows the first-order transition given by Clusel and Fortin [9] and the thin vertical line indicates the second-order bulk phase transition. The thin dashed line indicates a simple approximation of the first-order line, $H \approx 2 \frac{\sigma(T)\zeta}{m_0(T)}$, where $\sigma(T)$ and $m_0(T)$ are the interface tension and the spontaneous magnetization of the pure 2D Ising model, respectively. The sketches of the spin configurations illustrate the interface location in the three phases and the double-headed arrows show the parameters of the Monte Carlo simulations.

the spontaneous magnetization of the pure 2D Ising model, respectively. This approximation reproduces the exact low- T expansion, $H = 4\zeta - 4\zeta T e^{-2/T}$, and works very well for $T < 1$

Table 1. Summary of simulation parameters (PT: parallel-tempering algorithm, SC: single-cluster update).

ζ	H	T	$L_x \times L_y$	Method	Measurements
0.2	0.4	1.6–2.2	80–2000	PT	1×10^6
0.2	0.5	1.4–1.9	80–2000	PT	1×10^6
0.2	0.5	2.1–2.3	80–180 500	SC	1×10^6 – 5×10^6
0.2	0.7–0.9	1.0	80–640	PT	1×10^6
0.2	0.285–0.316	2.0	80–2000	PT	1×10^6
0.25	0.4	1.9–2.1	256–2500	PT	1×10^6
0.25	0.5	1.5–1.9	64–2000	PT	1×10^6
0.25	0.5	2.0–2.3	64–6400	SC	1×10^6 – 5×10^6
0.25	0.7–0.8	1.5	64–1600	PT	1×10^6
0.5	0.5	2.2–2.35	50–3200	PT	1×10^6
0.5	0.9–1.1	1.0	50–5000	PT	1×10^6
0.5	1.5–2.0	1.0	50–3872	PT	1×10^6

as one can see in figure 2 (thin dashed line). Since $\sigma(T)$ vanishes much faster than $m_0(T)$ as $T \rightarrow T_c$, also this point is reproduced exactly, but the slope of the approximate transition line at T_c does not diverge as for the exact solution. Due to the first-order transition induced by the inhomogeneous boundary magnetic field, the second-order phase transitions across the vertical line at $T = T_c$ are transitions from a region where an interface in the bulk separates two ordered domains of opposite magnetization from a disordered regime above the transition temperature. Therefore, the system undergoes a transition without a change in the magnetization $\langle m \rangle$ which is zero in both phases, cf figure 2, but the width of the magnetization distribution does change.

3. Numerical results

Since we are primarily interested in the location of the interface induced by the boundary field, we first performed simulations at low temperatures to generate a well-defined interface. To overcome the slow dynamics at low temperatures we developed a combination of the multimagnetic algorithm with the parallel tempering method [10] for which we used two different schemes: in the first scheme, we kept the magnetic field value H fixed and simulated $n = 32$ replica of the system at different temperatures T_i . In the second scheme, we kept the temperature T fixed and used $n = 32$ different values of the magnetic field H_i .

To construct the weight function for the multimagnetic part of the algorithm, we employed an accumulative recursion, described in detail in [10, 11]. Statistical averages were taken over runs of 10^6 Monte Carlo (MC) steps, where one MC step consists of one full multimagnetical lattice sweep for all 32 replica and one attempted parallel tempering exchange of all adjacent replica. With this method we were able to study systems with $N = L_x \times L_y = 50$ to 5000 spins for aspect ratios $\zeta = L_x/L_y = 0.2, 0.25 = \zeta_s$ and 0.5, for further details see table 1.

Let us first discuss the data obtained for the case $\zeta = 0.2 < \zeta_s$. For this value of the aspect ratio, the phase diagram as predicted by Clusel and Fortin [9] is shown in figure 2. The thick line indicates the first-order transitions from the fully magnetized state with $\langle m \rangle > 0$ to the mixed state with an interface extending across the bulk. To check the nature of these transitions we measured the probability density of the magnetization at four points along the transition line. In the first two cases, we kept the boundary magnetic field constant ($H = 0.4$ and 0.5) and varied the temperature to locate the transition point, and in the other two cases,




Figure 3. Left plot: the probability density for the magnetization as a function of temperature at a constant boundary magnetic field $H = 0.5$ for $N = 20 \times 100$. The vertical lines indicate the first-order transition temperature $T_0 \approx 1.5950$ according to (2) and the critical temperature $T_c \approx 2.2692$ of the bulk phase transition. Right plot: histograms of the magnetization in the vicinity of T_0 for various lattice sizes ranging from $L_y = 20$ to $L_y = 100$. Here the temperatures $T_0(L)$ are determined such that the peaks at $m \approx \pm m_0$ and $m = 0$ are of equal height.

we fixed the temperature ($T = 1.0$ and 2.0) and varied the boundary magnetic field. These points are indicated by the double-headed arrows in figure 2.

In the following we illustrate our procedure for obtaining the first-order transition point and the associated interface tension for the case of fixed $H = 0.5$. A level plot of the magnetization density $m = (1/N) \sum_{x,y=1}^{L_x,L_y} \sigma_{xy}$ as a function of temperature is shown in figure 3 (left). For each lattice size, a pseudo-transition point can be defined by varying the temperature until the peaks at $m \approx \pm m_0$ and $m = 0$ are of equal height, which can be achieved by histogram reweighting. The interface tension can then be estimated from [12]

$$F_L^s = \frac{1}{2L} \ln \left(\frac{P_L^{\max}}{P_L^{\min}} \right), \quad (3)$$

where P_L^{\max} is the value of the peaks and P_L^{\min} denotes the minimum in between, see figure 3 (right). The length of the interface is denoted by L , which is $L = L_x$ in the case of $\zeta < \zeta_s$.

The thus-defined pseudo-transition temperatures $T_0(L)$ approach the infinite-volume transition temperature T_0 as $1/L^2$, and for the final estimate of $F^s = \lim_{L \rightarrow \infty} F_L^s$, we performed a fit according to

$$F_L^s = F^s + \frac{a}{L} + \frac{b \ln(L)}{L}. \quad (4)$$

At fixed T one proceeds analogously by varying the magnetic field H , i.e., the roles of T and H are just interchanged. For all four cuts at constant surface field or temperature we find good agreement with the infinite-volume transition points derived from (2) and a clearly nonzero interface tension, see table 2.

We also checked the critical behaviour along the line of second-order transitions at $T = T_c = 2/\log(1 + \sqrt{2}) \approx 2.2692$. To this end, we run at $H = 0.5$ single-cluster simulations (suitably adapted to the surface field) for systems with $N = L_x \times L_y = 4 \times 20$ to 190×950 spins and performed a finite-size scaling (FSS) analysis to determine the transition point and some critical exponents. This particular value of the magnetic field has been chosen because of the relatively large temperature gap between the boundary field induced first-order transition and the bulk phase transition. Between each measurement we performed one sweep, which here consists of n single-cluster updates with n chosen such that $n \langle |S| \rangle \approx N$, where $\langle |S| \rangle$ is the average cluster size. For every run we generated $1 \times 10^6 - 5 \times 10^6$ sweeps, and




Figure 4. FSS of the specific-heat maxima C_{\max} . The logarithmic fit $C_{\max} = a + b \log(L_x)$ and an unbiased fit using the power-law ansatz $C_{\max} = a + bL_x^{\alpha/\nu}$ are almost indistinguishable on the scale of the figure.

Table 2. Results of the simulations close to the first-order transition line. For a given aspect ratio ζ we kept either the boundary magnetic field H or the temperature T fixed. The third column shows the measured transition points and the fourth column shows the exact infinite-volume values given by (2). The fifth column contains our numerical estimates for the extrapolated interface tensions.

ζ	H	T_0	T_0 (exact)	F_s
0.2	0.4	1.84 (1)	1.822 52...	0.18 (1)
0.2	0.5	1.60 (1)	1.594 97...	0.32 (2)
0.25	0.5	1.91 (1)	1.958 45...	0.09 (1)
ζ	T	H_0	H_0 (exact)	F_s
0.2	1.0	0.72 (1)	0.702 352...	0.82 (2)
0.2	2.0	0.305 (3)	0.305 928...	0.12 (1)
0.25	1.5	0.73 (1)	0.762 807...	0.24 (1)

recorded the time series of the energy density $e = E/N$ and the magnetization density. Using these time series, we can compute the specific heat, $C = N(\langle e^2 \rangle - \langle e \rangle^2)/T^2$, the (finite lattice) susceptibility, $\chi = N(\langle m^2 \rangle - \langle m \rangle^2)$, and the Binder cumulant $U = 1 - \langle m^4 \rangle/3\langle m^2 \rangle^2$ in the vicinity of the simulation point by reweighting.

In this way we can use the maxima of the (finite lattice) susceptibility to detect the pseudo-critical points and can obtain an estimate for T_c from a linear least-squares fit of their scaling behaviour, $T_{\max} - T_c \propto L_x^{-1/\nu} = L_x^{-1}$, assuming thus the exact value $\nu = 1$ according to the universality class of the 2D Ising model. This leads to an estimate for the critical temperature, $T_c = 2.2695(7)$, which is in very good agreement with the exact value. The FSS ansatz for the (finite lattice) susceptibility maxima χ_{\max} is taken as usual as $\chi_{\max} \propto L_x^{\gamma/\nu}$. From a (linear) least-squares fit, we find that $\gamma/\nu = 1.75(4)$ is in perfect agreement with the exact value $7/4$. Concerning the specific heat we expect in the case of the Onsager exponent $\alpha = 0$ a logarithmic divergence of the form $C_{\max} = a + b \log(L_x)$. Indeed, the data can be fitted nicely with this ansatz, cf figure 4. We also tried an unbiased fit using the power-law ansatz $C_{\max} = a + bL_x^{\alpha/\nu}$, which gives us $\alpha = 0.05(2)$, verifying the expected value.




Figure 5. Spin–spin correlation function $\langle \sigma_{1y} \sigma_{xy} \rangle$ for $\zeta = 0.2$, $L_y = 100$ and $H = 0.5$ measured in different regimes: (a) ordered regime at $T = 1.0$, (b) regime with an interface in the bulk at $T = 2.0$ and (c) disordered regime at $T = 2.5$.

For the aspect ratio $\zeta = 0.2$, we also sampled the spin–spin correlation functions $\langle \sigma_{1y} \sigma_{xy} \rangle$ and $\langle \sigma_{x1} \sigma_{xy} \rangle$ in x - and y -direction, respectively. In the ordered regime where nearly all spins are aligned in the same direction ($\sigma_{xy} = +1$, say) we find for $\langle \sigma_{1y} \sigma_{xy} \rangle$ and $1 \leq y \leq L_y/2$ (where $h_y = H > 0$) almost constant values near unity as one expects. The fast decay of the spin–spin correlation function to a slightly smaller value in the upper half of the system indicates that the interface is localized near the boundary, cf figure 5(a). In the regime with an interface in the bulk along the x -direction we find a symmetric shape of $\langle \sigma_{1y} \sigma_{xy} \rangle$ as a function of y which is a clear signal for the phase separation, see figure 5(b). The opening angle between the plus and minus phases for $1 \leq y \leq L_y/2$ and $L_y/2 + 1 \leq y \leq L_y$, respectively, is a measure for the fluctuations of the interface, e.g. a stiff interface shows an acute angle. For temperatures above the critical temperature, i.e. in the disordered phase, we observe a similar vanishing of the spin–spin correlation functions as in the pure 2D Ising model, see figure 5(c).

At zero temperature, for aspect ratios ζ larger than the critical ratio $\zeta_s = 0.25$ and strong fields $H > J(1 + 4/L_y)$, the interface is localized on the boundary, cf figure 1. Although for $\zeta > \zeta_s$ no real solution of (2) exists near T_c , one can solve the equation for small temperatures¹ and finds in the case $\zeta = 0.5$ the phase diagram shown in figure 6. The zero-temperature limit is consistent with the energetic arguments in [9], see also figure 1. The lower line starting at $T = 0$, $H = 1$ can also be detected by means of computer simulations, but as one can see in the right plot of figure 6, for $T = 1$ the dip between the two maxima in the boundary magnetization density $m_b = (1/L_y) \sum_{y=1}^{L_y} \sigma_{1,y}$ vanishes with increasing lattice sizes. Therefore, there is no signal for a first-order transition between these two regimes. One can argue that there is no phase transition at all, because in the infinite-volume limit only the ordered phase survives. The dashed line in the left plot of figure 6 starting at $T = 0$, $H = 4\zeta = 2$ is not visible in simulations, because this second solution of (2) would correspond to the boundary between the bulk magnetized state and configurations with an interface propagating inside the bulk which, however, have a higher free energy than configurations with an interface localized near the

¹ Here we are in disagreement with [9] where no solution was found, because of a mistake in the discriminant of (2).




Figure 6. Left plot: the finite-size phase diagram for a system with $\zeta = 0.5$. The thick line shows the transition line between the ordered phase and the region with an interface localized on the boundary. The dashed line indicates where a configuration with no interface and a configuration with an interface propagating inside the bulk have the same free energy (non-physical solution of (2), because this bulk interface configuration has a higher free energy than that with an interface localized on the boundary and hence is suppressed.). The thin vertical line indicates the second-order bulk phase transition. Right plot: probability density of the boundary magnetization m_b for $T = 1.0$ plotted for various lattice sizes ranging from $L_y = 16$ to $L_y = 88$ and boundary magnetic field values $H \approx 1$ where the peaks are of equal height.

boundary and hence are suppressed. Therefore the phase diagram for $\zeta > 0.25$ consists only of two phases, namely the ordered low-temperature phase and the disordered high-temperature phase as in the pure 2D Ising model. We hence conclude that for $\zeta > 0.25$ the inhomogeneous boundary magnetic field only leads to finite-size effects.

Finally, let us come to the special case of $\zeta = \zeta_s = 0.25$. While the transition line disappears for $\zeta > 1/4$ as the solutions of (2) move to the complex plane, for ζ_s we still do see different peaks associated with the two phases and, therefore, a finite interface tension, cf table 2. The numerically estimated transition points for $H = 0.5$ and $T = 1.5$ also contained in table 2 are again seen to be in good agreement with (2). Furthermore, we analysed for $T = 1.0$ the spin–spin correlation functions $\langle \sigma_{1y} \sigma_{xy} \rangle$ and $\langle \sigma_{x1} \sigma_{xy} \rangle$ in x - and y -direction, respectively. Slightly below the transition from the ordered phase with $\langle m \rangle = m_0$ to the phase with an interface propagating inside the bulk and therefore $\langle m \rangle = 0$, $\langle \sigma_{1y} \sigma_{xy} \rangle$ shows an asymmetric shape and a fast decay near the boundary, indicating that the interface is localized near the boundary, cf figure 7(a). With increasing boundary magnetic field we cross the first-order line at $H_0 \approx 0.925$, where the interface starts moving into the bulk and, therefore, the profile of $\langle \sigma_{1y} \sigma_{xy} \rangle$ becomes symmetric in y . Right at the transition line where we have two coexisting phases we find both the fast decay near the boundary and the opening angle between the two almost symmetric halves, see figure 7(b). When the boundary magnetic field is increased further, this mixed-phase effect vanishes and the interface in the bulk becomes stable. In this case we find a symmetric shape of the spin–spin correlation function, cf figure 7(c), similar to $\zeta = 0.2$ in figure 5(b).

4. Summary

Our Monte Carlo data clearly confirm the theoretical considerations of Clusel and Fortin [9] and extend their exact results by studying the cases ζ equal and larger than the critical




Figure 7. Spin–spin correlation function $\langle \sigma_{1y} \sigma_{xy} \rangle$ for $\zeta = 0.25$, $L_y = 80$ and $T = 1$ measured for different values of the boundary magnetic field: (a) $H = 0.9$, (b) $H = 0.925$ and (c) $H = 0.975$.

value $\zeta_s = 1/4$. The observed finite-size scaling behaviour fits nicely with their predictions for the infinite system, cf our results in table 2. We also find that for a large aspect ratio some interesting finite-size effects can be observed, such as, for example, a regime in the H – T plane with two states separated by an energy gap which vanishes in the infinite-volume limit. Furthermore, we studied the spin–spin correlation function for which analytical results are not yet available. Since this observable turned out to be quite sensitive to the interface location, it would be a challenging enterprise to pursue further analytical considerations in this direction.

Acknowledgments

We gratefully acknowledge financial support from the Deutsche Forschungsgemeinschaft (DFG) under grant no. JA 483/23-1, the DFG Research Group FOR877, and the EU RTN-Network ‘ENRAGE’: Random Geometry and Random Matrices: From Quantum Gravity to Econophysics under grant no. MRTN-CT-2004-005616.

References

- [1] Cahn J W 1977 *J. Chem. Phys.* **66** 3667
- [2] Ebner C and Saam W F 1977 *Phys. Rev. Lett.* **38** 1486
- [3] Nakanishi H and Fisher M E 1983 *J. Chem. Phys.* **78** 3279
- [4] McCoy B M and Wu T T 1973 *The Two-Dimensional Ising Model* (Cambridge, MA: Harvard University Press)
- [5] Zamolodchikov A B 1989 *Adv. Stud. Pure Math.* **19** 641
Zamolodchikov A B 1989 *Int. J. Mod. Phys. A* **4** 4235
- [6] Abraham D B 1980 *Phys. Rev. Lett.* **44** 1165
Abraham D B 1982 *Phys. Rev. B* **25** 4922
Abraham D B 1988 *Phys. Rev. B* **37** 3835
- [7] Au-Yang H and Fisher M E 1975 *Phys. Rev. B* **11** 3469
- [8] Clusel M and Fortin J-Y 2005 *J. Phys. A: Math. Gen.* **38** 2849

- [9] Clusel M and Fortin J-Y 2006 *J. Phys. A: Math. Gen.* **39** 995
- [10] Janke W 2003 Histograms and all that *Computer Simulations of Surfaces and Interfaces (NATO Science Series, II. Mathematics, Physics and Chemistry vol 114)* ed B Dünweg, D P Landau and A I Milchev (Dordrecht: Kluwer) pp 137–57
- [11] Berg B A 1996 *J. Stat. Phys.* **82** 323
- [12] Janke W 2003 First-order phase transitions *Computer Simulations of Surfaces and Interfaces (NATO Science Series, II. Mathematics, Physics and Chemistry vol 114)* ed B Dünweg, D P Landau and A I Milchev (Dordrecht: Kluwer) pp 111–35