

## Heisenberg–Weyl algebra revisited: combinatorics of words and paths

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys. A: Math. Theor. 41 415204

(<http://iopscience.iop.org/1751-8121/41/41/415204>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.152

The article was downloaded on 03/06/2010 at 07:16

Please note that [terms and conditions apply](#).

# Heisenberg–Weyl algebra revisited: combinatorics of words and paths

**P Blasiak<sup>1</sup>, G H E Duchamp<sup>2</sup>, A Horzela<sup>1</sup>, K A Penson<sup>3</sup>  
and A I Solomon<sup>3,4</sup>**

<sup>1</sup> H. Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences,  
ul. Radzikowskiego 152, PL 31342 Kraków, Poland

<sup>2</sup> LIPN, UMR 7030, Université Paris 13 – CNRS, 99 av. J-B Clément, F-93430 Villetaneuse,  
France

<sup>3</sup> Laboratoire de Physique Théorique de la Matière Condensée, Université Pierre et Marie Curie,  
CNRS UMR 7600, Tour 24-2ième ét., 4 pl. Jussieu, F 75252 Paris Cedex 05, France

<sup>4</sup> Physics and Astronomy Department, The Open University, Milton Keynes MK7 6AA, UK

E-mail: [pawel.blasiak@ifj.edu.pl](mailto:pawel.blasiak@ifj.edu.pl), [ghed@lipn-univ.paris13.fr](mailto:ghed@lipn-univ.paris13.fr), [andrzej.horzela@ifj.edu.pl](mailto:andrzej.horzela@ifj.edu.pl),  
[penon@lptl.jussieu.fr](mailto:penon@lptl.jussieu.fr) and [a.i.solomon@open.ac.uk](mailto:a.i.solomon@open.ac.uk)

Received 24 June 2008

Published 18 September 2008

Online at [stacks.iop.org/JPhysA/41/415204](http://stacks.iop.org/JPhysA/41/415204)

## Abstract

The Heisenberg–Weyl algebra, which underlies virtually all physical representations of quantum theory, is considered from the combinatorial point of view. We provide a concrete model of the algebra in terms of paths on a lattice with some decomposition rules. We also discuss the rook problem on the associated Ferrers board; this is related to the calculus in the normally ordered basis. From this starting point we explore combinatorial underpinning of the Heisenberg–Weyl algebra, which offers novel perspectives, methods and applications.

PACS numbers: 03.65.Fd, 02.10.Ox

## 1. Introduction

From a modern viewpoint, the formalism and structure of quantum theory are founded on the theory of Hilbert space [1, 2]. The physical content of the theory consists of representing physical quantities as operators which satisfy some algebraic relations. Virtually all correspondence schemes come endowed with the Heisenberg–Weyl algebra structure, be it the canonical quantization scheme, the occupation number representation in quantum mechanics or the second quantization formalism of quantum field theory. This derives from the analogy with classical mechanics whose Poissonian structure is reflected in the commutator of position and momentum observables [3]. Ubiquitous and profound, the Heisenberg–Weyl algebra has become the hallmark of non-commutativity in quantum theory.

An exemplary model of the Heisenberg–Weyl algebra involves combinations of derivative  $D$ , multiplication  $X$  and identity  $I$  operators acting on the space of polynomials. Physical examples are the position  $\hat{x}$  and momentum  $\hat{p}$  operators in the space of square integrable functions, or the annihilation  $a$  and creation  $a^\dagger$  operators in Fock space. Here, without loss of generality, we conform to the notation  $\{a, a^\dagger\}$  for the generators of the (associative) algebra<sup>5</sup>, satisfying

$$aa^\dagger = a^\dagger a + I, \tag{1}$$

where  $I$  is the multiplicative identity. We shall be interested in combinatorial aspects of this relation and discuss one of the ensuing models of the Heisenberg–Weyl algebra.

Indeed, the combinatorial properties of equation (1) were recognized early and successfully applied to the domain of algebraic enumeration, principally concerning the action of the operators  $X$  and  $D$  on generating series. From this point of view, these operators are auxiliary constructions facilitating enumeration of discrete structures [4–10].

However, in this paper we adopt another approach, which leads towards a combinatorial model of the algebra itself. Starting from the definition of the Heisenberg–Weyl algebra as the algebra of words in  $a$  and  $a^\dagger$  supplemented by the relation of equation (1), we will recast it in the language of paths on a lattice with some decomposition rules. We shall also consider a convenient choice of basis, here taken to be normally ordered monomials, which permits a direct link to the combinatorics of words and the related algebra of paths. In this way, algebraic problems may be expressed in the more concrete form of the decomposition and enumeration of paths. For illustration, we consider the normal ordering procedure which reduces to the familiar rook problem on the Ferrers board and then derive the structure constants of the algebra by a simple combinatorial argument.

## 2. Heisenberg–Weyl algebra

In this paper, we consider an *algebra*  $\mathcal{A}$  to be a linear vector space over a field  $\mathbb{K}$  with a bilinear multiplication law:

$$\mathcal{A} \times \mathcal{A} \ni (b, c) \longrightarrow bc \in \mathcal{A}, \tag{2}$$

which is associative and possesses a unit element  $I$ . More precisely, it is called an *associative algebra with unit* as distinct from an algebraic structure lacking associativity or a unit (e.g. Lie algebra). A *basis* of an algebra is a basis for its vector space structure. Each basis  $(b_i)_{i \in \Lambda}$  defines a unique family  $\gamma_{ij}^k \in \mathbb{K}$  such that for every ordered pair  $(i, j) \in \Lambda \times \Lambda$  the set of  $k \in \Lambda$  such that  $\gamma_{ij}^k \neq 0$  is finite and

$$b_i b_j = \sum_{k \in \Lambda} \gamma_{ij}^k b_k. \tag{3}$$

$\gamma_{ij}^k$  are called the *structure constants* of the algebra  $\mathcal{A}$  with respect to the basis  $(b_i)_{i \in \Lambda}$ , from which the multiplication law can be uniquely recovered.

The *Heisenberg–Weyl algebra*, denoted by  $\mathcal{H}$ , is the algebra generated by  $a, a^\dagger$ , satisfying the relation of equation (1). Elements of the algebra  $A \in \mathcal{H}$  are linear combinations of finite products of  $a$  and  $a^\dagger$  of the form

$$A = \sum_{\mathbf{r}, \mathbf{s}} \alpha_{\mathbf{r}, \mathbf{s}} a^{\dagger r_1} a^{s_1} a^{\dagger r_2} a^{s_2} \dots a^{\dagger r_k} a^{s_k}, \tag{4}$$

<sup>5</sup> We do not attach much weight to this particular realization, however, as we shall study algebraic properties only, for which the underlying Fock space plays no role. Our considerations hold true for any representation of the Heisenberg–Weyl algebra.

where  $\mathbf{r} = (r_1, \dots, r_k)$  and  $\mathbf{s} = (s_1, \dots, s_k)$  are non-negative integer multi-indices (with the convention  $a^0 = a^{\dagger 0} = I$ ). This representation is ambiguous, however, due to the commutation relation (1) which yields different representations of the same element of the algebra, e.g.  $aa^\dagger = a^\dagger a + I$ . The problem can be resolved by fixing the preferred order of the generators  $a$  and  $a^\dagger$ . Conventionally, it is done by choosing the *normally ordered* form in which all annihilators stand to the right of creators. In this case, each element of the algebra  $\mathcal{H}$  is uniquely written in the normally ordered form as

$$A = \sum_{r,s} \beta_{r,s} a^{\dagger r} a^s. \tag{5}$$

Hence, the normally ordered monomials  $a^{\dagger r} a^s$  constitute a natural basis for the Heisenberg–Weyl algebra:

$$\text{Basis of } \mathcal{H} : \quad b_{(r,s)} = a^{\dagger r} a^s, \tag{6}$$

indexed by pairs of integers  $r, s = 0, 1, 2, \dots$ , and equation (5) is the expansion of the element  $A$  in this basis. We should note that the normally ordered representation of the elements of the algebra suggests itself as the simplest one [11]. It is important and commonly used in practical applications in quantum optics [12–14] or quantum field theory [15, 16]. Working on this basis entails the reshuffling of  $a$  and  $a^\dagger$  to the normally ordered form, which in general is a nontrivial task [17]. This brings up the issue of efficient calculation methods and intuitive schemes providing insight into the ordering procedure itself. Below, we provide a combinatorial model for the Heisenberg–Weyl algebra and then propose a resolution of the problem from this starting point.

### 3. Combinatorics of the Heisenberg–Weyl algebra

#### 3.1. Words, paths and rook problems

We start by showing that each word in a two-letter alphabet can be uniquely encoded as a *staircase path* on a plane rectangular lattice. This observation will lead us to considering the associated Ferrers board and the rook problem. Rather than giving a formal construction, we shall illustrate it by an example from which the general scheme can be straightforwardly recovered.

Suppose we consider a word, say

$$w = aa^\dagger aa^\dagger a^\dagger a^\dagger aa^\dagger a \tag{7}$$

to which we assign a staircase path on a rectangular lattice. Starting from the point  $(0, 0)$ , it is constructed by reading the word  $w$  from the left and drawing a line to the right if the letter is  $a^\dagger$  and up if the letter is  $a$ , as shown in figure 1 on the left. We observe that this scheme provides a unique encoding of words.

With each path (or word), one can associate a *Ferrers board*  $B_w$  by retaining rectangular cells below the path, see figure 1. Note that this is a many-to-one procedure as paths differing by a horizontal line at the beginning and a vertical line at the end yield the same board [5, 10].

The *rook problem* for the given board  $B$  consists in enumerating non-capturing arrangements of  $k$  rooks on the board which defines a finite sequence  $r_B(k), k = 0, 1, 2, \dots$ , called the *rook numbers*. In our example, one has

$$r_B(k) = 1, 10, 23, 9, 0, 0, \dots \quad k = 0, 1, 2, 3, \dots \tag{8}$$

A rook sequence can be conventionally encoded in a *rook polynomial* defined by

$$R_B(x) = \sum_{k \geq 0} r_B(k) x^k. \tag{9}$$








Figure 3. Path defined by the word  $w = a^{+r} a^s a^{+k} a^l$  and the associated Ferrers board.

- (i) replacing an *upper-left-hand* corner ( $\ulcorner$ ) by a *lower-right-hand* corner ( $\llcorner$ ), and
- (ii) removing the row and column which intersect in the given cell ( $\cdot\cdot$ ).

Note that this reduction is exactly equivalent to the decomposition of the associated Ferrers board induced by the rook problem. In fact, any path can be uniquely decomposed into a finite sum of paths without steps (paths pertaining to monomials  $a^{+r} a^s$ ). The latter constitute the basis in  $\mathcal{P}/\{\ulcorner=\llcorner\}$  corresponding to the normally ordered basis in  $\mathcal{H}$ .

In this way, we obtain the combinatorial model of the Heisenberg Weyl algebra as the algebra of paths with relation (13), i.e.

$$\mathcal{H} = \mathcal{W}/\{aa^\dagger = a^\dagger a + I\} \cong \mathcal{P}/\{\ulcorner=\llcorner\}. \tag{14}$$

With this equivalence the algebraic structure of  $\mathcal{H}$  and its calculus are recast in combinatorial terms, thus providing different perspectives and making accessible intuitive combinatorial arguments.

### 3.4. Example: structure constants

To illustrate the above equivalence, we show how to calculate the structure constants equation (3) of the Heisenberg–Weyl algebra by simple combinatorial enumeration. In the form suited for our purposes here we multiply two elements of the basis of equation (6), say  $b_{(r,s)}$  and  $b_{(k,l)}$ , defining the structure constants  $\gamma_{(r,s)(k,l)}^{(p,q)}$  in the form

$$b_{(r,s)} b_{(k,l)} = \sum_{p,q} \gamma_{(r,s)(k,l)}^{(p,q)} b_{(p,q)}. \tag{15}$$

Expansion in this basis of the product  $b_{(r,s)} b_{(k,l)}$  essentially comes from the normal ordering of the word

$$w = a^{+r} a^s a^{+k} a^l. \tag{16}$$

Following the scheme of section 3.1, we draw the associated path and subsequently read off the Ferrers board  $\mathcal{B}_w$  which has a simple rectangular form; see figure 3. Enumeration of non-capturing rook arrangements on the board is a direct result of the following observations:

- (1) the maximum number of rooks on the board is  $\min\{s, k\}$ , and
- (2) there are as many possible arrangements of  $i$  rooks as unordered choices of  $i$  columns and rows in the board  $\mathcal{B}_w$ .

Hence, the rook numbers are

$$r_{B_{\mathcal{W}}}(i) = i! \binom{s}{i} \binom{k}{i}, \tag{17}$$

for  $i = 0 \dots \min\{s, k\}$  and zero otherwise. Consequently, equation (11) gives the normally ordered form of the word  $w$ :

$$a^{\dagger r} a^s a^{\dagger k} a^l = \sum_{i=0}^{\min\{k,s\}} i! \binom{s}{i} \binom{k}{i} a^{\dagger r+k-i} a^{s+l-i}. \tag{18}$$

Finally, from equation (15) one may read off the structure constants of the algebra  $\mathcal{H}$  in the normally ordered basis. For fixed  $(r, s)$  and  $(k, l)$ , the only non-vanishing  $\gamma$ 's are

$$\gamma_{(r,s)(k,l)}^{(r+k-i,s+l-i)} = i! \binom{s}{i} \binom{k}{i} \quad \text{for } i = 0, \dots, \min\{k, s\}. \tag{19}$$

Note that the right-hand side of equation (19) is, in fact, independent of  $r$  and  $l$  since the outer elements are not included in the commutation.

#### 4. Summary

We considered the Heisenberg–Weyl algebra  $\mathcal{H}$  starting from the (two generator) free algebra of words  $\mathcal{W}$  and imposing the relation of equation (1), i.e.

$$\mathcal{W} \xrightarrow{[a,a^{\dagger}]=I} \mathcal{H}$$

We showed that words can be uniquely encoded as staircase paths on a plane, thus providing a realization of  $\mathcal{W}$  as a combinatorial algebra of paths  $\mathcal{P}$ . This allowed us to construct a model of  $\mathcal{H}$  in terms of paths with the decomposition rule of equation (13) in  $\mathcal{P}$  which reflects the defining relation of equation (1) in  $\mathcal{W}$ . The following diagram illustrates the whole scheme:

$$\begin{array}{ccc} \mathcal{W} & \xleftrightarrow{\sim} & \mathcal{P} \\ \downarrow & & \downarrow \\ \mathcal{W}/_{aa^{\dagger}=a^{\dagger}a+I} & \xleftrightarrow{\sim} & \mathcal{P}/_{\{\tau=\downarrow+\cdot\}} \end{array}$$

We further looked at the normally ordered basis in  $\mathcal{H}$  and the corresponding basis in  $\mathcal{P}/_{\{\tau=\downarrow+\cdot\}}$ . We pointed out that the decomposition rule of equation (13), reducing the number of steps in a path, is closely related to the familiar rook problem on the associated Ferrers board. This permits a graphical illustration of the calculus in this basis and, more generally, of the normal ordering problem.

In this paper, we have advocated a combinatorial approach to the Heisenberg–Weyl algebra by showing that it can be conceived as having a purely combinatorial origin. This gives a new perspective on the whole algebraic framework which is easily amenable to the sophisticated methods of discrete mathematics. Finally, we should mention other models of the algebra based on various discrete structures such as set partitions, graphs or urn models [22–24] as well as results deriving from combinatorial methodology; see [17] and references therein.

#### Acknowledgments

We thank Hayat Cheballah for useful discussions. One of us (PB) acknowledges his appreciation of the warm hospitality and support of the Laboratoire d'Informatique de l'Université Paris-Nord in Villetaneuse where most of his research was carried out. PB and AH wish to acknowledge support from the Polish Ministry of Science and Higher Education under grants no. N202 061434 and N202 107 32/2832.



## References

- [1] Isham C J 1995 *Lectures on Quantum Theory: Mathematical and Structural Foundations* (London: Imperial College Press)
- [2] Peres A 2002 *Quantum Theory: Concepts and Methods* (New York: Academic)
- [3] Dirac P A M 1982 *The Principles of Quantum Mechanics* 4th edn (New York: Oxford University Press)
- [4] Riordan J 1984 *An Introduction to Combinatorial Analysis* (New York: Wiley)
- [5] Stanley R P 1999 *Enumerative Combinatorics* (Cambridge: Cambridge University Press)
- [6] Comtet L 1974 *Advanced Combinatorics* (Dordrecht: Reidel)
- [7] Wilf H S 1994 *Generating functionology* (New York: Academic)
- [8] Flajolet P and Sedgewick R 2008 *Analytic Combinatorics* (Cambridge: Cambridge University Press)
- [9] Bergeron F, Labelle G and Leroux P 1998 *Combinatorial Species and Tree-Like Structures* (Cambridge: Cambridge University Press)
- [10] Bryant V 1968 *Aspects of Combinatorics: A Wide-ranging Introduction* (Cambridge: Cambridge University Press)
- [11] Cahill K E and Glauber R J 1969 Ordered expansions in boson amplitude operators *Phys. Rev.* **177** 1857–81
- [12] Glauber R J 1963 The quantum theory of optical coherence *Phys. Rev.* **130** 2529–39
- [13] Schleich W P 2001 *Quantum Optics in Phase Space* (Berlin: Wiley)
- [14] Klauder J R and Skagerstam B-S 1985 *Coherent States. Application in Physics and Mathematical Physics* (Singapore: World Scientific)
- [15] Bjorken J D and Drell S D 1993 *Relativistic Quantum Fields* (New York: McGraw-Hill)
- [16] Mattuck R D 1992 *A Guide to Feynman Diagrams in the Many-Body Problem* 2nd edn (New York: Dover)
- [17] Blasiak P, Horzela A, Penson K A, Solomon A I and Duchamp G H E 2007 Combinatorics and boson normal ordering: a gentle introduction *Am. J. Phys.* **75** 639–46 (arXiv:0704.3116)
- [18] Navon A M 1973 Combinatorics and fermion algebra *Nuovo Cimento* **16B** 324–30
- [19] Solomon A I, Duchamp G H E, Blasiak P, Horzela A and Penson K A 2004 Normal order: combinatorial graphs *Proc. 3rd Int. Symp. on Quantum Theory and Symmetries (Cincinnati 2003)* (Singapore: World Scientific) pp 527–36 (arXiv:quant-ph/0402082)
- [20] Varvak A 2005 Rook numbers and the normal ordering problem *J. Comb. Theory A* **112** 292–307
- [21] Katriel J and Duchamp G 1995 Ordering relations for  $q$ -boson operators, continued fraction techniques and the  $q$ -BCH enigma *J. Phys. A: Math. Gen.* **28** 7209–25
- [22] Méndez M A, Blasiak P and Penson K A 2005 Combinatorial approach to generalized Bell and Stirling numbers and boson normal ordering problem *J. Math. Phys.* **46** 083511 (arXiv:quant-ph/0505180)
- [23] Blasiak P and Horzela A 2008 Graphs for Quantum Theory arXiv:0710.0266
- [24] Blasiak P Urn models & operator ordering procedures *Oberwolfach Preprints OWP* 2008-06, 2008. Available at <http://www.mfo.de/publications/owp>