

On the Kertész line: thermodynamic versus geometric criticality

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2008 J. Phys. A: Math. Theor. 41 085001

(<http://iopscience.iop.org/1751-8121/41/8/085001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.153

The article was downloaded on 03/06/2010 at 07:28

Please note that [terms and conditions apply](#).

On the Kertész line: thermodynamic versus geometric criticality

Ph Blanchard¹, D Gandolfo², L Laanait³, J Ruiz² and H Satz¹

¹ Fakultät für Physik, Universität Bielefeld, D-33501, Bielefeld, Germany

² Centre de Physique Théorique, UMR 6207, Universités Aix-Marseille et Sud Toulon-Var, Luminy Case 907, 13288 Marseille, France

³ Ecole Normale Supérieure de Rabat, BP 5118, Rabat, Morocco

Received 16 October 2007, in final form 13 December 2007

Published 12 February 2008

Online at stacks.iop.org/JPhysA/41/085001

Abstract

The critical behavior of the Ising model in the absence of an external magnetic field can be specified either through spontaneous symmetry breaking (thermal criticality) or through cluster percolation (geometric criticality). We extend this to finite external fields for the case of the Potts' model, showing that a geometric analysis leads to the same first-order/second-order structure as found in thermodynamic studies. We calculate the Kertész line, separating percolating and non-percolating regimes, both analytically and numerically for the Potts model in the presence of an external magnetic field.

PACS numbers: 05.50.+q, 64.60.C, 75.10.H, 05.70.Fh, 05.10.Ln

(Some figures in this article are in colour only in the electronic version)

1. Introduction

The critical behavior in certain spin systems, such as the Ising model, can be specified in two equivalent, though conceptually quite different ways. In the absence of an external magnetic field, decreasing the temperature leads eventually to the onset of spontaneous symmetry breaking and hence to the singular behavior of derivatives of the partition function. On the other hand, the average size of clusters of like-sign spins also diverges at a certain temperature, i.e., there is an onset of percolation. The relation between these two distinct forms of singular behavior has been studied extensively over the years, and it was shown that for the Ising model on the lattice \mathbb{Z}^d , with $d \geq 2$, implemented with a suitable cluster definition using temperature dependent bond weights, the two forms lead to the same criticality: the critical temperatures T_c as well as the corresponding critical exponents coincide in the two formulations [4, 7].

In the presence of an external field H , the Z_2 symmetry of the Ising model is explicitly broken and hence there is no more thermodynamic critical behavior. Geometric critical behavior persists, however; for $T \leq T_p(H)$, there is percolation, while for $T > T_p(H)$, the




Figure 1. Thermodynamic and geometric phase structure for a first-order transition.

average cluster size remains finite. In the $T - H$ plane, there thus exists a line $T_p(H)$, the so-called Kertész line, separating a percolating from a non-percolating ‘phase’ [11]. Given the mentioned correct cluster definition, it starts at $T_p(0) = T_c$, i.e., at the thermodynamic critical point.

We want to show here that the equivalence of thermodynamic and geometric critical behavior can be extended to the case $H \neq 0$. Since in the case of continuous thermodynamic transitions, such as those of the Ising model, the introduction of an external field excludes singular behavior (for the case $d = 2$ this is shown analytically [6, 9]), our problem makes sense only for first-order transitions, for which the discontinuity remains over a certain range of H , even though for $H \neq 0$, the symmetry is broken. The ideal tool for such a study is the q -state Potts’ model on a lattice \mathbb{Z}^d , with $q \geq 3$ and $d \geq 3$. In this case, we have a thermodynamic phase diagram of the type shown in figure 1(a), with a line of first-order transitions starting at $T_c(0)$ and ending at a second-order point $T_c(H_c)$ [10]; the transition at this endpoint is found to be in the universality class of the 3D Ising model. In terms of the energy density $\epsilon(H)$ of the system (the energy per lattice volume), the phase diagram has the form shown in figure 1(b); for $H = 0$, the coexistence range $\epsilon_2 \leq \epsilon(0) \leq \epsilon_1$ corresponds to the critical temperature $T_c(0)$. The average spin $m(\epsilon)$ as order parameter vanishes for $\epsilon \geq \epsilon_1$ and becomes finite for smaller ϵ . We want to show that in the temperature range $T_c(0) \leq T(H) \leq T_c(H_c)$, the corresponding Kertész line $T_p(H)$ (see figure 1(c)) coincides with that of the thermal discontinuity and that it also leads to the same first-order/second-order phase structure. Let us begin with a conceptual discussion of the situation.

The q -state Potts’ model in the absence of an external field provides $q + 1$ phases: the disordered phase at high temperature and q degenerate ordered low-temperature phases. Spontaneous symmetry breaking has the system fall into one of these as the temperature is

decreased. Turning on a small external field H aligns the spins in its direction and thus effectively removes the $q - 1$ ‘orthogonal’ low-temperature phases. Hence now only two phases remain: the ordered low-temperature state of spins aligned in the direction of H and the disordered high-temperature phase. The two are for $T < T_c(H_c)$ separated by a mixed-phase coexistence regime. At the endpoint $T = T_c(H_c)$, there is a continuous transition from a system in one (symmetry broken) ordered phase to the corresponding (symmetric) disordered phase. The behavior at $H = H_c$ in figure 1(b) is thus just that of the Ising model, and hence the endpoint transition is in its universality class.

In the geometric formulation for $H = 0$, with decreasing temperature or energy density there is formation of finite clusters of q different orientations; the clusters here are defined using the temperature-dependent F–K bond weights. At $\epsilon(0) = \epsilon_1$, the Z_q symmetry is spontaneously broken: for one of the q directions, there now are percolating clusters and the percolation strength $P(\epsilon)$ becomes finite for $\epsilon < \epsilon_1$. However, the disordered phase also still forms a percolating medium (for $d \geq 3$). A further decrease of the energy density reduces the fraction of space in disordered state, and for $\epsilon(0) \leq \epsilon_2$, there is no more disordered percolation. Embedded in the disordered phase are at all times finite clusters of a spin orientation ‘orthogonal’ to the one chosen by spontaneous symmetry breaking. In our treatment, we will therefore divide the set of clusters into three classes: disordered, ordered in the direction of symmetry breaking and ordered orthogonal to the latter. While for $H = 0$, any of the q directions could be the given orientation, for $H \neq 0$, the external field specifies the alignment direction, making the $q - 1$ sets of ‘orthogonal’ clusters essentially irrelevant. It is for this reason that at the endpoint of a line of first-order transitions one generally encounters the universality class of the Ising model. Whatever the original symmetry of the system was, at the endpoint there remains only the aligned and the disordered ground states.

The plan of the paper is as follows. In the following section, we recall the cluster treatment of the Potts’ model and specify our method to identify the different cluster types. This will be followed by an analytic study valid for small external fields and by numerical calculations for different q up to asymptotic values of H . Formal details of the analytic calculation are given in the appendix.

2. The model

We consider a finite-volume q -state Potts model on the lattice \mathbb{Z}^d ($d \geq 2$), at inverse temperature $\beta = 1/T$ and subject to an external ordering field h . It is defined by the Boltzmann weight

$$\omega_{\text{Potts}}(\boldsymbol{\sigma}) = \prod_{\langle i,j \rangle} e^{\beta(\delta_{\sigma_i, \sigma_j} - 1)} \prod_i e^{h\delta_{\sigma_i, 1}}, \quad (1)$$

where the spins σ_i take on the values of the set $\{1, \dots, q\}$, and where the first product is over nearest neighbor pairs (n, n) . If we want to study the behavior of clusters, in the sense of F–K clusters, we turn to the corresponding Edwards–Sokal formulation [5], given by the Boltzmann weight

$$\omega_{\text{ES}}(\boldsymbol{\sigma}, \boldsymbol{\eta}) = \prod_{\langle i,j \rangle} [e^{-\beta} \delta_{\eta_{ij}, 0} + (1 - e^{-\beta}) \delta_{\eta_{ij}, 1} \delta_{\sigma_i, \sigma_j}] \prod_i e^{h\delta_{\sigma_i, 1}}, \quad (2)$$

where the edge variables η_{ij} belong to $\{0, 1\}$. This ‘site-bond’ model can be thought of as follows. Given a certain spin configuration, one puts between two neighboring sites $\sigma_i = \sigma_j$ an edge or bond with the probability $1 - e^{-\beta}$, and no edge with the probability $e^{-\beta}$; for $\sigma_i \neq \sigma_j$, no bond is present. When the field is infinite, all $\sigma_i = 1$, and we are left with a classical bond

percolation problem, while for finite field, one has a random bond percolation model in the random media given by the spin configuration.

In the presence of an external field, we find it convenient for the study of the Kertész's line to consider a modified version of the Edwards–Sokal formulation. We have three different types of spin combination: (0) two adjacent spins i, j are not equal, (1) two adjacent spins i, j are equal and parallel to h (we denote this direction as 1) or (2) two adjacent spins i, j are equal but not parallel to h . Correspondingly, we 'color' the edge between i and j in three different colors $n_{i,j}$, where the edge variables n_{ij} belong to $\{0, 1, 2\}$. The resulting Boltzmann weight becomes

$$\omega_{\text{CES}}(\boldsymbol{\sigma}, \mathbf{n}) = \prod_{(i,j)} \left[e^{-\beta} \delta_{n_{ij},0} + (1 - e^{-\beta}) \delta_{n_{ij},1} \chi(\sigma_i = \sigma_j = 1) + (1 - e^{-\beta}) \delta_{n_{ij},2} \chi(\sigma_i = \sigma_j \neq 1) \right] \prod_i e^{h \delta_{\sigma_i,1}}, \quad (3)$$

where the characteristic function $\chi(\sigma_i = \sigma_j = 1)$ is unity for $\sigma_i = \sigma_j = 1$ (parallel spins in the direction of h) and zero otherwise, while $\chi(\sigma_i = \sigma_j \neq 1)$ is unity for parallel spins not in the direction of h and zero otherwise. The summation over the spin variables then leads to the following tricolor-edge representation:

$$\omega_{\text{TER}}(\mathbf{n}) = \prod_{(i,j)} e^{-\beta \delta_{n_{ij},0}} (1 - e^{-\beta})^{(\delta_{n_{ij},1} + \delta_{n_{ij},2})} e^{h S_1(\mathbf{n})} (q - 1)^{C_2(\mathbf{n})} (q - 1 + e^h)^{|\Lambda| - S_1(\mathbf{n}) - S_2(\mathbf{n})}. \quad (4)$$

Here, $S_1(\mathbf{n})$ and $S_2(\mathbf{n})$ denote the number of sites that belong to edges of color 1 and color 2, respectively, while $C_2(\mathbf{n})$ denotes the number of connected components of the set of edges of color 2, and $|\Lambda|$ is the number of sites of the lattice under consideration.

Let us mention that such kind of graphical representation has already been considered for various spin models in the presence of an external field [3].

Let $p_\Lambda(i \leftrightarrow j)$ be the probability that the site i is connected to j by a path of edges of color 1. As (geometric) order parameter we will consider the following mass gap (inverse correlation length):

$$m(\beta, h) = - \lim_{|i-j| \rightarrow \infty} \frac{1}{|i-j|} \ln \lim_{\Lambda \uparrow \mathbb{Z}^d} p_\Lambda(i \leftrightarrow j), \quad (5)$$

where i and j belong to some line parallel to an axis of the lattice. As (thermodynamic) order parameter, we shall consider the mean energy $E(\beta, h) = -\frac{1}{\beta} \frac{\partial}{\partial \beta} f(\beta, h)$, where $f(\beta, h)$ is the free energy of the model⁴.

3. Analytic results

Let us first have a look at the diagram of ground-state configurations of the TER representation which are the translation invariant configurations maximizing the Boltzmann weight (4).

For the color $a = 0, 1, 2$, let b_a be the value of the Boltzmann weight of the ground-state configuration of color a per unit site. One finds $b_0 = e^{-\beta d} (q - 1 + e^h)$, $b_1 = (1 - e^{-\beta})^d e^h$, $b_2 = (1 - e^{-\beta})^d$.

Note that $b_0 = b_1$ on the line

$$\beta_0(h) = \ln[1 + (1 + (q - 1) e^{-h})^{1/d}] \quad (6)$$

and that $b_0 = b_1 = b_2$ at the point $\beta_0(0)$.

The diagram of ground-state configurations, inferred from the values of the weights b_0, b_1, b_2 is shown in figure 2 (in the (h, β) plane).

⁴ Note that all partition functions, and hence the free energies, of models (1)–(4) coincide.




Figure 2. Diagram of ground-state configurations: all the ground-state configurations coexist at $(0, \beta_0(0))$. Below $\beta_0(h)$, the 0-state dominates. Above $\beta_0(h)$, the 1-state dominates; it coexists with the 0-state on the line $\beta_0(h)$, and with the 2-state on the line $h = 0, \beta \geq \beta_0(0)$.

When q is large enough and h not too large, the TER representation (4) can be analyzed rigorously by a perturbative approach. Namely, by using the standard machinery of Pirogov–Sinai theory, we will show that the model undergoes a thermodynamic first-order phase transition in the sense that the mean energy (as well as the magnetization) is discontinuous at some $\beta_c(h) \sim \beta_0(h)$. We also find for these values of the parameters, that the phase diagram of this model reproduces the diagram of ground-state configurations (figure 2), see the appendix for more details.

In addition, the model exhibits a geometric (first-order) transition, in the sense that, on the same critical line, the mass gap is discontinuous.

Theorem 1. Assume $d \geq 2, q$ and h such that

$$c_d(1 + (q - 1)e^{-h})^{-1/2d} < 1 \tag{7}$$

holds, where c_d is a given number (depending only on the dimension), then there exists a unique $\beta_c(h) = \beta_0(h) + O(1 + (q - 1)e^{-h})^{-1/2d}$ such that

- (1) $\Delta E(\beta_c(h), h) = E(\beta_c^-(h), h) - E(\beta_c^+(h), h) > 0$
- (2) $m(\beta, h) > 0$ for $\beta \leq \beta_c(h)$ and $m(\beta, h) = 0$ for $\beta > \beta_c(h)$.

The proof is given in the appendix.

Let us recall that it has already been shown that the Potts model (1) undergoes, for q large and h small, a first-order phase transition on a critical line [1], where both the mean energy and the magnetization are discontinuous. Since, as already mentioned, the free energies of models (1) and (4) are the same, this critical line coincides with the one mentioned in the theorem.

In the absence of an external magnetic field, the statements of the theorem have been shown previously [12–14].

Condition (7) restricts the range of values of parameters to which our rigorous analysis applies. Moreover, we do not expect thermodynamic first-order transitions when h is sufficiently enhanced. In the following section, we turn to numerical study on a wider range of values.




Figure 3. $\beta_c(h)$ for several values of q , with ‘first-order’ behavior in red and ‘second order’ in blue. The first-order behavior is both thermodynamic and geometric. The second-order behavior is only geometric.

4. Numerical simulations

We have implemented a generalization of the Swendsen–Wang algorithm for our colored Edwards–Sokal model (3).

First, given a spin configuration, we put between any two neighboring spins of the same color, an edge colored 0 with probability (w.p.) $e^{-\beta}$, and w.p. $1 - e^{-\beta}$, an edge colored 1 if these spins are of color 1, and colored 2 otherwise. When two neighboring spins disagree, the corresponding edge is colored 0.

Then, starting from an edge configuration, a spin configuration is constructed as follows. Isolated sites (endpoints of 0-bonds only) are colored 1 w.p. $e^h/(q - 1 + e^h)$ and colored $c \in \{2, \dots, q\}$ w.p. $1/(q - 1 + e^h)$. Non-isolated sites are colored 1 (w.p. 1) if they are endpoints of 1-bonds and colored $c \in \{2, \dots, q\}$ w.p. $1/(q - 1)$.

This algorithm allows us to compute both quantities associated with spin configurations and those associated with edge configurations.

The numerical results for $d = 2$ are presented in figure 3. For $q \leq 4$, we found a whole geometric transition line $\beta_c(h)$ for which $m(\beta, h) > 0$ when $\beta < \beta_c(h)$, and $m(\beta, h) = 0$ when $\beta \geq \beta_c(h)$. The mass gap is continuous at $\beta_c(h)$. For $\beta < \beta_c(h)$, the mean cluster sizes remain finite, while for $\beta \geq \beta_c(h)$ the size of 1-edge clusters diverges. The energy density as well as the magnetization do not show any singular behavior.

For $q \geq 5$, some critical h_c appears for which the geometric transition line $\beta_c(h)$ becomes first order when $h < h_c$, i.e. $m(\beta, h) > 0$ for $\beta \leq \beta_c(h)$ and $m(\beta, h) = 0$ for $\beta > \beta_c(h)$. In addition, on this part of the line, we find that the mean energy is discontinuous⁵. When $h \geq h_c$, only a geometric transition occurs and the scenario is the same as for $q \leq 4$. Thus our numerics show that the geometric and thermodynamic transitions coincide up to h_c , similarly to what we got analytically but only at (very) small field (and large q), see figure 3.

⁵ The Swendsen–Wang algorithm allows us to compute both associated order parameters (mass gap and mean energy).

Let us mention that the numerics are in accordance with the theory for vanishing and infinite fields: $\beta_c(0) = \ln(1 + \sqrt{q})$ and $\beta_c(\infty) = \ln 2$.

The system size in these calculations was $L = 50, d = 2$. The ‘first-order’ part of the transition lines has been determined via Binder cumulants [2]. The Hoshen–Kopelman algorithm [8] was used to study cluster statistics. For each value of q , more than 2×10^5 iterations were performed. Data have been binned in order to control errors in measurements.

5. Concluding remarks

For the Potts model in the presence of an external magnetic field, we have shown that when the Kertész line is first order, it coincides with the usual thermodynamic critical line. This property holds up to some critical point $(h_c, \beta_c(h_c))$, beyond which the thermodynamic disappears. Such behavior may well appear also for a broader class of models exhibiting first-order transition in the presence of an external field. We believe that the behavior at the above critical point also belongs to the universality class of the Ising model, as it is the case in the 3-state Potts model in three dimensions [10].

Acknowledgments

It is a pleasure to thank János Kertész for interesting comments. The BIBOS research center (Bielefeld) and the Centre de Physique Théorique (CNRS Marseille) are gratefully acknowledged for warm hospitality and financial support. The authors are indebted to an anonymous referee for constructive remarks.

Appendix

We first introduce the partition function of the TER representation with boundary conditions $a \in \{0, 1, 2\}$ in a box Λ ⁶:

$$Z_a(\Lambda) = \sum_{\mathbf{n}} \prod_{i \in \Lambda} \omega_i(\mathbf{n}) q^{C_2(\mathbf{n}) - \delta_{a,2}} \prod_{i \in \partial \Lambda} \prod_{j \sim i} \delta_{n_{ij}, a}, \tag{A.1}$$

where the sum is over all configurations $\mathbf{n} = \{n_{ij}\}_{ij \cap \Lambda \neq \emptyset}$, $\partial \Lambda$ is the boundary of Λ (set of sites of Λ with a n.n. in $\mathbb{Z}^d \setminus \Lambda$), the notation $i \sim j$ means that i and j are n.n., and

$$\omega_i(\mathbf{n}) = (1 - e^{-\beta})^{(\delta_{n_{ij},1} + \delta_{n_{ij},2})/2} e^{-\beta \delta_{n_{ij},0}/2} e^{h \chi(i \in '1')} (q - 1 + e^h)^{\prod_{j \sim i} \delta_{n_{ij},0}}, \tag{A.2}$$

where $\chi(i \in '1')$ means that the site i belongs to some edge of color 1. Next, consider a configuration \mathbf{n} on the envelope of Λ : $E(\Lambda) = \{\langle i, j \rangle \cap \Lambda \neq \emptyset\}$. A site $i \in \Lambda$ is called *correct* if for all $j \sim i$, n_{ij} takes the same value, and called *incorrect* otherwise. Denote $I(\mathbf{n})$ the set of incorrect sites of the configuration \mathbf{n} . A couple $\Gamma = \{\text{Supp } \Gamma, \mathbf{n}(\Gamma)\}$ where the support of Γ ($\text{Supp } \Gamma$) is a maximal connected subset of $I(\mathbf{n})$, and $\mathbf{n}(\Gamma)$ the restriction of \mathbf{n} to the envelope of Λ is called *contour* of the configuration \mathbf{n} (here, a set of sites is called connected if the graph that joins all the sites of this set at distance $d(i, j) = \max_{k=1, \dots, d} |i_k - j_k| \leq 1$ is connected). A couple $\Gamma = \{\text{Supp } \Gamma, \mathbf{n}(\Gamma)\}$ where $\text{Supp } \Gamma$ is a connected set of sites is called contour if there exists a configuration \mathbf{n} such that Γ is a contour of \mathbf{n} . For a contour Γ , let \mathbf{n}_Γ denote the configuration having Γ as unique contour, $\text{Ext } \Gamma$ denotes the unique infinite component of $\mathbb{Z}^d \setminus (\text{Supp } \Gamma)$, $\text{Int } \Gamma = \mathbb{Z}^d \setminus (\text{Ext } \Gamma \cup \text{Supp } \Gamma)$ and $\text{Int}_m \Gamma$ denote the set of sites of

⁶ The reader should not be confused by the fact that (A.1), called diluted partition function in PS theory, differs from the usual one by an unimportant boundary term which makes the expansions (A.3) and (A.4) easier to write.

Int Γ corresponding to the color $m \in \{0, 1, 2\}$ for the configuration \mathbf{n}_Γ . Two contours Γ_1 and Γ_2 are said to be compatible if their union is not connected and are called external contours if furthermore $\text{Int } \Gamma_1 \subset \text{Ext}_\Lambda \Gamma_2$ and $\text{Int } \Gamma_2 \subset \text{Ext}_\Lambda \Gamma_1$. For a family $\theta = \{\Gamma_1, \dots, \Gamma_n\}_{\text{ext}}$ of external contours, let $\text{Ext}_\Lambda \theta$ denote the intersection $\Lambda \cap \bigcap_{k=1}^n \text{Ext}_\Lambda \Gamma_k$. With these definitions and notations, one gets the following expansion of the partition functions over families of external contours:

$$Z_a(\Lambda) = \sum_{\theta = \{\Gamma_1, \dots, \Gamma_n\}_{\text{ext}}} b_a^{|\text{Ext}_\Lambda \theta|} \prod_{k=1}^n \rho(\Gamma_k) \prod_{m=0,1,2} Z_m(\text{Int}_m \Gamma_k), \quad (\text{A.3})$$

where $\rho(\Gamma) = \prod_{i \in \text{supp } \Gamma} \omega_i(\mathbf{n}_\Gamma) q^{C(\mathbf{n}_\Gamma) - \delta_{a,2}}$. From (A.3), we get

$$Z_a(\Lambda) = b_a^{|\Lambda|} \sum_{\{\Gamma_1, \dots, \Gamma_n\}_{\text{comp}}} \prod_{k=1}^n z_a(\Gamma_k), \quad (\text{A.4})$$

where the sum is now over families of compatible contours and the activities $z_a(\Gamma)$ of contours are given by $z_a(\Gamma) = \rho(\Gamma) b_a^{-|\text{supp } \Gamma|} \prod_{m \neq a} \frac{Z_m(\text{Int}_m \Gamma)}{Z_a(\text{Int}_m \Gamma)}$.

It is easy to prove the following Peierls' estimate:

$$\rho(\Gamma) (\max_{a=0,1,2} b_a)^{-|\text{Supp } \Gamma|} \leq e^{-\tau |\text{Supp } \Gamma|}, \quad (\text{A.5})$$

where $e^{-\tau} = (1 + (q - 1) e^{-h})^{-1/2d}$. Indeed, first note that an incorrect site i is either of color 1 or of color 2. In the first case one has $\sum_{j \sim i} (\delta_{n_{ij},0} + \delta_{n_{ij},1}) = 2d$, so that $\omega_i(\mathbf{n}_\Gamma)/b_1 = (e^\beta - 1)^{-\sum_{j \sim i} \delta_{n_{ij},0}/2}$, implying

$$\omega_i(\mathbf{n}_\Gamma) / \max_{a=0,1,2} b_a \leq (1 + (q - 1) e^{-h})^{-\sum_{j \sim i} \delta_{n_{ij},0}/2d}.$$

Thus since $1 \leq \sum_{j \sim i} \delta_{n_{ij},0} \leq 2d - 1$, each incorrect site of color 1 gives at most a contribution $e^{-\tau}$ to the LHS of (A.5). In the second case, one has $\sum_{j \sim i} (\delta_{n_{ij},0} + \delta_{n_{ij},2}) = 2d$, so that $\omega_i(\mathbf{n}_\Gamma)/b_2 = (e^\beta - 1)^{-\sum_{j \sim i} \delta_{n_{ij},0}/2}$, implying

$$\omega_i(\mathbf{n}_\Gamma) / \max_{a=0,1,2} b_a \leq (q - 1 + e^h)^{-\sum_{j \sim i} \delta_{n_{ij},0}/2d}.$$

We then use again that $1 \leq \sum_{j \sim i} \delta_{n_{ij},0} \leq 2d - 1$ and that $C_2(\mathbf{n}_\Gamma) \leq \sum_{i \in \text{Supp } \Gamma} \chi(1 \leq \delta_{n_{ij},2})/2^{\sum_{j \sim i} \delta_{n_{ij},2}}$ (see [12]) to obtain that each incorrect site of color 2 gives at most a contribution $(e^h + q - 1)^{-1/2+1/2d} \leq e^{-\tau}$ to the LHS of (A.5).

When the assumptions of the theorem are satisfied, the Peierls' estimate (A.5) provides a good control of the system by using Pirogov–Sinai theory [15]. We introduce the truncated activity

$$z'_a(\Gamma) = \begin{cases} z_a(\Gamma) & \text{if } z_a(\Gamma) \leq (c_0 e^{-\tau})^{|\text{Supp } \Gamma|} \\ (c_0 e^{-\tau})^{|\text{Supp } \Gamma|} & \text{otherwise,} \end{cases}$$

where c_0 is a numerical constant, and we call a contour stable if $z_a(\Gamma) = z'_a(\Gamma)$. Let $Z'_a(\Lambda)$ be the partition function obtained from (A.4) by leaving out unstable contours, i.e., by taking the activities $z'_a(\Gamma)$ in (A.4), and let us introduce the metastable free energies $f_a^{\text{met}}(\beta, h) = -\lim_{\Lambda \uparrow \mathbb{Z}^d} (1/|\Lambda|) \ln Z'_a(\Lambda)$. The leading term of these metastable free energies equals $-\ln b_a$. The corrections can be expressed by free energies of contour models which can be controlled by convergent cluster expansions. As a standard result of Pirogov–Sinai theory, one gets that the phase diagram of the system is a small perturbation of the diagram of ground-state configurations. Namely, there exists a unique point $\beta_c(0)$ given by the solution of $f_0^{\text{met}}(\beta, h) = f_1^{\text{met}}(\beta, h) = f_2^{\text{met}}(\beta, h)$ for which all contours are stable and such that

$Z_a(\Lambda) = Z'_a(\Lambda)$ for $a = 0, 1, 2$. There exists a line $\beta_c(h)$ given by the solution of $f_0^{\text{met}}(\beta, h) = f_1^{\text{met}}(\beta, h)$ when $h > 0$ and such that, $Z_a(\Lambda) = Z'_a(\Lambda)$ for $a = 0, 1$. For $\beta < \beta_c(h)$ one has $Z_0(\Lambda) = Z'_0(\Lambda)$, and for $\beta > \beta_c(h)$ one has $Z_1(\Lambda) = Z'_1(\Lambda)$. For $h = 0$ and $\beta \geq \beta_c(0)$, one has in addition $Z_2(\Lambda) = Z'_2(\Lambda)$.

For the color $a = 0, 1, 2$, denote by $\langle \cdot \rangle^a(\beta, h)$ the expectation value under the a -boundary condition. As a consequence of the above expansions and analysis, we obtain by standard Peierls' estimates that for $h \geq 0$,

$$\langle \delta_{nij,1} \rangle^1(\beta, h) \geq 1 - O(e^{-\tau}) \quad \text{for } \beta \geq \beta_c(h) \quad (\text{A.6})$$

$$\langle \delta_{nij,0} \rangle^0(\beta, h) \geq 1 - O(e^{-\tau}) \quad \text{for } \beta \leq \beta_c(h), \quad (\text{A.7})$$

while in addition we also get for $h = 0$,

$$\langle \delta_{nij,2} \rangle^2(\beta, 0) \geq 1 - O(e^{-\tau}) \quad \text{for } \beta \geq \beta_c(0). \quad (\text{A.8})$$

By definition of the mean energy, one has that $\Delta E = E(\beta^-, h) - E(\beta^+, h)$ is proportional to the difference $\langle \delta_{nij,0} \rangle^0(\beta^-, h) - \langle \delta_{nij,0} \rangle^1(\beta^+, h)$, and the first statement of the theorem follows immediately from these properties.

To prove the second statement, we remark that if one imposes that the site i is connected to j by a path made up of edges of color 1, then under the boundary condition 0, there exists necessarily an external contour that encloses both the sites i and j . As a consequence of the above analysis the probability of external contours Γ decays like $(c_0 e)^{-\tau|\text{Supp } \Gamma|}$ when the 0-contours are stable, i.e. when $Z_0(\Lambda) = Z'_0(\Lambda)$. One thus gets $p_\Lambda(i \leftrightarrow j) \leq (\text{Cte } e^{-\tau})^{|i-j|}$ when $\beta \leq \beta_c(h)$ from which the first statement of the theorem follows. On the other hand, under the boundary condition 1, the probability that the site i is not connected to j can be bounded from above by a small number $O(e^{-\tau})$ when $Z_1(\Lambda) = Z'_1(\Lambda)$. This also follows from a Peierls-type arguments and implies that the probability that the site i is connected to j under the boundary condition 1 is greater than $1 - O(e^{-\tau})$ for $\beta \geq \beta_c(h)$. It gives also that the probability $p_\Lambda(i \leftrightarrow j)$ for the site i to be connected with j under the boundary condition 0 is also greater than $1 - O(e^{-\tau})$ for $\beta > \beta_c(h)$, implying the second statement.

References

- [1] Bakchich A, Benyoussef A and Laanait L 1989 *Ann. Inst. Henri Poincaré* **50** 17
- [2] Binder K and Landau D P 1984 *Phys. Rev. B* **30** 1477
Challa M S S, Binder K and Landau D P 1986 *Phys. Rev. B* **34** 1841
- [3] Chayes L and Machta J 1997 *Physica A* **239** 542
- [4] Coniglio A and Klein W 1980 *J. Phys. A: Math. Gen.* **13** 2775
- [5] Edwards R G and Sokal A D 1988 *Phys. Rev. D* **38** 2009
- [6] Friedli S and Pfister C-E 2004 *Phys. Rev. Lett.* **92** 015702
Friedli S and Pfister C-E 2004 *Commun. Math. Phys.* **245** 69
- [7] Fortuin C M and Kasteleyn P W 1969 *J. Phys. Soc. Japan* **26** 11
Fortuin C M and Kasteleyn P W 1972 *Physica* **57** 536
- [8] Hoshen J and Kopelman R 1976 *Phys. Rev. B* **14** 3438
- [9] Isakov S N 1984 *Commun. Math. Phys.* **95** 427
- [10] Karsch F and Stickan S 2000 *Phys. Lett. B* **488** 319
- [11] Kertész J 1989 *Physica A* **161** 58
- [12] Kotecký R, Laanait L, Messenger A and Ruiz J 1990 *J. Stat. Phys.* **58** 199
- [13] Laanait L, Messenger A, Miracle-Solé S, Ruiz J and Shlosman S 1991 *Commun. Math. Phys.* **140** 81
- [14] Laanait L, Messenger A and Ruiz J 1986 *Commun. Math. Phys.* **105** 527
- [15] Sinai Ya G 1982 *Theory of Phase Transitions: Rigorous Results* (Oxford: Pergamon)
Zahradník M 1984 *Commun. Math. Phys.* **93** 359