

Walking on fractals: diffusion and self-avoiding walks on percolation clusters

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2009 J. Phys. A: Math. Theor. 42 015001

(<http://iopscience.iop.org/1751-8121/42/1/015001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.153

The article was downloaded on 03/06/2010 at 07:30

Please note that [terms and conditions apply](#).

Walking on fractals: diffusion and self-avoiding walks on percolation clusters

V Blavatska^{1,2} and W Janke¹

¹ Institut für Theoretische Physik and Centre for Theoretical Sciences (NTZ),
Universität Leipzig, Postfach 100 920, D-04009 Leipzig, Germany

² Institute for Condensed Matter Physics, National Academy of Sciences of Ukraine,
UA-79011 Lviv, Ukraine

E-mail: Viktoria.Blavatska@itp.uni-leipzig.de, viktoria@icmp.lviv.ua and
Wolfhard.Janke@itp.uni-leipzig.de

Received 24 July 2008

Published 19 November 2008

Online at stacks.iop.org/JPhysA/42/015001

Abstract

We consider random walks (RWs) and self-avoiding walks (SAWs) on disordered lattices directly at the percolation threshold. Applying numerical simulations, we study the scaling behavior of the models on the incipient percolation cluster in space dimensions $d = 2, 3, 4$. Our analysis yields estimates of universal exponents, governing the scaling laws for configurational properties of RWs and SAWs.

PACS numbers: 67.80.dj, 36.20.-r, 64.60.ah, 07.05.Tp

1. Introduction

The model of a random walk (RW) provides a good description of diffusion processes, such as, for example, encountered for electrons in metals or colloidal particles in solution [1]. The averaged mean-square displacement of the diffusive particle at time t (or, equivalently, after $t = N$ steps on a lattice) scales as

$$\langle R^2 \rangle \sim t^{2\nu_{\text{RW}}}, \quad (1)$$

where in a non-disordered medium $\nu_{\text{RW}} = 1/2$, independently of the space dimension d . A RW is a fractal object, with fractal dimension $d_{\text{RW}} = 1/\nu_{\text{RW}}$. The number of all possible trajectories C_N for a randomly walking particle of N steps can be found exactly: $C_N = z_0^N$, where z_0 is the coordination number of the corresponding lattice.

Forbidding the trajectory of a random walk to cross itself, we obtain a self-avoiding walk (SAW), which is one of the most successful in describing the universal configurational properties of a long, flexible single polymer chain in good solvent [2]. The average squared

Table 1. Critical concentration p_c of site-diluted lattices and fractal dimensions of percolation clusters, $d_{p_c}^F$, and the backbone of percolation clusters, $d_{p_c}^B$, for different space dimensions d .

d	p_c	$d_{p_c}^F$	$d_{p_c}^B$
2	0.592 746 [14]	91/49 [13]	1.650 ± 0.005 [18]
3	0.311 60 [15]	2.51 ± 0.02 [17]	1.86 ± 0.01 [18]
4	0.196 88 [16]	3.05 ± 0.05 [17]	1.95 ± 0.05 [18]

end-to-end distance $\langle R^2 \rangle$ and the number of configurations C_N of SAWs with N steps on the underlying lattice obey the scaling laws:

$$\langle R^2 \rangle \sim N^{2\nu_{\text{SAW}}}, \quad C_N \sim z^N N^{\gamma_{\text{SAW}}-1}, \quad (2)$$

where ν_{SAW} , γ_{SAW} are universal exponents that only depend on the space dimension d , and z is a non-universal fugacity, counting the average number of accessible nearest-neighbor sites. The properties of SAWs on a regular lattice have been studied in detail both in analytical approaches [3–6] and computer simulations [7–12]. For example, in the space dimension $d = 3$ one finds within the frame of the field-theoretical renormalization group approach $\nu_{\text{SAW}} = 0.5882 \pm 0.0011$ [6] and Monte Carlo simulations give $\nu_{\text{SAW}} = 0.5877 \pm 0.0006$ [11]. For space dimensions d above the upper critical dimension $d_{\text{up}} = 4$, the effect of self-avoidance becomes irrelevant and SAWs behave effectively as random walks with exponents $\nu_{\text{RW}} = 1/2$, $\gamma_{\text{RW}} = 1$.

The problem of random walks in disordered media is of great interest since it is connected with a large amount of physical phenomena: transport properties in fractures and porous rocks, the anomalous density of states in randomly diluted magnetic systems, silica aerogels and in glassy ionic systems, diffusion-controlled fusion of excitations in porous membrane films, etc (see, e.g., [13] for a review). Similarly, SAWs on randomly diluted lattices may serve as a model of linear polymers in a porous medium.

Much of our understanding of disordered systems comes from percolation theory [19]. A disordered medium can be modelled as a randomly diluted lattice, with a given concentration p of lattice sites allowed for walking. Most interesting is the case, when p equals the critical concentration p_c , the site-percolation threshold (see table 1) and an incipient percolation cluster can be found in the system. Studying properties of percolative lattices, one encounters two possible statistical averages. In the first, one considers only percolation clusters with linear size much larger than the typical length of the physical phenomenon under discussion. The other statistical ensemble includes all the clusters which can be found in a percolative lattice. For the latter ensemble of all clusters, the walks can start on any of the clusters, and for an N -step walk, performed on the i th cluster, we have $\langle R^2 \rangle \sim l_i^2$, where l_i is the averaged size of the i th cluster. In what follows, we will be interested in the former case, when trajectories of walks reside only on the percolation cluster. In this regime, the scaling laws (1), (2) hold with new exponents $\nu_{\text{RW}}^{p_c} \neq \nu_{\text{RW}}$, $\gamma_{\text{RW}}^{p_c} \neq \gamma_{\text{RW}}$ [20–35], $\nu_{\text{SAW}}^{p_c} \neq \nu_{\text{SAW}}$, $\gamma_{\text{SAW}}^{p_c} \neq \gamma_{\text{SAW}}$ [36–58]. A hint to the physical understanding of this phenomenon is given by the fact that weak disorder does not change the dimension of a lattice, whereas the percolation cluster itself is a fractal object with fractal dimension $d_{p_c}^F$ dependent on d (see table 1). In this way, scaling law exponents of residing walks change with the dimension $d_{p_c}^F$ of the (fractal) lattice on which the walk resides. Since $d_{\text{up}} = 6$ for percolation [19], the exponents $\nu_{\text{SAW}}^{p_c}(d \geq 6) = 1/2$, $\gamma_{\text{SAW}}^{p_c}(d \geq 6) = 1$.

Our present paper aims to supplement the studies of random and self-avoiding walks on percolative lattices by obtaining numerical values for exponents, governing the scaling behavior of the models, up to $d = 4$ by computer simulations. The layout of the paper is




Figure 1. Procedure of site labeling and extracting the percolation cluster.

as follows: in the following section, we will present in detail the procedure of extracting the percolation cluster and its backbone on disordered lattices at the percolation threshold. In section 3, we describe the pruned-enriched Rosenbluth algorithm, applied to study the scaling of self-avoiding walks, and present the results obtained. In section 4, we consider the method for studying random walks on percolation clusters. In section 5, we end up by giving conclusions and an outlook.

2. Construction of percolation clusters

We consider site percolation on a regular lattice of edge length $L = 400, 200, 50$ in dimensions $d = 2, 3, 4$, respectively. Each site of the lattice is assigned to be occupied with probability p_c (values of critical concentration in different dimensions are given in table 1), and empty otherwise. To describe the procedure of extracting the percolation cluster, let us consider schematically the two-dimensional case. We apply an algorithm based on the one proposed by Hoshen and Kopelman [59]. As a first step, a label is prescribed to each of the occupied sites. Such a labeling process is regulated, we start, for example, from the first ‘column’ of the lattice, label the occupied sites upward, and then turn to the next ‘column’, as shown in figure 1, left. Next, we start the procedure of burning the occupied sites. Again, in the same order, starting from the bottom of the first ‘column’ of the lattice, for each of the labeled sites (say, i), we check whether its nearest neighbors are also occupied or not. If yes, two possibilities appear: (1) the label of the neighbor is larger than the label of site i . In this case, we change the label of the neighbor to that of site i . (2) The label of the neighbor is smaller than that of i . Then, we change the label of site i to that of the neighbor.

Such a burning procedure is applied until no more change of site labels is needed. As a result, we end up with groups of clusters of occupied sites with the same labels (figure 1, right). Finally, we check whether there exists a cluster that wraps around the lattice in all d directions. If yes, we have found the percolation cluster (figure 2). If not, this disordered lattice is rejected and a new one is constructed. Note that on finite lattices the definition of spanning clusters is not unique (e.g., one could consider clusters connecting only two opposite borders), but all definitions are characterized by the same fractal dimension and are thus equally legitimate. The here employed definition has the advantage of yielding the most isotropic clusters. Note also that directly at $p = p_c$ more than one spanning cluster can be found in the system, and the probability $P(k)$ for at least k separated clusters grows with the space dimension as $P(k) \sim \exp(-\alpha k^{d/(d-1)})$ [60]. In our study, we take into account only one percolation




Figure 2. Percolation cluster on a $d = 2$ -dimensional regular lattice of edge length $L = 50$.




Figure 3. For all sites of a percolation cluster the chemical distance from the starting site is calculated. The minimal paths from all the sites on the edge of the percolation cluster to this starting point are found, which form the elastic backbone of the percolation cluster.

cluster per each disordered lattice constructed, in order to avoid presumable correlations of the data.

Aiming to investigate the scaling of SAWs on percolative lattices, we are interested in the backbone of percolation clusters, which can be defined as follows. Assume that each bond (or site) of the cluster is a resistor and that an external potential drop is applied at two ends of the cluster. The backbone is the subset of the cluster consisting of all bonds (or sites) through which the current flows; i.e., it is the structure left when all ‘dangling ends’ are eliminated from the cluster. The SAWs can be trapped in ‘dangling ends’, therefore infinitely long chains can only exist on the backbone of the cluster. The algorithm for extracting the backbone of obtained percolation clusters was first introduced in [61] and improved in [62]. We choose the




Figure 4. Elastic and geometrical backbones of the percolation cluster depicted in figure 2.

starting point—‘seed’—at the center of the cluster, and find the chemical distance l of all the sites belonging to the percolation cluster to this starting point. In figure 3, the starting point is numbered with 0, and the chemical distance of all the other sites are depicted. The burning algorithm is divided into two parts. First, we start from some site of the edge of the lattice belonging to the percolation cluster and consider it as a burning site. At the next step, if the nearest neighbor of this site has the chemical distance smaller than the burning site itself, the nearest neighbor site is burnt. Such a procedure ends when the ‘seed’ site is reached. All the thus obtained burnt sites are located along the shortest path between the ‘seed’ and the given site at the edge of the percolation cluster, as is shown in figure 3. The same algorithm is applied successively to all the edge sites. As a result, we obtain the so-called skeleton or elastic backbone [63], shown in figure 4, left. In the second part of the algorithm, we consider successively each site of the elastic backbone, and check whether a ‘loop’ starts from this site. A ‘loop’ is a path of sites, belonging to the percolation cluster, which is connected with the elastic backbone by no less than two sites. All sites of the elastic backbone together with the sites of ‘loops’ form finally the geometric backbone of the cluster (see figure 4, right).

The results for fractal dimensions of the percolation cluster and its geometrical backbone in $d = 2, 3, 4$ are compiled in table 1.

3. Self-avoiding walks on percolation clusters

3.1. The method

For the sampling of SAWs, we use the pruned-enriched Rosenbluth method (PERM), proposed in the work of Grassberger [64]. The algorithm is based on ideas from the very first days of Monte Carlo simulations, the Rosenbluth–Rosenbluth (RR) method [7] and enrichment [66]. Let us consider the growing polymer chain, i.e. the n th monomer is placed at a randomly chosen neighbor site of the last placed $(n-1)$ th monomer ($n \leq N$, where N is total length of polymer). In order to obtain correct statistics, if this new site is occupied, any attempt to place a monomer at it results in discarding the entire chain. This leads to an exponential ‘attrition’, the number of discarded chains grows exponentially with the chain length, which makes the method useless for long chains. In the RR method, occupied neighbors are avoided without discarding the chain, but the bias is corrected by means of giving a weight $W_n \sim \left(\prod_{l=1}^n m_l\right)$ to each sample configuration at the n th step, where m_l is the number of free lattice sites to place the l th monomer. When the chain of total length N is constructed, the new one starts from the same starting point, until the desired number of chain configurations are obtained.

The configurational averaging, e.g., for the end-to-end distance $r(N) \equiv \sqrt{R^2(N)}$, then has the form:

$$\langle r(N) \rangle = \frac{1}{Z_N} \sum_{\text{conf}} W_N^{\text{conf}} |\vec{r}_N - \vec{r}_0|, \quad Z_N = \sum_{\text{conf}} W_N^{\text{conf}}, \quad (3)$$

where \vec{r}_0 is the position of the starting point of the growing chains, \vec{r}_N is the position of the last monomer and Z_N is the partition sum.

The Rosenbluth method, however, also suffers from attrition: if all the next neighbors at some step ($n < N$) are occupied, i.e., the chain is running into a ‘dead end’, the complete chain has to be discarded and the growth process has to be restarted. Combining the chain growth algorithm with population control, such as PERM [64] leads to a considerable improvement of the efficiency by increasing the number of successfully generated chains. The weight fluctuations of the growing chain are suppressed in PERM by pruning configurations with too small weights, and by enriching the sample with copies of high-weight configurations [64]. These copies are made while the chain is growing and continue to grow independently of each other. Pruning and enrichment are performed by choosing thresholds $W_n^<$ and $W_n^>$ depending on the estimate of the partition sum for the n -monomer chain. These thresholds are continuously updated as the simulation progresses. The zeroes iteration is a pure chain-growth algorithm without reweighting. After the first chain of full length has been obtained, we switch to $W_n^<$, $W_n^>$. If the current weight W_n of an n -monomer chain is less than $W_n^<$, a random number $r = 0, 1$ is chosen; if $r = 0$, the chain is discarded, otherwise it is kept and its weight is doubled. Thus, low-weight chains are pruned with probability $1/2$. If W_n exceeds $W_n^>$, the configuration is doubled and the weight of each identical copy is taken as half the original weight. For a value of the weight lying between the thresholds, the chain is simply continued without enriching or pruning the sample. For updating the threshold values we apply similar rules as in [67, 68]: $W_n^> = C(Z_n/Z_1)(c_n/c_1)^2$ and $W_n^< = 0.2W_n^>$, where c_n denotes the number of created chains having length n , and the parameter C controls the pruning–enrichment statistics. After a certain number of chains of total length N is produced, the given tour is finished and a new one starts. We adjust the pruning–enrichment control parameter such that on average 10 chains of total length N are generated per each tour [68]. Also, what is even more important for efficiency is that in almost all iterations at least one such chain was created. The number of different trajectories of SAWs with N steps can be then estimated as averaged weight:

$$C_N = \frac{1}{t} \sum_{\text{conf}} W_N^{\text{conf}}, \quad (4)$$

where t is the number of successful tours. PERM has been applied to a wide class of problems, in particular studies of the Θ -transition in homopolymers [64, 65], trapping of random walkers on absorbing lattices [69], studies of protein folding [68, 70], etc.

3.2. Results

To study scaling properties of SAWs on percolating lattices, we have to perform two types of averaging: the first average is performed over all SAW configurations on a single percolation cluster according to (3); the second average is carried out over different realizations of disorder, i.e. over all percolation clusters constructed:

$$\overline{\langle r \rangle} = \frac{1}{M} \sum_{i=1}^M \langle r \rangle_i, \quad (5)$$




Figure 5. Disorder averaged end-to-end distance versus number of steps in double logarithmic scale for SAWs on the backbone of percolation clusters in $d = 2$ (pluses), $d = 3$ (stars), $d = 4$ (squares). Lines represent linear fitting, statistical error bars are of the size of symbols.

$$\overline{C_N} = \frac{1}{M} \sum_{i=1}^M C_{N,i}, \tag{6}$$

where M is the number of different clusters and the index i means that a given quantity is calculated on the cluster i .

The SAW statistics crucially depends on the type of disorder averaging, namely, whether the disorder is considered to be ‘annealed’ (positions of defects are in thermodynamical equilibrium with the system) or ‘quenched’ (positions of defects are out of equilibrium). As was pointed out in [71], the correctness of results, obtained in the picture of ‘quenched’ disorder, depends on whether the location of the starting point of a SAW is fixed while the configurational averaging is performed, or not. In the latter case, one has to average over all locations and effectively this corresponds to the case of annealed disorder.

An interesting question arises: what is the difference in statistics between SAWs walking on percolation clusters and the backbone of percolation clusters, after eliminating all the ‘dead ends’? First Kim [38] claimed, based on a scaling argument, that the critical behavior on the percolation cluster is the same as that on the backbone, namely $\nu_{SAW}^{pc} = \nu_{SAW}^B$. This equality was also assumed by Rammal [50] in deriving a Flory formula for SAWs on fractal substrates. This can be easily explained: since the walks, which visit the dead ends are eventually terminated after a certain number of steps, the walks that survive in the limit $N \rightarrow \infty$ are those confined within the backbone. However, in a numerical study [39] it was found that $\nu_{SAW}^B > \nu_{SAW}^{pc}$ and, moreover, that ν_{SAW}^{pc} almost equals the value for SAWs on pure lattices. It was argued that the averaged end-to-end distance of SAWs on the backbone is significantly enhanced in comparison with averaging on the full percolation cluster. We have checked this, comparing results obtained by us for the averaged end-to-end distance $\langle r(N) \rangle$ on percolation clusters and the backbone of percolation clusters. We conclude that there is practically no difference in scaling for SAWs on both types of clusters, the SAW statistics is determined by the backbone of percolation clusters.

Table 2. The exponent ν_{SAW}^{pc} for a SAW on a percolation cluster. FL: Flory-like theories, EE: exact enumerations, RS, RG: real-space and field-theoretic renormalization group, MC: Monte Carlo simulations. For SAWs on the regular lattice one has $\nu_{\text{SAW}}(d = 2) = 3/4$ [4], $\nu_{\text{SAW}}(d = 3) = 0.5882(11)$ [6], $\nu_{\text{SAW}}(d \geq 4) = 1/2$.

$\nu_{\text{SAW}}^{pc} \setminus d$	2	3	4
FL equation (7)	7/9	0.665	0.594
[49]	0.778	0.662	0.593
[52]	0.77	0.66	0.62
[53]	0.770	0.656	0.57
[54]	0.76	0.65	0.58
EE	[43] 0.745(10)	0.635(10)	
[45]	0.770(5)	0.660(5)	
[46]	0.778(15)	0.66(1)	
RS	[49] 0.778	0.724	
[55]	0.767		
RG	[56] 0.785	0.678	0.595
[57]	0.796	0.669	0.587
MC	[39] 0.77(1)		
[40]	0.783(3)		
[41]		0.62–0.63	0.56–0.57
[46]	0.787(10)	0.662(6)	
Our results	0.782 ± 0.003	0.667 ± 0.003	0.586 ± 0.003

To study the scaling properties of SAWs on the backbone of percolation clusters, we choose as the starting point the ‘seed’ of the cluster, and apply the PERM algorithm, taking into account that a SAW can have its steps only on the sites belonging to the backbone of the percolation cluster. We used lattices of size up to $L_{\text{max}} = 400, 200, 50$ in $d = 2, 3, 4$, respectively, and performed averages over 1000 percolation clusters in each case. Estimates for the critical exponents ν_{SAW}^{pc} were obtained by linear least-square fitting (see tables 6–8 in the appendix). Note that since we can construct lattices only of a finite size L , it is not possible to perform very long SAWs on it. For each L , the scaling for $\langle r(N) \rangle$ holds only up to some ‘marginal’ number of SAWs steps $N_{\text{marg}} \sim L^{1/\nu_{\text{SAW}}^{pc}}$ [58]. We take this into account when analyzing the data obtained; for each lattice size we are interested only in values of $N < N_{\text{marg}}$, thus avoiding distortions, caused by finite-size effects. Our results for the scaling exponent ν_{SAW}^{pc} for SAWs on the backbone of percolation clusters [58] are given in table 2 and compared with previous estimates obtained by a variety of different techniques. We see that the value of ν_{SAW}^{pc} is larger than the corresponding exponent on the pure lattice; the presence of disorder leads to stretching of the trajectory of self-avoiding walks.

A simple modified Flory formula for the exponent of a SAW on a percolation cluster, proposed a long time ago by Kremer [36],

$$\nu_{\text{SAW}}^{pc} = 3 / (d_{pc}^F + 2), \tag{7}$$

gives numbers in an astonishingly good agreement with our numerical data (see table 2). For the estimates we have used the values of the fractal dimension of percolation clusters from table 1. Since $d_{\text{up}} = 6$ for percolation and $d_{pc}^F(d \geq 6) = 4$ [19], we receive from equation (7) that the exponent $\nu_{\text{SAW}}^{pc}(d \geq 6) = 1/2$. Note that there exists a whole family of more sophisticated Flory-like theories [38, 49–54].




Figure 6. Disorder averaged distribution function $r\overline{P}(r, N)$ versus the scaling variable $r/N^{v_{SAW}^{pc}}$ in $d = 2$ dimensions. Lattice size $L = 400$, number of SAW steps $N = 140$ (squares), $N = 120$ (triangles), $N = 100$ (crosses).

Table 3. The connectivity constant z^{pc} for a SAW on a percolation cluster. SS: series studies, EE: exact enumerations, MC: Monte Carlo simulations. For SAWs on the regular lattice one has $z(d = 2) = 2.6385 \pm 0.0001$ [73], $z(d = 3) = 4.68404 \pm 0.00009$ [10], $z(d = 4) = 6.77507 \pm 0.00001$ [9].

$z^{pc} \setminus d$	2	3	4
SS [48]	1.31 ± 0.03		
EE [43]	1.53 ± 0.05		
	1.565 ± 0.005	1.462 ± 0.005	
MC [39]	1.459 ± 0.003		
	1.456 ± 0.005	1.317 ± 0.005	
Our results	1.566 ± 0.005	1.459 ± 0.005	1.340 ± 0.005
$z \cdot p_c$	1.564	1.460	1.333

The disorder averaged distribution function, defined via

$$\overline{P}(r, N) = \frac{1}{M} \sum_{i=1}^M P_i(r, N) \tag{8}$$

can be written in terms of the scaled variables $r/\langle r \rangle$ as

$$r\overline{P}(r, N) \sim f(r/\langle r \rangle) \sim f(r/N^{v_{SAW}^{pc}}). \tag{9}$$

The distribution function is normalized according to $\sum_r \overline{P}(r, N) = 1$, such that: $\langle r \rangle = \sum_r r\overline{P}(r, N)$. The numerical results for the distribution function in $d = 2, 3$ and 4 are shown in figures 6–8 for different N . When plotted against the scaling variable $r/N^{v_{SAW}^{pc}}$, the data are indeed found to nicely collapse onto a single curve, using our values for the exponent v_{SAW}^{pc} reported in table 2.

Let us now turn our attention to estimating the number of different possible SAW configurations \overline{C}_N , defined by equation (2). First, let us note, that the fugacity or connectivity constant z is obviously affected by introducing disorder into the lattice. For the case, when the SAW is not confined only to the percolation cluster, namely when averaging over all the clusters is performed, then $z^{pc} = zp_c$ exactly. In table 3 we present results of this estimate,




Figure 7. Disorder averaged distribution function $r\overline{P}(r, N)$ versus the scaling variable $r/N^{\nu_{SAW}^{pc}}$ in $d = 3$ dimensions. Lattice size $L = 200$, number of SAW steps $N = 80$ (squares), $N = 60$ (triangles), $N = 50$ (crosses).




Figure 8. Disorder averaged distribution function $r\overline{P}(r, N)$ versus the scaling variable $r/N^{\nu_{SAW}^{pc}}$ in $d = 4$ dimensions. Lattice size $L = 50$, number of SAW steps $N = 30$ (squares), $N = 20$ (triangles), $N = 15$ (crosses).

Table 4. The exponent γ_{SAW}^{pc} for a SAW on a percolation cluster. FL: Flory-like theories, EE: exact enumerations, MC: Monte Carlo simulations. For a SAW on the regular lattice one has $\gamma_{SAW}(d = 2) = 43/32$ [75], $\gamma_{SAW}(d = 3) = 1.1596 \pm 0.0002$ [6], $\gamma_{SAW}(d \geq 4) = 1$.

$\gamma_{SAW}^{pc} \setminus d$	2	3	4
FL [53]	1.384	1.379	1.27
EE [43]	1.3 ± 0.1		
	1.34 ± 0.05	1.29 ± 0.05	
MC [37]	1.31 ± 0.03	1.40 ± 0.02	
	1.26 ± 0.05	1.19 ± 0.05	
Our results	1.350 ± 0.008	1.269 ± 0.008	1.250 ± 0.008




Figure 9. Averaged connectivity constant for SAWs on the backbone of percolation clusters in $d = 2$ (triangles), $d = 3$ (squares), $d = 4$ (stars).




Figure 10. Disorder averaged number of SAWs configurations versus number of steps for SAWs on the backbone of percolation clusters in $d = 2$ (triangles), $d = 3$ (squares), $d = 4$ (stars).

taking values for p_c from table 1. However, since each existing bond on the infinite percolation cluster has a non-trivial (correlated) probability of occurrence, a similar argument cannot be applied to the SAWs confined to the infinite percolation cluster only. However, enumeration estimates [72] suggested $z^{p_c} \simeq zp_c$ for SAWs on the percolation cluster. It turns out that any difference from the linear dependence for incipient clusters is subtle and could hardly be detected. We have estimated z_{p_c} as the averaged number of possibilities to perform the next step in the PERM procedure for SAWs on the backbone of percolation clusters (see figure 9); results are presented in table 3.

In the analytical study of [74], it was argued that the exponent γ , governing the scaling of the number of SAW configurations, is not changed by the presence of disorder even at $p = p_c$. This was supported by an exact enumeration study [43]. However, this argument disagrees with results of studies [37, 46, 53], where averaging over single percolation clusters was performed and different values for $\gamma_{SAW}^{p_c}$ were found. In [53] it was argued, using scaling arguments, that at $p = p_c$ the exponents γ_{SAW} will crossover to $\gamma_{SAW}^{p_c} = \gamma_{SAW} + d(\nu_{SAW}^{p_c} - \nu_{SAW})$ at $p = p_c$; the estimates based on this equality are given in the first row of table 4.




Figure 11. Averaged end-to-end distance versus number of steps in a double logarithmic scale of RWs on percolation clusters (crosses) and the backbone of percolation clusters (pluses) in $d = 3$.

We obtained numerical estimates for γ_{SAW}^p , studying the behavior of the quantity

$$\ln \overline{C_N}/N = \ln(A)/N + \ln(z^{p_c}) + (\gamma_{SAW}^{p_c} - 1) \frac{\ln N}{N}, \quad (10)$$

where A is a constant. Figure 10 shows these values for the backbone of percolation clusters in $d = 2, 3, 4$. Estimates for $\gamma_{SAW}^{p_c}$ are obtained by linear least-square fits (see tables 9, 10 and 11 in the appendix). Our final results are presented in table 4.

4. Random walks on percolation clusters

To simulate the diffusion process in a disordered medium, the picture of the ‘ant in the labyrinth’, proposed by de Gennes [76] is traditionally used. Here the walker (an ‘ant’) starts at an arbitrary point on the diluted lattice and tries to move randomly to the nearest site. If the randomly chosen direction leads to an empty site, it moves and the steps increment by 1. If the chosen site is occupied by a defect (in our case, does not belong to the percolation cluster) it stays at the current position for this time step. The growth is stopped, if the total number of steps N is performed, than the next trajectory is started to grow.

After averaging the end-to-end distance over RW configurations on a single percolation cluster, the disorder average is carried out as in equation (5) over all constructed percolation clusters. Let us note that, in contrast to the SAW problem discussed above, the scaling behavior of RWs on a percolation cluster is different from that on its backbone. Let us recall that the statistics of long SAWs on percolation clusters is nevertheless determined by its backbone, since the walks, which visit the ‘dead ends’ are eventually terminated after a certain number of steps. Simple random walks cannot be trapped in ‘dead ends’, however, visiting these parts of a cluster will lead to some ‘slowing down’ of the diffusion process in comparison with the behavior on the backbone where all the dead ends are removed. This is really confirmed by analyzing our results for the averaged end-to-end distance of random walks on a percolation cluster and its backbone (see figure 11).

We have studied RWs both on the percolation cluster and its backbone, performing 10^7 realizations on each cluster and average over 1000 clusters in each space dimension $d = 2, 3, 4$. Estimates of scaling exponents $\nu_{RW}^{p_c}$ and ν_{RW}^B , describing scaling of walks on percolation cluster and backbone, respectively, are obtained by linear least-square fitting and given in table 5.




Figure 12. Disorder averaged distribution function $r\overline{P}(r, N)$ versus the scaling variable $r/\langle r \rangle$ in $d = 2$ dimensions, left: percolation cluster, right: backbone of percolation cluster. Lattice size $L = 400$, number of RW steps $N = 180$ (squares), $N = 160$ (triangles), $N = 140$ (crosses).




Figure 13. Disorder averaged distribution function $r\overline{P}(r, N)$ versus the scaling variable $r/\langle r \rangle$ in $d = 3$ dimensions, left: percolation cluster, right: backbone of percolation cluster. Lattice size $L = 200$, number of RW steps $N = 100$ (squares), $N = 90$ (triangles), $N = 80$ (crosses).

One can see that the inequality $\nu_{RW}^{pc} < \nu_{RW}^B$ holds, and that the quantitative difference between these two values grows with increasing the space dimension d . On the other hand, both values are *smaller* than the corresponding exponent $\nu_{RW} = 1/2$, governing scaling of random walks on the pure lattice: the presence of disorder slows down the diffusion process. The reason for this subdiffusive behavior is intuitively clear: due to the presence of defects, the randomly walking particle returns back to already visited sites more often, thus its walking distance is shorter than on the pure lattice. This has also been observed in recent studies of less-disordered deterministic fractals such as two-dimensional Sierpinski carpet composites [77].

The disorder averaged distribution function, defined in equation (8) and rewritten in terms of the scaled variables $r/\langle r \rangle$ as

$$r\overline{P}(r, N) \sim f(r/\langle r \rangle) \sim f(r/N^{\nu^{pc}}), \tag{11}$$

is shown in figures 12–14 for $d = 2, 3, 4$, both for the cases of the percolation cluster and for the backbone. When plotted against the scaling variable $r/N^{\nu^{pc}}$, the data are indeed found to




Figure 14. Disorder averaged distribution function $r\overline{P}(r, N)$ versus the scaling variable $r/\overline{\langle r \rangle}$ in $d = 4$ dimensions, left: percolation cluster, right: backbone of percolation cluster. Lattice size $L = 50$, number of RW steps $N = 60$ (squares), $N = 50$ (triangles), $N = 40$ (crosses).

Table 5. Exponents governing the scaling law of the end-to-end distance for RWs on percolation clusters and the backbone of percolation clusters. RS: real-space renormalization-group, EE: exact enumerations, MC: Monte Carlo simulations.

	2	3	4
$\nu_{RW}^{pc} \setminus d$			
RS [20]	0.356	0.285	
EE [21]	0.349 ± 0.002		
[22]		0.266 ± 0.01	
Analytic [23]	0.352	0.268	
MC [24]	0.352 ± 0.006	0.271 ± 0.004	
[25]	0.352 ± 0.006	0.271 ± 0.004	
[26]	0.392 ± 0.007	0.282 ± 0.003	
[27]	0.348 ± 0.009	0.274 ± 0.007	
[28]			0.222 ± 0.007
Our results	0.353 ± 0.003	0.273 ± 0.003	0.231 ± 0.003
$\nu_{RW}^B \setminus d$			
Analytic [29]	0.371 ± 0.001		
EE [30]	0.372 ± 0.005		
MC [31]	0.370 ± 0.003		
Our results	0.372 ± 0.002	0.306 ± 0.002	0.262 ± 0.002

Table 6. Results of linear fitting of obtained results for $\overline{\langle r \rangle}$ for SAWs in $d = 2$ dimensions on the backbone of percolation clusters, $L = 400$.

N_{min}	ν_{SAW}^{pc}	A	χ^2/DF
11	0.790 ± 0.005	0.829 ± 0.003	2.396
16	0.786 ± 0.005	0.841 ± 0.005	1.910
21	0.782 ± 0.004	0.847 ± 0.005	1.479
26	0.783 ± 0.003	0.842 ± 0.006	1.262
31	0.782 ± 0.003	0.840 ± 0.007	0.839

Table 7. Same as table 6 for $d = 3, L = 200$.

N_{\min}	ν_{SAW}^{pc}	A	χ^2/DF
11	0.668 ± 0.003	0.935 ± 0.004	2.269
16	0.669 ± 0.003	0.930 ± 0.004	2.054
21	0.669 ± 0.003	0.924 ± 0.004	1.345
26	0.667 ± 0.002	0.930 ± 0.006	0.743
31	0.668 ± 0.002	0.934 ± 0.008	0.844

Table 8. Same as table 6 for $d = 4, L = 50$.

N_{\min}	ν_{SAW}^{pc}	A	χ^2/DF
8	0.588 ± 0.002	1.025 ± 0.004	2.615
10	0.587 ± 0.002	1.023 ± 0.006	1.777
12	0.586 ± 0.003	1.021 ± 0.01	0.978
14	0.586 ± 0.003	1.031 ± 0.01	0.767

Table 9. Results of linear fitting of obtained results for $\overline{C_N}$ for SAWs in $d = 2$ dimensions on the backbone of percolation clusters, $L = 400$.

N_{\min}	γ_{SAW}^{pc}	A	χ^2/DF
16	1.341 ± 0.005	1.219 ± 0.003	3.135
21	1.349 ± 0.005	1.189 ± 0.003	2.682
26	1.351 ± 0.007	1.168 ± 0.002	1.913
31	1.352 ± 0.008	1.172 ± 0.002	1.621
36	1.350 ± 0.008	1.163 ± 0.001	0.704

Table 10. Same as table 9 for $d = 3, L = 200$.

N_{\min}	γ_{SAW}^{pc}	A	χ^2/DF
11	1.265 ± 0.004	1.82 ± 0.003	2.767
16	1.268 ± 0.005	1.192 ± 0.003	2.135
21	1.270 ± 0.006	1.184 ± 0.003	1.968
26	1.267 ± 0.008	1.176 ± 0.002	1.513
31	1.269 ± 0.008	1.172 ± 0.002	0.762

Table 11. Same as table 9 for $d = 4, L = 50$.

N_{\min}	γ_{SAW}^{pc}	A	χ^2/DF
8	1.251 ± 0.005	1.77 ± 0.003	1.767
10	1.252 ± 0.007	1.182 ± 0.003	1.135
12	1.250 ± 0.008	1.184 ± 0.003	0.968

nicely collapse onto a single curve, using our values for the exponent $\nu_{\text{RW}}^{pc}, \nu_{\text{RW}}^B$ reported in table 5.

5. Conclusions

We studied the scaling behavior of simple random walks and self-avoiding walks on disordered lattices. Both models are of great interest: RWs provide a good description of diffusion

processes, SAWs are successful in describing the universal properties of long flexible polymer macromolecules in a good solvent.

We consider the case, when the concentration p of lattice sites allowed for walking equals the critical concentration p_c and an incipient percolation cluster can be found in the system. Studying properties of percolative lattices, one encounters two possible statistical averages: in the first, one considers only percolation clusters with linear size much larger than the typical length of the physical phenomenon under discussion, the other statistical ensemble includes all the clusters, which can be found on a percolative lattice. In our study, we considered only the first case, being interested in random and self-avoiding walks on a percolation cluster, which has a fractal structure. In this regime, the critical behavior of both models belongs to a new universality class: scaling law exponents change with the dimension of the (fractal) lattice on which the walk resides.

We performed numerical simulations of random and self-avoiding walks on percolation clusters and the backbone of percolation clusters on lattice sizes $L = 400, 200, 50$ in space dimensions $d = 2, 3, 4$, respectively. Our results bring about the estimates for critical exponents, governing the scaling behavior of the models. We found that the statistics of SAWs is governed by the same scaling exponent both on a percolation cluster and its backbone: since the walks, which visit the dead ends are eventually terminated after a certain number of steps, the walks that survive in the limit $N \rightarrow \infty$ on a percolation cluster are those confined within its backbone. For simple random walks, however, the picture is different: they cannot be trapped in ‘dead ends’. However, visiting these parts of a cluster will lead to some ‘slowing down’ of the diffusion process in comparison with the behavior on the backbone where all the dead ends are removed. We found that the inequality $v_{\text{RW}}^{p_c} < v_{\text{RW}}^B$ holds and the quantitative difference between these two values grows with increasing space dimension d .

The presence of disorder leads to a stretching of the trajectory of self-avoiding walks: the value of $v_{\text{SAW}}^{p_c}$ is larger than the corresponding exponent on the pure lattice. However, the exponent $v_{\text{RW}}^{p_c}$ governing scaling of random walks on percolative lattices is smaller than that on a pure lattice: the presence of disorder slows down the diffusion process. This can be explained as follows: due to the presence of defects, the randomly walking particle returns back to already visited sites more often, thus its walking distance is shorter than on the pure lattice.

Acknowledgments

Work supported in part by the German Science Foundation (DFG) through the Research Group FOR877 and by the EU RTN-Network ‘ENRAGE’: *Random Geometry and Random Matrices: From Quantum Gravity to Econophysics* under grant no. MRTN-CT-2004-005616. VB is grateful for support through the ‘Marie Curie International Incoming Fellowship’ EU Programme and to the Institut für Theoretische Physik, Universität Leipzig, for hospitality.

Appendix

To estimate the critical exponents $v_{\text{SAW}}^{p_c}, \gamma_{\text{SAW}}^{p_c}$, linear least-square fits with varying lower cutoff for the number of steps N_{min} are used in order to detect possible corrections to scaling. For estimating $v_{\text{SAW}}^{p_c}$ we use linear fits for the averaged end-to-end distance $\ln(\overline{r(N)}) = \ln(A) + v_{\text{SAW}}^{p_c} \ln N$, and for $\gamma_{\text{SAW}}^{p_c}$ we employ equation (10). Since this is an important aspect for assessing the quality of our final exponent estimates discussed in the main text, we have compiled in this appendix these more detailed results in tables 6–11.

The χ^2 value (sum of squares of normalized deviation from the regression line) divided by the number of degrees of freedom, DF, given in the last rows, serves as a test of the goodness of fit.

References

- [1] See e.g. Shlesinger M F and West B (ed) 1984 *Random Walks and their Applications in the Physical and Biological Sciences* (AIP Conf. Proc. vol 109) (New York: AIP)
- Spitzer F 1976 *Principles of Random Walk* (Berlin: Springer)
- [2] des Cloizeaux J and Jannink G 1990 *Polymers in Solution* (Oxford: Clarendon)
- de Gennes P-G 1979 *Scaling Concepts in Polymer Physics* (Ithaca, NY: Cornell University Press)
- [3] Le Guillou J C and Zinn-Justin J 1980 *Phys. Rev. B* **21** 3976
- [4] Nienhuis B 1982 *Phys. Rev. Lett.* **49** 1062
- [5] Le Guillou J C and Zinn-Justin J 1985 *J. Physique Lett.* **46** L127
- Le Guillou J C and Zinn-Justin J 1989 *J. Physique* **50** 1365
- [6] Guida R and Zinn-Justin J 1998 *J. Phys. A: Math. Gen.* **31** 8104
- [7] Rosenbluth M N and Rosenbluth A W 1955 *J. Chem. Phys.* **23** 356
- [8] Madras N and Sokal A D 1988 *J. Stat. Phys.* **50** 109
- [9] MacDonald D, Hunter D L, Kelly K and Jan N 1992 *J. Phys. A: Math. Gen.* **25** 1429
- [10] MacDonald D, Joseph S, Hunter D L, Moseley L L, Jan N and Guttmann A J 2000 *J. Phys. A: Math. Gen.* **33** 5973
- [11] Li B, Madras N and Sokal A D 1995 *J. Stat. Phys.* **80** 661
- [12] Caracciolo S, Causo M S and Pelissetto A 1998 *Phys. Rev. E* **57** R1215
- [13] Havlin S and Abraham D Ben 1987 *Phys. Adv.* **36** 695
- [14] Ziff R M 1994 *Phys. Rev. Lett.* **72** 1942
- [15] Grassberger P 1992 *J. Phys. A: Math. Gen.* **25** 5867
- [16] Paul G, Ziff R M and Stanley H E 2001 *Phys. Rev. E* **64** 026115
- [17] Grassberger P 1986 *J. Phys. A: Math. Gen.* **19** 1681
- [18] Moukarzel C 1998 *Int. J. Mod. Phys. C* **8** 887
- [19] Stauffer D and Aharony A 1992 *Introduction to Percolation Theory* (London: Taylor and Francis)
- [20] Sahimi M and Jerauld J 1983 *J. Phys. C: Solid State Phys.* **16** L1043
- [21] Majid I, Ben-Avraham D, Havlin S and Stanley H E 1984 *Phys. Rev. B* **30** 1626
- [22] Pandey R B and Stauffer D 1983 *Phys. Rev. Lett.* **51** 527
- [23] Alexander S and Orbach R 1982 *J. Phys. Paris Lett.* **43** L635
- [24] Ben-Avraham D and Havlin S 1982 *J. Phys. A: Math. Gen.* **15** L691
- Havlin S, Ben-Avraham D and Sompolinsky D 1982 *Phys. Rev. A* **27** 1730
- [25] Havlin S and Ben-Avraham D 1983 *J. Phys. A: Math. Gen.* **16** L483
- [26] Argyrakis P and Kopelman P 1984 *Phys. Rev. B* **29** 511
- [27] McCarthy J F 1988 *J. Phys. A: Math. Gen.* **21** 3379
- [28] Lee S B and Nakanishi H 2000 *J. Phys. A: Math. Gen.* **33** 2943
- [29] Bug A L, Grest G S, Cohen M H and Webman I 1986 *J. Phys. A: Math. Gen.* **19** L323
- [30] Hong D, Havlin S and Herrmann H 1984 *Phys. Rev. B* **7** 4083
- [31] Mastorakos J and Argyrakis P 1993 *Phys. Rev. E* **48** 4847
- [32] Webman I 1981 *Phys. Rev. Lett.* **47** 1497
- [33] Gefen Y, Aharony A and Alexander S 1983 *Phys. Rev. Lett.* **50** 77
- [34] Rammal R and Toulouse G 1983 *J. Phys. Paris Lett.* **44** L13
- [35] Mukherjee S, Jacobs D and Nakanishi H 1995 *J. Phys. A: Math. Gen.* **28** 291
- [36] Kremer K 1981 *Z. Phys. B* **45** 149
- [37] Lee S B and Nakanishi H 1988 *Phys. Rev. Lett.* **61** 2022
- Lee S B, Nakanishi H and Kim Y 1989 *Phys. Rev. B* **39** 9561
- [38] Kim Y 1990 *Phys. Rev. A* **41** 4554
- [39] Woo K Y and Lee S B 1991 *Phys. Rev. A* **44** 999
- [40] Grassberger P 1993 *J. Phys. A: Math. Gen.* **26** 1023
- [41] Lee S B 1996 *J. Korean Phys. Soc.* **29** 1
- [42] Meir Y and Harris A B 1989 *Phys. Rev. Lett.* **63** 2819
- [43] Lam P M 1990 *J. Phys. A: Math. Gen.* **23** L831
- [44] Nakanishi H and Moon J 1992 *Physica A* **191** 309
- [45] Rintoul M D, Moon J and Nakanishi H 1994 *Phys. Rev. E* **49** 2790

- [46] Ordemann A, Porto M, Roman H E, Havlin S and Bunde A 2000 *Phys. Rev. E* **61** 6858
- [47] Nakanishi H and Lee S B 1991 *J. Phys. A: Math. Gen.* **24** 1355
- [48] Barat K, Karmakar S N and Chakrabarti B K 1991 *J. Phys. A: Math. Gen.* **24** 851
- [49] Sahimi M 1984 *J. Phys. A: Math. Gen.* **17** L379
- [50] Rammal R, Toulouse G and Vannimenus J 1984 *J. Phys. Paris* **45** 389
- [51] Kim Y 1987 *J. Phys. A: Math. Gen.* **20** 1293
- [52] Roy A K and Blumen A 1990 *J. Stat. Phys.* **59** 1581
- [53] Roy A K and Chakrabarti B K 1987 *J. Phys. A: Math. Gen.* **20** 215
- [54] Aharony A and Harris A B 1989 *J. Stat. Phys.* **54** 1091
- [55] Lam P M and Zhang Z Q 1984 *Z. Phys. B* **56** 155
- [56] von Ferber C, Blavatska V, Folk R and Yu H 2004 *Phys. Rev. E* **70** 035104
- [57] Janssen H-K and Stenull O 2007 *Phys. Rev. E* **75** 020801
- [58] Blavatska V and Janke W 2008 *Europhys. Lett.* **82** 66006
- [59] Hoshen J and Kopelman R 1976 *Phys. Rev. B* **14** 3438
- [60] Aizenmann M 1997 *Nucl. Phys. B* **485** 551
Shchur L N and Rostunov T 2002 *JETP Lett.* **76** 475
- [61] Herrmann H J, Hong D C and Stanley H E 1984 *J. Phys. A: Math. Gen.* **17** L261
- [62] Porto M, Bunde A, Havlin S and Roman H E 1997 *Phys. Rev. E* **56** 1667
- [63] Havlin S, Nossal R, Trus B and Weiss G H 1984 *J. Phys. A: Math. Gen.* **17** L957
- [64] Grassberger P 1997 *Phys. Rev. E* **56** 3682
- [65] Vogel T, Bachmann M and Janke W 2007 *Phys. Rev. E* **76** 061803
- [66] Wall F T and Erpenbeck J J 1959 *J. Chem. Phys.* **30** 634
- [67] Hsu H P, Mehra V, Nadler W and Grassberger P 2007 *J. Chem. Phys.* **118** 444
- [68] Bachmann M and Janke W 2003 *Phys. Rev. Lett.* **91** 208105
Bachmann M and Janke W 2004 *J. Chem. Phys.* **120** 6779
- [69] Mehra V and Grassberger P 2002 *Physica D* **168** 244
- [70] Frauenkron H, Bastolla U, Gerstner E, Grassberger P and Nadler W 1998 *Phys. Rev. Lett.* **80** 3149
- [71] Le Doussal P and Machta J 1991 *J. Stat. Phys.* **64** 541
- [72] Chakrabarti B K, Bhadra K, Roy A K and Karmakar S N 1983 *Phys. Lett. A* **93** 434
- [73] Guttmann A J and Wang J 1991 *J. Phys. A: Math. Gen.* **24** 3107
- [74] Lyklema J W and Kremer K 1984 *Z. Phys. B* **55** 41
- [75] Nienhuis B 1984 *J. Stat. Phys.* **34** 731
- [76] de Gennes P G 1976 *La Recherche* **7** 919
- [77] Anh D H N, Hoffmann K H, Seeger S and Tarafdar S 2005 *Europhys. Lett.* **70** 109
Anh D H N, Blaudeck P, Hoffmann K H, Prehl J and Tarafdar S 2007 *J. Phys. A: Math. Theor.* **40** 11453