

Maximally entangled mixed states of two atoms trapped inside an optical cavity

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2009 J. Phys. A: Math. Theor. 42 245302

(<http://iopscience.iop.org/1751-8121/42/24/245302>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.154

The article was downloaded on 03/06/2010 at 07:53

Please note that [terms and conditions apply](#).

Maximally entangled mixed states of two atoms trapped inside an optical cavity

Shang-Bin Li^{1,2} and Jing-Bo Xu¹

¹ Zhejiang Institute of Modern Physics and Department of Physics, Zhejiang University, Hangzhou 310027, People's Republic of China

² Shanghai research center of Amertron-global, Zhangjiang High-Tech Park, 299 Lane, Bisheng Road, No. 3, Suite 202, Shanghai, 201204, People's Republic of China

E-mail: stephenli74@yahoo.com.cn

Received 22 September 2008, in final form 17 February 2009

Published 26 May 2009

Online at stacks.iop.org/JPhysA/42/245302

Abstract

In some off-resonant cases, the reduced density matrix of two atoms symmetrically coupled with an optical cavity can very approximately approach maximally entangled mixed states or maximal Bell violation mixed states in their evolution. The influence of a phase decoherence on the generation of a maximally entangled mixed state is also discussed.

PACS numbers: 03.67.-a, 03.65.Ud

Quantum entanglement plays a crucial role in quantum information processes [1]. In the past few years, much attention has been paid to the preparation of maximally entangled mixed states [2–4]. The properties of maximally entangled mixed states have been studied by many authors [5–7]. Maximally entangled mixed states are those states that, for a given mixedness, achieve the greatest possible entanglement. For two-qubit systems and for various combinations of entanglement and mixedness measures, the forms of the corresponding maximally entangled mixed states are different [7]. Using correlated photons from parametric down-conversion, maximally entangled mixed states in the linear entropy–concurrence plane have been created and characterized [2]. Generation and characterization of two-photon polarization maximally entangled mixed states in the linear entropy–concurrence plane have also been carried out, which is based on the peculiar spatial characteristics of a high brilliance source of entangled pairs [3]. The preparation of maximally entangled mixed states of two atoms asymmetrically on-resonance coupled with an optical cavity has also been proposed [4]. Recently, Clark and Parkins [8] have proposed a scheme to controllably entangle the internal states of two atoms trapped in a high-finesse optical cavity by employing quantum-reservoir engineering. By applying the on-resonance atom–cavity couplings which are time dependent, Olaya-Castro *et al* have also presented an efficient scheme for controlled generation of entangled states of two atoms inside an optical cavity [9]. However, truly resonant coupling is not available in a

realistic physical system. It is desirable to investigate how the off-resonance coupling affects the preparation of maximally entangled mixed states of two atoms. Based on our previous analytical results in [10], in which the entanglement behaviors of two atoms inside an optical cavity in the presence of a phase decoherence have been derived, we can easily analyze the feasibility for preparing maximally entangled mixed states in such a system. For keeping this paper self-contained, we briefly outline the basics about two two-level atoms inside an optical cavity. Here, we investigate two two-level atoms symmetrically coupling to a single-mode optical cavity and show that in some off-resonant cases, the maximally entangled mixed states in the plane of concurrence versus linear entropy of two atoms can be very approximately generated. It is shown that the long-time entanglement behavior of two atoms is sensitive to the ratio of the detuning and the coupling strength. The influence of the initial mixedness of the atoms and phase decoherence is also analyzed.

Considering the system that two atoms are trapped inside a single-mode optical cavity initially prepared in the vacuum state, the Hamiltonian for the system can be given by [11, 12] ($\hbar = 1$),

$$H = \frac{\omega_0}{2} \sum_i^2 \sigma_z^{(i)} + \omega a^\dagger a + g \sum_i^2 (a \sigma_+^{(i)} + a^\dagger \sigma_-^{(i)}), \quad (1)$$

where $\sigma_z^{(i)}, \sigma_\pm^{(i)}$ ($i = 1, 2$) are atomic operators, ω_0 is the atomic transition frequency, g is the coupling constant of an individual atom to the cavity field and a (a^\dagger) is the annihilation (creation) operator of the cavity field with frequency ω . The generation of the entangled state in the system (1) in the laboratory has been implemented [12]. Various modifications and generalizations of the system (1) have been studied for preparing entangled states or realizing various kinds of quantum information processes [13–16]. It is assumed that the cavity fields are prepared initially in the vacuum state $|0\rangle$, and atom 1 is prepared in the mixed state $\lambda|e\rangle\langle e| + (1 - \lambda)|g\rangle\langle g|$ and atom 2 is in the ground state $|g\rangle$, i.e.,

$$\rho(0) = |0\rangle\langle 0| \otimes [\lambda|e\rangle\langle e| + (1 - \lambda)|g\rangle\langle g|] \otimes |g\rangle\langle g|. \quad (2)$$

The time evolution of $\rho(t)$ can be derived as follows:

$$\begin{aligned} \rho(t) = & \frac{\lambda}{8} \left[1 + \frac{\Delta^2}{\Omega^2} + \left(1 - \frac{\Delta^2}{\Omega^2} \right) \cos \Omega t \right] |0\rangle\langle 0| \otimes |B_+\rangle\langle B_+| \\ & + \frac{g^2 \lambda}{\Omega^2} [1 - \cos \Omega t] |1\rangle\langle 1| \otimes |gg\rangle\langle gg| + \frac{\lambda}{4} |0\rangle\langle 0| \otimes |B_-\rangle\langle B_-| \\ & + \frac{\sqrt{2}g\lambda}{2\Omega} \left[\frac{\Delta}{\Omega} (1 - \cos \Omega t) + i \sin \Omega t \right] |0\rangle\langle 1| \otimes |B_+\rangle\langle gg| \\ & + \frac{\sqrt{2}g\lambda}{2\Omega} \left[\exp\left(\frac{i\Omega t - i\Delta t}{2}\right) - \exp\left(\frac{-i\Omega t - i\Delta t}{2}\right) \right] |0\rangle\langle 1| \otimes |B_-\rangle\langle gg| \\ & + \frac{\lambda}{4} \left\{ \left(1 - \frac{\Delta}{\Omega} \right) e^{i(\Omega+\Delta)t/2} + \left(1 + \frac{\Delta}{\Omega} \right) e^{-i(\Omega-\Delta)t/2} \right\} |0\rangle\langle 0| \otimes |B_+\rangle\langle B_-| \\ & + \frac{1 - \lambda}{2} |0\rangle\langle 0| \otimes |gg\rangle\langle gg| + \text{h.c.}, \end{aligned} \quad (3)$$

where $\Delta = \omega_0 - \omega$ is the detuning between the atoms and cavity field, $\Omega = (\Delta^2 + 8g^2)^{1/2}$, and $|B_\pm\rangle = \frac{\sqrt{2}}{2}(|eg\rangle \pm |ge\rangle)$ are the Bell states. By tracing out the degree of freedom of the cavity field, we obtain the reduced density matrix $\rho_s(t)$ describing the subsystem containing

only two atoms,

$$\begin{aligned} \rho_s(t) = & \frac{\lambda}{8} \left[1 + \frac{\Delta^2}{\Omega^2} + \left(1 - \frac{\Delta^2}{\Omega^2} \right) \cos \Omega t \right] |B_+\rangle\langle B_+| \\ & + \frac{\lambda}{4} |B_-\rangle\langle B_-| + \left\{ \frac{g^2 \lambda}{\Omega^2} [1 - \cos \Omega t] + \frac{1 - \lambda}{2} \right\} |gg\rangle\langle gg| \\ & + \frac{\lambda}{4} \left\{ \left(1 - \frac{\Delta}{\Omega} \right) e^{i(\Omega+\Delta)t/2} + \left(1 + \frac{\Delta}{\Omega} \right) e^{-i(\Omega-\Delta)t/2} \right\} |B_+\rangle\langle B_-| + \text{h.c.} \end{aligned} \quad (4)$$

First, we analyze the feasibility of preparing maximally entangled mixed states of two atoms in this cavity QED system. The concurrence [17] is adopted to quantify the bipartite entanglement between the two atoms, and the linear entropy defined by $M = \frac{4}{3}(1 - \text{Tr} \rho_s^2)$ of the reduced density matrix is used to quantify the mixedness. In the situation with a rational value of $\frac{\Delta}{\Omega}$, the evolving density matrix is periodic. In the case with an irrational value of $\frac{\Delta}{\Omega}$, the evolving state is not periodic. The explicit analytical expression of the concurrence $C_s(t)$ characterizing the entanglement in $\rho_s(t)$ can be obtained as

$$\begin{aligned} C_s(t) &= \lambda(A^2 + B^2)^{\frac{1}{2}}, \\ A &= \frac{\Delta^2}{4\Omega^2} - \frac{1}{4} + \frac{1}{4} \left(1 - \frac{\Delta^2}{\Omega^2} \right) \cos \Omega t, \\ B &= \frac{1}{2} \left(1 - \frac{\Delta}{\Omega} \right) \sin(\Omega + \Delta)t/2 - \frac{1}{2} \left(1 + \frac{\Delta}{\Omega} \right) \sin(\Omega - \Delta)t/2. \end{aligned} \quad (5)$$

In the case with $\lambda = 1$, the two atoms can be in pure states at some specific times. The entanglements characterized by the concurrence of those pure states are given by $C = |\sin \frac{\Delta k \pi}{\Omega}|$ which are achieved at discrete times denoted by $t = 2k\pi/\Omega$ ($k = 1, 2, \dots$). If $\frac{\Delta}{\Omega}$ is a rational number, the series $|\sin \frac{\Delta k \pi}{\Omega}|$ ($k = 1, 2, \dots$) have finite and discrete values. While for the case that $\frac{\Delta}{\Omega}$ is an irrational number, the series $|\sin \frac{\Delta k \pi}{\Omega}|$ ($k = 1, 2, \dots$) have infinite numbers of values, and this series can very approximately approach any values between 0 and 1 according to Hurwitz's theorem in number theory. It means pure two-qubit states with any desired degree of entanglement can be very approximately generated for those cases with the irrational values of $\frac{\Delta}{\Omega}$.

In the large detuning limit, i.e., $g/|\Delta| \ll 1$, the population of the single-mode cavity field will be very small in the time evolution, which leads to very small entanglement between the atoms and the cavity field. Therefore the mixedness of the subsystem containing two atoms is very small. In the small detuning limit, i.e. $|\Delta|/g \ll 1$ but not zero, and simultaneously $\frac{\Delta}{\Omega}$ is an irrational number, the trajectories of the reduced density operator of two atoms in the concurrence versus linear-entropy plane exhibit a kind of 'quasi-ergodic' property, roughly speaking, where 'quasi-ergodic' means there are no distinct interspaces in the pattern formed by the trajectory of the evolving state in the concurrence versus linear-entropy plane.

In figure 1, the concurrence versus mixedness of the two atoms is depicted for different values of detuning. In the resonant case, the concurrence of the two atoms increases (decreases) with the increase (decrease) of mixedness of their reduced density matrix. In the resonant situation, the evolving reduced density matrix $\rho_s(t)$ in equation (4) cannot become any one of the maximally entangled mixed states in the plane of linear entropy concurrence. Interestingly, in the off-resonant case, part of the frontier of the concurrence versus linear entropy can be very approximately reached by the evolving reduced density matrix of the two atoms. However, the region in the frontier which can be approximately approached by the evolving reduced density matrix reduces with the increase of the detuning. Approximately, two atoms can acquire the




Figure 1. The concurrence versus mixedness of two atoms is depicted. The trajectory is chosen from the scaled time $gt \in [0, 50]$. The dashed line and dotted line in (a), (b) and (c) represent the Werner state and the maximally entangled mixed state (the frontier of the concurrence versus linear entropy) respectively. (a) $\Delta = 0$; (b) $\Delta = 0.5g$, $\Delta/\Omega = 1/\sqrt{33}$ (an irrational number); (c) $\Delta = 5g$, $\Delta/\Omega = 5/\sqrt{33}$ (an irrational number). In the three cases, $\lambda = 1$, i.e. the starting point is the origin.

desired pure state concurrence between 1 and 0 for both the small detuning case and large detuning case under the precondition that $\frac{\Delta}{\Omega}$ is an irrational number.

From figure 2, we can understand the influence of initial mixedness of two atoms on the entanglement and mixedness of their evolving reduced density matrix. It is shown that the range of the frontier that can approximately be approached is reduced when the initial mixedness of the atoms increases. For the case with $\lambda = 0.6$, two atoms can evolve into a state with smaller mixedness than their initial state which is different from other cases with $\lambda = 0.9$ and $\lambda = 0.7$. One can also find that the patterns formed by the trajectories are mirror symmetric with the horizontal axis labeled by half of the concurrence of the maximally entangled mixed state corresponding to the initial linear entropy.

Bell's inequality test with entangled atoms inside a cavity has been extensively studied [18]. The most commonly discussed Bell inequality is the CHSH inequality [19, 20]. The CHSH operator reads

$$\hat{B} = \vec{a} \cdot \vec{\sigma} \otimes (\vec{b} + \vec{b}') \cdot \vec{\sigma} + \vec{a}' \cdot \vec{\sigma} \otimes (\vec{b} - \vec{b}') \cdot \vec{\sigma}, \quad (6)$$

where $\vec{a}, \vec{a}', \vec{b}, \vec{b}'$ are unit vectors. In the above notation, the Bell inequality reads

$$|\langle \hat{B} \rangle| \leq 2. \quad (7)$$

The maximal amount of Bell violation of a state ρ is given by [21]

$$|\mathcal{B}|_{\max} = 2\sqrt{\kappa + \tilde{\kappa}}, \quad (8)$$

where κ and $\tilde{\kappa}$ are the two largest eigenvalues of $T_\rho^\dagger T_\rho$. The matrix T_ρ is determined completely by the correlation functions being a 3×3 matrix whose elements are $(T_\rho)_{nm} = \text{Tr}(\rho \sigma_n \otimes \sigma_m)$. Here, $\sigma_1 \equiv \sigma_x, \sigma_2 \equiv \sigma_y$ and $\sigma_3 \equiv \sigma_z$ denote the usual Pauli matrices. The quantity $|\mathcal{B}|_{\max}$ is called the maximal violation measure, which indicates the Bell violation when $|\mathcal{B}|_{\max} > 2$ and the maximal violation when $|\mathcal{B}|_{\max} = 2\sqrt{2}$. For the density operator ρ_s in equation (4), $\kappa + \tilde{\kappa}$ can be written as follows

$$\kappa + \tilde{\kappa} = \zeta + \max[\zeta, \zeta], \quad (9)$$

where

$$\zeta = \frac{4g^4}{\Omega^4} (1 - \cos \Omega t)^2 + \frac{1}{4} \left[\left(1 - \frac{\Delta}{\Omega} \right) \sin \frac{(\Omega + \Delta)t}{2} - \left(1 + \frac{\Delta}{\Omega} \right) \sin \frac{(\Omega - \Delta)t}{2} \right]^2 \quad (10)$$

$$\zeta = \left(\frac{\Delta^2 + 4g^2}{\Omega^2} + \frac{4g^2}{\Omega^2} \cos \Omega t \right)^2$$

in the case with $\lambda = 1$. In [7], the analytical form of the mixed states which possess the maximal value of $|\mathcal{B}|_{\max}$ of two qubits for a given linear entropy has been derived. Here, part of the frontier of the maximal Bell violation versus the linear entropy can also be very approximately approached by the evolving state of the two atoms (see figure 3(b)). In figure 3(a), our calculations show that the two atoms cannot violate the Bell-CHSH inequality in the resonant case, though they could get entangled. While in the off-resonant case, the Bell violation of atom 1 and atom 2 can emerge in their long-time evolution, even though the detuning Δ is very very small.

If the pure phase decoherence mechanism is considered, the master equation governing the time evolution of the system under the Markovian approximation is given by [22, 23]

$$\frac{d\rho}{dt} = -i[H, \rho] - \frac{\gamma}{2}[H, [H, \rho]], \quad (11)$$




Figure 2. The concurrence versus mixedness of two atoms is displayed in the cases in which one of the atoms is initially in three different mixed states: (a) $\lambda = 0.9$; (b) $\lambda = 0.7$; (c) $\lambda = 0.6$. The trajectories are also chosen from the scaled time $gt \in [0, 500]$. The dashed line and dotted line in (a), (b) and (c) represent the Werner state and the maximally entangled mixed state, respectively. It is shown that two atoms can approximately approach part of the maximally entangled mixed states though the range of the frontier that can be approached is reduced when the initial mixedness of the atom increases. In the three cases, $\Delta = 0.5g$, $\Delta/\Omega = 1/\sqrt{33}$ (an irrational number).




Figure 3. The maximal Bell violation $|B|_{\max}$ versus the mixedness of two atoms is displayed for three different values of the detuning: (a) $\Delta = 0$; (b) $\Delta = 0.01g$, $\Delta/\Omega = \frac{1}{3\sqrt{8889}}$ (an irrational number); (c) $\Delta = 5g$, $\Delta/\Omega = 5/\sqrt{33}$ (an irrational number). The trajectory is chosen from the scaled time $gt \in [0, 500]$. The dashed line represents the frontier of maximal Bell violation versus the linear entropy, namely, for a given linear entropy, the maximal value of $|B|_{\max}$ of two atoms can not exceed the dashed line. In the three cases, $\lambda = 1$.

where γ is the phase decoherence rate. The explicit analytical expression of the concurrence $C_\gamma(t)$ characterizing the entanglement of the two atoms in the presence of a phase decoherence




Figure 4. We display the concurrence versus mixedness of two atoms in the presence of phase decoherence. The trajectories are chosen from the scaled time $gt \in [0, 500]$. The dashed line and dotted line in (a), (b) and (c) represent the Werner state and the maximally entangled mixed state, respectively. (a) $\Delta = 0$; (b) $\Delta = 0.5g$, $\Delta/\Omega = 1/\sqrt{33}$ (an irrational number), it can be observed that an appropriate detuning and decoherence rate can make two atoms possess the ability to approach the wider region of the frontier; (c) $\Delta = g$, $\Delta/\Omega = 1/3$ (a rational number). In the three cases, $\gamma = 0.01/g$ and $\lambda = 1$.

can be obtained as

$$\begin{aligned}
C_\gamma(t) &= \lambda(A_\gamma^2 + B_\gamma^2)^{\frac{1}{2}}, \\
A_\gamma &= \frac{\Delta^2}{4\Omega^2} - \frac{1}{4} + \frac{1}{4} \left(1 - \frac{\Delta^2}{\Omega^2}\right) \cos \Omega t \exp\left(-\frac{\gamma t}{2}\Omega^2\right), \\
B_\gamma &= \frac{1}{2} \left(1 - \frac{\Delta}{\Omega}\right) \sin(\Omega + \Delta)t/2 \exp\left[-\frac{\gamma t}{8}(\Omega + \Delta)^2\right] \\
&\quad - \frac{1}{2} \left(1 + \frac{\Delta}{\Omega}\right) \sin(\Omega - \Delta)t/2 \exp\left[-\frac{\gamma t}{8}(\Omega - \Delta)^2\right], \tag{12}
\end{aligned}$$

if the system is initially in the same state as $\rho(0)$ in equation (2). From equation (12), we can easily know that the phase decoherence does not completely destroy the entanglement but generates a stationary entangled state of the two atoms. The concurrence $C_\gamma(t)$ is not greater than 0.5 in the resonant case. The entanglement of the stationary state decreases with the increase of the detuning. The phase decoherence changes trajectories in the plane of concurrence versus linear entropy of the evolving state and makes the trajectories chaotic. In figure 4, we display the concurrence versus mixedness of the two atoms in the presence of phase decoherence. The evolving reduced density matrix of the two atoms can approximately approach a wider region of the maximally entangled mixed states, if both the ratio Δ/g and the decoherence rate γ are appropriate.

In summary, we have investigated a possible scheme for generating the maximally entangled mixed state of two atoms which are symmetrically coupled to a single-mode optical cavity field. It is shown that the two atoms cannot achieve the maximally entangled mixed state in the resonant case. In the off-resonant case, the reduced density matrix of the two atoms can approximately approach the maximally entangled state in their evolution. The distinct roles of the rational values or irrational values of $\frac{\Delta}{\Omega}$ in the long-time behaviors of entanglement and mixedness of the two atoms have been clarified. The influence of the phase decoherence and the initial mixedness of the atoms is also discussed. These results presented here may have potential applications in the domain of quantum information and quantum communication and in the field dealing with the fundamental tests of quantum mechanics.

References

- [1] DiVincenzo D P 1995 *Science* **270** 255
 Cirac J I and Zoller P 2000 *Nature* **404** 579
 Nielsen M A and Chuang I L 2000 *Quantum Computation and Quantum Information* (Cambridge: Cambridge University Press)
- [2] Peters N A *et al* 2004 *Phys. Rev. Lett.* **92** 133601
- [3] Barbieri M *et al* 2004 *Phys. Rev. Lett.* **92** 177901
- [4] Li S-B 2007 *Phys. Rev. A* **75** 054304
- [5] Ishizaka S and Hiroshima T 2000 *Phys. Rev. A* **62** 022310
- [6] Verstraete F, Audenaert K and De Moor B 2001 *Phys. Rev. A* **64** 012316
- [7] Wei T C *et al* 2003 *Phys. Rev. A* **67** 022110
- [8] Clark S G and Parkins A S 2003 *Phys. Rev. Lett.* **90** 047905
- [9] Olaya-Castro A, Johnson N F and Quiroga L 2004 *Phys. Rev. A* **70** 020301
 Olaya-Castro A, Johnson N F and Quiroga L 2004 *J. Opt. B: Quantum Semiclass. Opt.* **6** S730
- [10] S-B Li and Xu J-B 2005 *Phys. Rev. A* **72** 022332
- [11] Zheng S B and Guo G C 2000 *Phys. Rev. Lett.* **85** 2392
- [12] Hagle E, Maître X, Nogues G, Wunderlich C, Brune M, Raimond J M and Haroche S 1997 *Phys. Rev. Lett.* **79** 1
 Raimond J M, Brune M and Haroche S 2001 *Rev. Mod. Phys.* **73** 565
- [13] van Enk S J, Cirac J I and Zoller P 1997 *Phys. Rev. Lett.* **79** 5178
 Cabrillo C, Cirac J I, García-Fernández P and Zoller P 1999 *Phys. Rev. A* **59** 1025

- Duan L-M, Kuzmich A and Kimble H J 2002 *Phys. Rev. A* **67** 032305
- [14] Plenio M, Huelga S F, Beige A and Knight P L 1999 *Phys. Rev. A* **59** 2468
Jane E, Plenio M B and Jonathan D 2002 *Phys. Rev. A* **65** 050302
Browne D E, Plenio M B and Huelga S F 2003 *Phys. Rev. Lett.* **91** 067901
- [15] Osnaghi S *et al* 2001 *Phys. Rev. Lett.* **87** 037902
- [16] Sørensen A S and Mølmer K 2003 *Phys. Rev. Lett.* **90** 127903
Solano E, Agarwal G S and Walther H 2003 *Phys. Rev. Lett.* **90** 027903
Unanyan R G and Fleischhauer M 2003 *Phys. Rev. Lett.* **90** 133601
Hong J and Lee H-W 2002 *Phys. Rev. Lett.* **89** 237901
Sørensen A S and Mølmer K 2003 *Phys. Rev. Lett.* **91** 097905
Clark S, Peng A, Gu M and Parkins S 2003 *Phys. Rev. Lett.* **91** 177901
- [17] Wootters W K 1998 *Phys. Rev. Lett.* **80** 2245
- [18] Beige A, Munro W J and Knight P L 2000 *Phys. Rev. A* **62** 052102
Majumdar A S and Nayak N 2001 *Phys. Rev. A* **64** 013821
- [19] Bell J S 1965 *Physics (N. Y.)* **1** 195
- [20] Clauser J F, Horne M A, Shimony A and Holt R A 1969 *Phys. Rev. Lett.* **23** 880
- [21] Horodecki M, Horodecki P and Horodecki R 1995 *Phys. Lett. A* **200** 340
- [22] Gardiner C W and Zoller P 2000 *Quantum Noise* (Berlin: Springer)
- [23] Milburn G J 1991 *Phys. Rev. A* **44** 5401
and Schneider S and Milburn G J 1998 *Phys. Rev. A* **57** 3748
Schneider S and Milburn G J 1999 *Phys. Rev. A* **59** 3766