

Tachyon vacuum solution in open string field theory with a constant B field

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2009 J. Phys. A: Math. Theor. 42 395402

(<http://iopscience.iop.org/1751-8121/42/39/395402>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.155

The article was downloaded on 03/06/2010 at 08:10

Please note that [terms and conditions apply](#).

Tachyon vacuum solution in open string field theory with a constant B field

Akira Ishida¹, Chanju Kim², Yoonbai Kim³, O-Kab Kwon³ and D D Tolla³

¹ Graduate School of Mathematics, Nagoya University, Nagoya 464-6802, Japan

² Institute for the Early Universe and Department of Physics, Ewha Womans University, Seoul 120-750, Korea

³ Department of Physics, BK21 Physics Research Division, Institute of Basic Science, and College University, Sungkyunkwan University, Suwon 440-746, Korea

E-mail: ishida@math.nagoya-u.ac.jp, cjkim@ewha.ac.kr, yoobai@skku.edu, okab@skku.edu and ddtolla@skku.edu

Received 1 June 2009, in final form 13 August 2009

Published 8 September 2009

Online at stacks.iop.org/JPhysA/42/395402

Abstract

We show that Schnabl's tachyon vacuum solution is an exact solution of the equation of motion of Witten's open bosonic string field theory in the background of a constant antisymmetric two-form field. The action computed at the vacuum solution is given by the Dirac–Born–Infeld factor multiplied to that without the antisymmetric tensor field.

PACS number: 11.25.Sq

1. Introduction

Recently there has been much interest in Witten's open string field theory (OSFT) for bosonic strings [1]. It is a natural framework to discuss the nonperturbative physics of D-branes as demonstrated in [2]. Several conjectures were made on tachyon condensation in [3], [4], and the open-string tachyon was identified with the unstable mode of the D-brane. The first conjecture states that the tension of the D-brane is given by the height of the tachyon potential from the tachyon vacuum. In OSFT, Schnabl proved it by constructing an analytic tachyon vacuum solution [5]. Since then, Schnabl's method [6, 7] has been further developed to prove Sen's third conjecture [8] as well as to study related problems analytically in OSFT, including rolling tachyon [9, 10–21].

In this paper, we would like to consider Witten's OSFT in the background of a constant antisymmetric two-form field B and find the tachyon vacuum solution. This background has attracted attention since, in the presence of the B field, the end points of the open strings become noncommutative and low energy dynamics of the D-branes is described by noncommutative

gauge theories [22]. Witten’s OSFT in the constant B field was studied in [23–25], and it was shown that the action has the same form as the corresponding action on trivial background but it has Moyal-type noncommutativity in addition to the ordinary Witten’s star product and string coupling constant [23, 24]. Since Schnabl’s tachyon vacuum solution [5] belongs to the universal subspace of the string field spanned by the modes of the bc ghosts and the Virasoro generators [4], one may expect that it should still be a solution even in the background of the constant B field. In this paper, we verify this explicitly by showing that Schnabl’s solution satisfies the equation of motion of the OSFT with Moyal-type noncommutativity. Then, by direct substitution of the solution into the action, we identify the tension of D25-brane in the presence of the constant B field, which is nothing but the D25-brane tension multiplied by the Dirac–Born–Infeld (DBI) factor $\sqrt{-\det(\eta + 2\pi\alpha' B)}$. This result is obtained through a nontrivial combination of the overall normalization factor in the OSFT action which comes from the relation between the open string couplings with and without B [22], and the Jacobian factor coming from the integration of zero modes.

Since the B field appears only through the gauge invariant combination $B+F$ where F is the field strength on the D-brane, this result is consistent with the DBI action obtained by summing over all disk diagrams with the external constant F .

2. Constant B field and tachyon vacuum

Dynamics of bosonic strings including off-shell contributions is described by Witten’s OSFT [1], in which the action is

$$S(\Phi) = -\frac{1}{\alpha'^3 g_0^2} \int \left(\frac{1}{2} \Phi * Q_B \Phi + \frac{1}{3} \Phi * \Phi * \Phi \right) \tag{2.1}$$

$$= -\frac{1}{\alpha'^3 g_0^2} \left(\frac{1}{2} {}_{12} \langle V_2 || \Phi \rangle_1 Q_B | \Phi \rangle_2 + \frac{1}{3} {}_{123} \langle V_3 || \Phi \rangle_1 \Phi \rangle_2 \Phi \rangle_3 \right), \tag{2.2}$$

where g_0 , Φ , $*$ and Q_B in the first line 2.1 are the open string coupling constant, a string field, Witten’s star product and the BRST charge, respectively. In the second line, ${}_{12\dots n} \langle V_n |$ is the n -string overlap vertex and $| \Phi \rangle_n$ is the n th string state. From (2.2), the equation of motion is

$${}_{12} \langle V_2 | Q_B | \Phi \rangle_2 + {}_{123} \langle V_3 || \Phi \rangle_2 | \Phi \rangle_3 = 0. \tag{2.3}$$

The analytic tachyon vacuum solution is known to be [5]

$$| \Psi \rangle = \lim_{N \rightarrow \infty} \left(\sum_{n=0}^N \psi'_n - \psi_N \right), \tag{2.4}$$

where ψ_n is expressed in terms of ghosts and wedge state $|n\rangle$ ($n = 0, 1, \dots$),

$$\psi_n = \frac{2}{\pi} c_1 |0\rangle * |n\rangle * B_1^L c_1 |0\rangle, \tag{2.5}$$

and $\psi'_n \equiv \frac{d\psi_n}{dn}$. B_1^L is represented in terms of b ghost as

$$B_1^L = \int_{C_L} \frac{d\xi}{2\pi i} (1 + \xi^2) b(\xi), \tag{2.6}$$

where the contour C_L runs counterclockwise along the unit circle with $\text{Re}(\xi) < 0$. In (2.4), the coefficient of the so-called phantom piece ψ_N has to be -1 for the solution to satisfy the equation of motion when contracted with itself [6, 7]:

$${}_{12} \langle V_2 || \Psi \rangle_1 Q_B | \Psi \rangle_2 + {}_{123} \langle V_3 || \Psi \rangle_1 | \Psi \rangle_2 | \Psi \rangle_3 = 0. \tag{2.7}$$

Note that this solution belongs to the universal subspace of the string field spanned by the modes of the bc ghosts and the Virasoro generators.

Evaluation of the action for the Schnabl's solution gives

$$S(\Psi) = \frac{1}{6\alpha'^3 g_0^2} {}_{123}\langle V_3 | | \Psi \rangle_1 | \Psi \rangle_2 | \Psi \rangle_3 = \frac{\text{Vol}_{26}}{2\pi^2 \alpha'^3 g_0^2} \equiv T_{25} \text{Vol}_{26}, \quad (2.8)$$

where $\text{Vol}_{26} = \int d^{26}x$ represents the spacetime volume factor and T_{25} is the tension of 25-dimensional space-filling brane. Apart from the volume factor, this coincides with the tension of D25-brane, consistent with Sen's conjecture [3].

In the presence of a constant antisymmetric two-form field, the string worldsheet action is written as

$$\frac{1}{2\pi\alpha'} \int d^2z (g_{MN} - 2\pi\alpha' B_{MN}) \partial X^M \bar{\partial} X^N + S_{\text{gh}}, \quad (2.9)$$

where S_{gh} is the b, c ghost contribution. We denote by the Greek indices μ, ν the directions that the antisymmetric tensor field is nonzero, $B_{\mu\nu} \neq 0$. The classical solution of the equation of motion under the boundary condition $E_{\mu\nu} \partial_{\bar{z}} X^\nu = (E^T)_{\mu\nu} \partial_z X^\nu$ with $E_{\mu\nu} = g_{\mu\nu} + 2\pi\alpha' B_{\mu\nu}$ is given by

$$\begin{aligned} X^\mu(z, \bar{z}) = & \tilde{x}^\mu - i\alpha' [(E^{-1})^{\mu\nu} \ln \bar{z} + (E^{-1T})^{\mu\nu} \ln z] p_\nu \\ & + i\sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \frac{1}{n} [(E^{-1})^{\mu\nu} \alpha_{n,\nu} \bar{z}^{-n} + (E^{-1T})^{\mu\nu} \alpha_{n,\nu} z^{-n}], \end{aligned} \quad (2.10)$$

where \tilde{x}^μ 's are noncommutative coordinates satisfying

$$[\tilde{x}^\mu, \tilde{x}^\nu] = i\theta^{\mu\nu}, \quad \theta^{\mu\nu} = -(2\pi\alpha')^2 (E^{-1} B E^{-1T})^{\mu\nu}. \quad (2.11)$$

The commutation relations among operators are

$$[\alpha_{n,\nu}, \alpha_{m,\nu}] = n G_{\mu\nu} \delta_{n+m,0}, \quad [x^\mu, p_\nu] = i\delta_\nu^\mu, \quad (2.12)$$

where $G_{\mu\nu}$ is the inverse matrix of $G^{\mu\nu} = (E^{-1} g E^{-1T})^{\mu\nu}$ and $x^\mu = \tilde{x}^\mu + \frac{1}{2}\theta^{\mu\nu} p_\nu$ is the center of mass coordinate.

The corresponding OSFT action in the constant B field background is [23, 24]

$$\begin{aligned} S_B = & -\frac{1}{\alpha'^3 G_0^2} \int \left(\frac{1}{2} \Phi \star \tilde{Q}_B \Phi + \frac{1}{3} \Phi \star \Phi \star \Phi \right) \\ = & -\frac{1}{\alpha'^3 G_0^2} \left(\frac{1}{2} {}_{12}\langle \hat{V}_2 | | \Phi \rangle_1 \tilde{Q}_B | \Phi \rangle_2 + \frac{1}{3} {}_{123}\langle \hat{V}_3 | | \Phi \rangle_1 | \Phi \rangle_2 | \Phi \rangle_3 \right), \end{aligned} \quad (2.13)$$

where G_0 is the open string coupling in the presence of the constant B field [22],

$$G_0^2 = g_0^2 \sqrt{\frac{\det(g + 2\pi\alpha' B)}{\det g}}. \quad (2.14)$$

In S_B , \star in the first line is Witten's star product modified by the Moyal-type noncommutativity and ${}_{12\dots n}\langle \hat{V}_n |$ in the second line is the n -string overlap vertex corresponding to the product \star in the B field background. The BRST charge \tilde{Q}_B is not affected by the B field except that the metric in \tilde{Q}_B changes to the open string metric $G_{\mu\nu}$. The equation of motion of (2.13) is

$${}_{12}\langle \hat{V}_2 | \tilde{Q}_B | \Phi \rangle_2 + {}_{123}\langle \hat{V}_3 | | \Phi \rangle_2 | \Phi \rangle_3 = 0, \quad (2.15)$$

where the mode expansion of ${}_{12}\langle \hat{V}_2 |$, \tilde{Q}_B and ${}_{123}\langle \hat{V}_3 |$ are expressed in terms of $G_{\mu\nu}$, $\alpha_{n,\mu}$, p_μ , x^μ , b_n and c_n defined above.

Now, we find a nontrivial homogeneous solution of this equation which represents the tachyon vacuum. We expect that Schnabl's solution will satisfy the equation of motion, since it is in the universal subspace as seen above. Let us first introduce new modes [23]

$$\hat{\alpha}_n^\mu = (E^{-1T})^{\mu\nu} \alpha_{n,\nu}, \quad \hat{p}^\mu = (E^{-1T})^{\mu\nu} p_\nu, \quad \hat{x}_\mu = E_{\mu\nu} x^\nu. \quad (2.16)$$

Then the commutation relations become those without the B field,

$$[\hat{\alpha}_n^\mu, \hat{\alpha}_m^\nu] = n g^{\mu\nu} \delta_{n+m,0}, \quad [\hat{x}_\nu, \hat{p}^\mu] = i \delta_\nu^\mu, \quad (2.17)$$

and the bilinear form of α_n^μ with metric $G_{\mu\nu}$ is replaced by that of the transformed modes $\hat{\alpha}_n^\mu$ with metric $g_{\mu\nu}$, i.e.

$$G^{\mu\nu} \alpha_{n,\mu} \alpha_{m,\nu} = g_{\mu\nu} \hat{\alpha}_n^\mu \hat{\alpha}_m^\nu, \quad (2.18)$$

where $\alpha_{0,\mu} = \sqrt{2\alpha'} p_\mu$. Then, ${}_{12}\langle \hat{V}_2 |$ and ${}_{123}\langle \hat{V}_3 |$ are rewritten in terms of the new modes, $g_{\mu\nu}$, $\hat{\alpha}_n^\mu$, \hat{p}^μ , \hat{x}_μ , b_n and c_n , and are related to those in the absence of the B field as [23, 24]

$${}_{12}\langle \hat{V}_2 | = {}_{12}\langle V_2 |, \quad {}_{123}\langle \hat{V}_3 | = {}_{123}\langle V_3 | \exp\left(-\frac{i}{2} \sum_{r<s} \theta^{\mu\nu} p_\mu^{(r)} p_\nu^{(s)}\right). \quad (2.19)$$

Also, \tilde{Q}_B has the same form as Q_B with the oscillators replaced by the new ones.

With these forms of overlap vertices and the BRST charge, it is not difficult to see that the Schnabl's tachyon vacuum solution (2.4) is still the solution of the equation of motion (2.15) even in the presence of the constant B field. For this, we need to show that (2.15) holds when contracted with any state in the Fock space as well as with the solution itself [6, 7]. First note that the kinetic term reduces to that in the absence of the B field due to (2.19) and (2.17). Furthermore, since the tachyon vacuum solution is a homogeneous solution [5], we have $p_\mu^{(r)} |\Psi\rangle_r = 0$ and hence the Moyal-type noncommutativity disappears in the cubic term. Then, the resulting equation coincides with the equation without the background B field 2.7 for homogeneous string fields. This is true for the contraction with either any state in the Fock space or the string field itself. This completes the proof.

From now on, let us discuss the physical properties of the solution. In the absence of the B field, the calculation of the action (2.1) for the Schnabl's analytic vacuum solution gives the exact value of the tension T_{25} of 25-dimensional space-filling brane [5] as in (2.8). Since all the B field dependence in the action (2.13) appears only through the open string coupling constant G_o and the transformed center of mass coordinate \hat{x}^μ , the value of the action (2.13) at the tachyon vacuum solution with the constant B field gives

$$\begin{aligned} S_B(\Phi = \Psi) &= \frac{1}{6\alpha'^3 G_o^2} {}_{123}\langle V_3 | |\Psi\rangle_1 |\Psi\rangle_2 |\Psi\rangle_3 = \frac{1}{2\pi^2 \alpha'^3 G_o^2} \int d^{26} \hat{x} \\ &= \frac{\text{Vol}_{26}}{2\pi^2 \alpha'^3 G_o^2} |\det(\eta + 2\pi\alpha' B)|, \end{aligned}$$

where $\eta_{\mu\nu}$ is the flat spacetime metric and $|\det(\eta + 2\pi\alpha' B)|$ is the Jacobian factor for the coordinate transformation given in (2.16). Now, we use the relation $G_o^2 = \sqrt{-\det(\eta + 2\pi\alpha' B)} g_o^2$ from (2.14), which gives

$$S_B(\Phi = \Psi) = T_{25} \text{Vol}_{26} \sqrt{-\det(\eta + 2\pi\alpha' B)}. \quad (2.21)$$

Since we are considering the case of a constant B field, this result is consistent with the DBI action obtained by summing over all the disk diagrams with external gauge field legs of constant electromagnetic field strength $F_{\mu\nu}$ as boundary terms [26], considering the fact that the B field appears only through the gauge invariant combination $B_{\mu\nu} + F_{\mu\nu}$. The result (2.21) is natural because Schnabl's tachyon vacuum solution in the presence of the constant B field background is an exact analytic solution of the classical SFT equation and does not involve any off-shell contribution.

3. Conclusion

In this paper, we showed that Schnabl's tachyon vacuum solution is an exact solution of the string field equation in the presence of the constant B field. Since the effect of the constant B field appears only in the open string coupling and the Jacobian factor of spacetime integration, the value of the SFT action is easily computed and gives the DBI Lagrangian density multiplied by the D25-brane tension and spacetime volume, which coincides with the result of disc amplitude computation in string theory.

The obtained result may create a new direction in OSFT including both constant B field background and marginal deformations of either time or spatial dependence. It would be intriguing if the treatment of the B field becomes applicable to the cases with time and spatial dependence, which lead to higher-derivative corrections [27]. Since fundamental strings couple minimally to the antisymmetric tensor field B in string theories, our work on the OSFT in the presence of the constant B field may contribute to the understanding of fundamental strings in the context of OSFT.

Acknowledgments

This work was supported by Astrophysical Research Center for the Structure and Evolution of the Cosmos (ARCSEC) and grant no R01-2006-000-10965-0 from the Basic Research Program through the Korea Science and Engineering Foundation (AI, OK), by the World Class University grant R32-2008-000-10130-0 (CK), by the Science Research Center Program of the Korea Science and Engineering Foundation through the Center for Quantum Spacetime (CQeST) with grant numbers R11-2005-021 and KRF-2006-352-C00010 (CK, DT) and by Faculty Research Fund, Sungkyunkwan University, 2007 (YK).

References

- [1] Witten E 1986 Noncommutative geometry and string field theory *Nucl. Phys. B* **268** 253
- [2] Sen A and Zwiebach B 2000 Tachyon condensation in string field theory *J. High Energy Phys.* [JHEP03\(2000\)002](#) (arXiv:[hep-th/9912249](#))
- [3] Sen A 1999 Descent relations among bosonic D-branes *Int. J. Mod. Phys. A* **14** 4061 (arXiv:[hep-th/9902105](#))
- [4] Sen A 1999 Universality of the tachyon potential *J. High Energy Phys.* [JHEP12\(1999\)027](#) (arXiv:[hep-th/9911116](#))
- [5] Schnabl M 2006 Analytic solution for tachyon condensation in open string field theory *Adv. Theor. Math. Phys.* **10** 433 (arXiv:[hep-th/0511286](#))
- [6] Okawa Y 2006 Comments on Schnabl's analytic solution for tachyon condensation in Witten's open string field theory *J. High Energy Phys.* [JHEP04\(2006\)055](#) (arXiv:[hep-th/0603159](#))
- [7] Fuchs E and Kroyter M 2006 On the validity of the solution of string field theory *J. High Energy Phys.* [JHEP05\(2006\)006](#) (arXiv:[hep-th/0603195](#))
- [8] Ellwood I and Schnabl M 2007 Proof of vanishing cohomology at the tachyon vacuum *J. High Energy Phys.* [JHEP02\(2007\)096](#) (arXiv:[hep-th/0606142](#))
- [9] Rastelli L and Zwiebach B 2008 Solving open string field theory with special projectors *J. High Energy Phys.* [JHEP01\(2008\)020](#) (arXiv:[hep-th/0606131](#))
Okawa Y, Rastelli L and Zwiebach B 2006 Analytic solutions for tachyon condensation with general projectors arXiv:[hep-th/0611110](#)
- [10] Schnabl M 2007 Comments on marginal deformations in open string field theory *Phys. Lett. B* **654** 194 (arXiv:[hep-th/0701248](#))
- [11] Kiermaier M, Okawa Y, Rastelli L and Zwiebach B 2008 Analytic solutions for marginal deformations in open string field theory *J. High Energy Phys.* [JHEP01\(2008\)028](#) (arXiv:[hep-th/0701249](#))
- [12] Erler T 2007 Marginal solutions for the superstring *J. High Energy Phys.* [JHEP07\(2007\)050](#) (arXiv:[0704.0930](#) [hep-th])

- [13] Okawa Y 2007 Analytic solutions for marginal deformations in open superstring field theory *J. High Energy Phys.* **JHEP09(2007)084** (arXiv:0704.0936 [hep-th])
Okawa Y 2007 Real analytic solutions for marginal deformations in open superstring field theory *J. High Energy Phys.* **JHEP09(2007)082** (arXiv:0704.3612 [hep-th])
- [14] Ellwood I 2007 Rolling to the tachyon vacuum in string field theory *J. High Energy Phys.* **JHEP12(2007)028** (arXiv:0705.0013 [hep-th])
- [15] Fuchs E, Kroyter M and Potting R 2007 Marginal deformations in string field theory *J. High Energy Phys.* **JHEP09(2007)101** (arXiv:0704.2222 [hep-th])
Fuchs E and Kroyter M 2007 Marginal deformation for the photon in superstring field theory *J. High Energy Phys.* **JHEP11(2007)005** (arXiv:0706.0717 [hep-th])
- [16] Kiermaier M and Okawa Y 2007 Exact marginality in open string field theory: a general framework arXiv:0707.4472 [hep-th]
Kiermaier M and Okawa Y 2007 General marginal deformations in open superstring field theory arXiv:0708.3394 [hep-th]
- [17] Erler T 2008 Tachyon vacuum in cubic superstring field theory *J. High Energy Phys.* **JHEP01(2008)013** (arXiv:0707.4591 [hep-th])
- [18] Lee B H, Park C and Tolla D D 2007 Marginal deformations as lower dimensional D-brane solutions in open string field theory arXiv:0710.1342 [hep-th]
- [19] Kwon O K 2008 Marginally deformed rolling tachyon around the tachyon vacuum in open string field theory arXiv:0801.0573 [hep-th]
- [20] Bagchi A and Sen A 2008 Tachyon condensation on separated brane-antibrane system arXiv:0801.3498 [hep-th]
- [21] HELLERMAN S and SCHNABL M 2008 Light-like tachyon condensation in open string field theory arXiv:0803.1184 [hep-th]
- [22] Seiberg N and Witten E 1999 String theory and noncommutative geometry *J. High Energy Phys.* **JHEP09(1999)032** (arXiv:hep-th/9908142)
- [23] Sugino F 2000 Witten's open string field theory in constant B-field background *J. High Energy Phys.* **JHEP03(2000)017** (arXiv:hep-th/9912254)
- [24] Kawano T and Takahashi T 2000 Open string field theory on noncommutative space *Prog. Theor. Phys.* **104** 459 (arXiv:hep-th/9912274)
- [25] Kawano T and Takahashi T 2000 Open-closed string field theory in a background B field *Prog. Theor. Phys.* **104** 1267 (arXiv:hep-th/0005080)
Schnabl M 2000 String field theory at large B-field and noncommutative geometry *J. High Energy Phys.* **JHEP11(2000)031** (arXiv:hep-th/0010034)
Bonora L, Mamone D and Salizzoni M 2002 B field and squeezed states in vacuum string field theory *Nucl. Phys. B* **630** 163 (arXiv:hep-th/0201060)
Bonora L, Mamone D and Salizzoni M 2002 Vacuum string field theory with B field *J. High Energy Phys.* **JHEP04(2002)020** (arXiv:hep-th/0203188)
Furuuchi K 2002 Non-commutative space and Chan–Paton algebra in open string field algebra *Nucl. Phys. B* **640** 145 (arXiv:hep-th/0202200)
Maccaferri C, Scherer Santos R J and Tolla D D 2005 Time-localized projectors in string field theory with E-field *Phys. Rev. D* **71** 066007 (arXiv:hep-th/0501011)
- [26] Fradkin E S and Tseytlin A A 1985 Nonlinear electrodynamics from quantized strings *Phys. Lett. B* **163** 123
- [27] Coletti E, Sigalov I and Taylor W 2003 Abelian and nonabelian vector field effective actions from string field theory *J. High Energy Phys.* **JHEP09(2003)050** (arXiv:hep-th/0306041)