

## Ternutator identities

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2009 J. Phys. A: Math. Theor. 42 475209

(<http://iopscience.iop.org/1751-8121/42/47/475209>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

### Download details:

IP Address: 171.66.16.156

The article was downloaded on 03/06/2010 at 08:24

Please note that [terms and conditions apply](#).

# Ternutator identities

Chandrashekar Devchand<sup>1</sup>, David Fairlie<sup>2</sup>, Jean Nuyts<sup>3</sup> and Gregor Weingart<sup>4</sup>

<sup>1</sup> Institut für Mathematik der Universität Potsdam, Am Neuen Palais 10, D-14469 Potsdam, Germany

<sup>2</sup> Department of Mathematical Sciences, University of Durham, Science Laboratories, South Rd, Durham DH1 3LE, UK

<sup>3</sup> Physique Théorique et Mathématique, Université de Mons-Hainaut, 20 Place du Parc, B-7000 Mons, Belgium

<sup>4</sup> Instituto de Matemáticas, Universidad Nacional Autónoma de México, 62210 Cuernavaca, Morelos, Mexico

E-mail: [devchand@math.uni-potsdam.de](mailto:devchand@math.uni-potsdam.de), [david.fairlie@durham.ac.uk](mailto:david.fairlie@durham.ac.uk), [jean.nuyts@umh.ac.be](mailto:jean.nuyts@umh.ac.be) and [gw@matcuer.unam.mx](mailto:gw@matcuer.unam.mx)

Received 27 August 2009, in final form 7 October 2009

Published 6 November 2009

Online at [stacks.iop.org/JPhysA/42/475209](http://stacks.iop.org/JPhysA/42/475209)

## Abstract

The ternary commutator or ternutator, defined as the alternating sum of the product of three operators, has recently drawn much attention as an interesting structure generalizing the commutator. The ternutator satisfies cubic identities analogous to the quadratic Jacobi identity for the commutator. We present various forms of these identities and discuss the possibility of using them to define ternary algebras.

PACS number: 02.10.Ud

Mathematics Subject Classification: 16R10

## 1. Introduction

Some time ago Nambu [1] introduced antisymmetric brackets depending on three functions and a derivation, generalizing the Poisson bracket and satisfying the Nambu identity, which generalizes the Jacobi identity. These brackets have been discussed, for example, in [2–4]. Nambu also considered the possible quantization of his brackets: for three arbitrary operators  $A_1, A_2, A_3$  (with an associative product) in an arbitrary vector space, he defined the related ternary commutator  $[A_1, A_2, A_3]$ , which we call the *ternutator*, by the alternating sum over the permutations of (1, 2, 3),

$$\begin{aligned}
 [A_1, A_2, A_3] &:= A_1 A_2 A_3 + A_2 A_3 A_1 + A_3 A_1 A_2 - A_1 A_3 A_2 - A_2 A_1 A_3 - A_3 A_2 A_1 \\
 &=: \sum_{\pi \in \mathcal{S}_3} \text{sgn}(\pi) A_{\pi(1)} A_{\pi(2)} A_{\pi(3)}. \tag{1}
 \end{aligned}$$

This can be expressed in terms of the commutator as

$$\begin{aligned} [A_1, A_2, A_3] &= A_1[A_2, A_3] + A_2[A_3, A_1] + A_3[A_1, A_2] \\ &= [A_2, A_3]A_1 + [A_3, A_1]A_2 + [A_1, A_2]A_3. \end{aligned} \tag{2}$$

The ternutator is trilinear in the operators and has the property of being totally skew-symmetric,

$$[A_1, A_2, A_3] = -[A_2, A_1, A_3] = -[A_3, A_2, A_1], \tag{3}$$

which leads to the cyclic property,  $[A_1, A_2, A_3] = [A_2, A_3, A_1] = [A_3, A_1, A_2]$ . This bracket, as a particular case of those introduced by Fillipov in [5], has recently attracted renewed interest in the physics literature (e.g. [6–11]). Earlier physical applications of ternary algebraic structures are reviewed in [12].

The ternutator generalizes the commutator, which is the binary operation underlying Lie algebras. The Jacobi identity plays a crucial role in the classification of Lie algebras. In this paper, we present identities for the ternutator, analogous to the Jacobi identity for the commutator. This problem has also been discussed by others, e.g. [13, 14]. All  $n$ -linear brackets defined by the alternating sum of the product of any odd number  $n \geq 5$  of operators have recently been shown [15] to satisfy similar identities. Just as the Jacobi identity provides the defining relations for Lie algebras, we discuss the possibility of using the ternutator identities to define ternary algebras. Some specific examples of sets of structure constants satisfying the ternutator identity are given in section 4.

## 2. Ternutators and their identities

The Jacobi identity for commutators, i.e. the equality of the two ways of writing the ternutator in terms of the commutator in (2), can be expressed as an alternating sum over permutations of (1, 2, 3),

$$\sum_{t \in S_3} \epsilon_{t_1 t_2 t_3} [[A_{t_1}, A_{t_2}], A_{t_3}] = 2 \sum_{\substack{t \in S_3 \\ t_1 < t_2}} \epsilon_{t_1 t_2 t_3} [[A_{t_1}, A_{t_2}], A_{t_3}] = 0, \tag{4}$$

where the second sum is over all permutations  $(t_1, t_2, t_3)$  of (1, 2, 3) satisfying the condition  $t_1 < t_2$  and  $\epsilon_{t_1 t_2 t_3}$  is the completely antisymmetric Levi Civita symbol.

The following result for the ternutator (1) is well known (e.g. [13]).

**Lemma 1.** *There are no identities of second order in the ternutator.*

**Proof.** The most general second-order identity is

$$\sum_{\substack{t \in S_5 \\ t_1 < t_2 < t_3, t_4 < t_5}} c(t_1, t_2, t_3, t_4, t_5) [[A_{t_1}, A_{t_2}], A_{t_3}], A_{t_4}, A_{t_5}] = 0, \tag{5}$$

where the sum is over all permutations  $(t_1, \dots, t_5)$  of (1, \dots, 5) satisfying the conditions  $t_1 < t_2 < t_3$  and  $t_4 < t_5$ . This has ten summands, multilinear in the five operators  $A_i, i = 1, \dots, 5$ . It is easy to check that these ten terms are linearly independent. Hence the ten coefficients  $c(t_1, t_2, t_3, t_4, t_5)$  must be zero.  $\square$

There are two types of monomials cubic in the ternutator, both involving seven operators. Label any set of seven operators (not necessarily linearly independent)  $A_1, \dots, A_7$  and define

$$T_1(A_1, \dots, A_7) := [[[A_1, A_2, A_3], A_4, A_5], A_6, A_7] \tag{6}$$

$$T_2(A_1, \dots, A_7) := [[A_1, A_2, A_3], [A_4, A_5, A_6], A_7]. \tag{7}$$

In virtue of the symmetry properties of the ternutator, there are 210 independent monomials of the form  $T_1(A_{l_1}, A_{l_2}, A_{l_3}, A_{l_4}, A_{l_5}, A_{l_6}, A_{l_7})$  labelled by the permutations of  $(l_1, \dots, l_7)$  of  $(1, \dots, 7)$  satisfying  $(l_1 < l_2 < l_3)$ ,  $(l_4 < l_5)$  and  $(l_6 < l_7)$ , and 70 monomials of the form  $T_2(A_{l_1}, A_{l_2}, A_{l_3}, A_{l_4}, A_{l_5}, A_{l_6}, A_{l_7})$ , with conditions  $(l_1 < l_2 < l_3)$ ,  $(l_4 < l_5 < l_6)$ ,  $(l_1 < l_4)$ . An identity cubic in ternutators, if it exists, is a linear dependence amongst these 280 terms:

$$\sum_{\substack{\iota \in S_7 \\ \iota_1 < \iota_2 < \iota_3, \iota_4 < \iota_5, \iota_6 < \iota_7}} c_1(l_1, l_2, l_3, l_4, l_5) T_1(A_{l_1}, A_{l_2}, A_{l_3}, A_{l_4}, A_{l_5}, A_{l_6}, A_{l_7}) + \sum_{\substack{\iota \in S_7 \\ \iota_1 < \iota_2 < \iota_3, \iota_1 < \iota_4 < \iota_5 < \iota_6}} c_2(l_1, l_2, l_3, l_7) T_2(A_{l_1}, A_{l_2}, A_{l_3}, A_{l_4}, A_{l_5}, A_{l_6}, A_{l_7}) = 0, \tag{8}$$

where the sums are over all permutations  $\iota \in S_7, \iota : n \mapsto \iota_n, n \in \{1, \dots, 7\}$ , satisfying the given restrictions. The following result was also known to Bremner [13].

**Theorem 1.** *Among any seven operators, there exist precisely seven independent identities cubic in the ternutator.*

In order to prove this, we have used REDUCE and independently MAPLE to find that the 280 coefficients in (8) are constrained by 273 independent linear relations leaving a seven-parameter space of identities. Hence there is a basis of seven independent identities, which generalize the usual Jacobi identity for commutators.

Consider seven arbitrary operators and label them  $A_i, i = 1, \dots, 7$ . Pick out one of them, say  $A_7$ , and consider the following alternating sums of  $T_1$  and  $T_2$ , (skew) symmetric under permutations of the remaining six operators:

$$I_1(l_1, \dots, l_6; 7) = \sum_{\substack{\iota \in S_6 \\ \iota_1 < \iota_2 < \iota_3, \iota_4 < \iota_5 < \iota_6}} \epsilon_{\iota_1 \iota_2 \iota_3 \iota_4 \iota_5 \iota_6} [[A_{l_{\iota_1}}, A_{l_{\iota_2}}, A_{l_{\iota_3}}], A_{l_{\iota_4}}, A_7], A_{l_{\iota_5}}, A_{l_{\iota_6}}] \\ I_2(l_1, \dots, l_6; 7) = \sum_{\substack{\iota \in S_6 \\ \iota_1 < \iota_2 < \iota_3, \iota_4 < \iota_5}} \epsilon_{\iota_1 \iota_2 \iota_3 \iota_4 \iota_5 \iota_6} [[A_{l_{\iota_1}}, A_{l_{\iota_2}}, A_{l_{\iota_3}}], [A_{l_{\iota_4}}, A_{l_{\iota_5}}, A_7], A_{l_{\iota_6}}]. \tag{9}$$

The identity is then

$$I = I_1 + I_2 \equiv 0. \tag{10}$$

An alternative way of writing the identity is as follows. Take seven operators  $A_a, A_b, A_c, A_d, A_e, A_f, A_g$  and single out  $A_g$ . We can write the identity  $I$  as the alternating sum over all permutations of the six indices  $(a, b, c, d, e, f)$ ,

$$12 I = \sum_{S(a,b,c,d,e,f)} \text{sgn}(\pi) ([[[A_a, A_b, A_c], A_d, A_g], A_e, A_f] + [[A_a, A_b, A_c], [A_d, A_e, A_g], A_f]) \equiv 0, \tag{11}$$

where  $\text{sgn}(\pi)$  is the sign of the permutation. The factor 12 here is the number of times a given term is repeated in the sum over all permutations. The seven independent identities correspond to the seven possibilities of singling out any one, say  $A_g$ . These seven identities transform under  $S_7$  as the direct sum of the representations associated with the Young

tableaux

$$\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} \oplus \begin{array}{|c|} \hline \square & \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} \tag{12}$$

These identities were known to Bremner [13] in a slightly different form. They were independently rediscovered in this explicit form by one of us (JN). There exist further equivalent expressions for these identities. Again singling out  $A_7$ , we may write

$$\begin{aligned} & 3 \sum_{\substack{\mu \in S_6 \\ \mu_1 < \mu_2 < \mu_3, \mu_4 < \mu_5 < \mu_6, \mu_1 < \mu_4}} \epsilon_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6} [[A_{\mu_1}, A_{\mu_2}, A_{\mu_3}], [A_{\mu_4}, A_{\mu_5}, A_{\mu_6}], A_7] \\ &= \sum_{\substack{\mu \in S_7 \\ \mu_1 < \mu_2 < \mu_3, \mu_4 < \mu_5 < 7, \mu_6 < \mu_7}} \epsilon_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6 \mu_7} [[[A_{\mu_1}, A_{\mu_2}, A_{\mu_3}], A_{\mu_4}, A_{\mu_5}], A_{\mu_6}, A_{\mu_7}] \\ &\quad - 2 \sum_{\substack{\mu \in S_7 \\ \mu_1 < \mu_2 < \mu_3, \mu_5 = 7, \mu_6 < \mu_7}} \epsilon_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6 \mu_7} [[[A_{\mu_1}, A_{\mu_2}, A_{\mu_3}], A_{\mu_4}, A_{\mu_5}], A_{\mu_6}, A_{\mu_7}]. \end{aligned} \tag{13}$$

The left-hand side has ten summands. In the first term of the right-hand side there are 150 summands and in the second term 60 summands. This linear dependence amongst monomials cubic in the ternutator has alternative expression:

$$\begin{aligned} & 3 \sum_{\substack{\mu \in S_6 \\ \mu_1 < \mu_2 < \mu_3, \mu_4 < \mu_5 < \mu_6, \mu_1 < \mu_4}} \epsilon_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6} [[A_{\mu_1}, A_{\mu_2}, A_{\mu_3}], [A_{\mu_4}, A_{\mu_5}, A_{\mu_6}], A_7] \\ &= \sum_{\substack{\mu \in S_6 \\ \mu_1 < \mu_2, \mu_3 < \mu_4, \mu_5 < \mu_6}} \epsilon_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6} [[[A_{\mu_1}, A_{\mu_2}, A_7], A_{\mu_3}, A_{\mu_4}], A_{\mu_5}, A_{\mu_6}] \\ &\quad + \sum_{\substack{\mu \in S_6 \\ \mu_1 < \mu_2 < \mu_3, \mu_4 < \mu_5}} \epsilon_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6} [[[A_{\mu_1}, A_{\mu_2}, A_{\mu_3}], A_{\mu_4}, A_{\mu_5}], A_{\mu_6}, A_7] \\ &\quad - 2 \sum_{\substack{\mu \in S_6 \\ \mu_1 < \mu_2 < \mu_3, \mu_5 < \mu_6}} \epsilon_{\mu_1 \mu_2 \mu_3 \mu_4 \mu_5 \mu_6} [[[A_{\mu_1}, A_{\mu_2}, A_{\mu_3}], A_{\mu_4}, A_7], A_{\mu_5}, A_{\mu_6}]. \end{aligned} \tag{14}$$

Here, there are ten summands on the left-hand side, 90 terms in the first part of the right-hand side, 60 terms in the second part and 60 in the third part.

### 3. Ternary algebras

A ternary algebra is a vector space in which a ternary composition is given by the ‘ternutation relations’,

$$[e_i, e_j, e_k] = t_{ijk}^m e_m, \tag{15}$$

where  $e_i$  are basis elements and the structure constants  $t_{ijk}^m$  are completely antisymmetric in  $i, j, k$  and transform as (3, 1)-tensors under a nonsingular change of basis,  $e'_j = S_j^k e_k$ ,

$\det(S) \neq 0$ . Inserting these basis elements in the polynomials  $I_1$  and  $I_2$  in (9) yields cubic polynomials in the structure constants,

$$\begin{aligned}
 I_1(t; a, b, c, d, e, f; g, q) &= \sum_{S(a,b,c,d,e,f)} \text{sgn}(\pi) t_{abc}^p t_{pdg}^n t_{nef}^q \\
 I_2(t; a, b, c, d, e, f; g, q) &= \sum_{S(a,b,c,d,e,f)} \text{sgn}(\pi) t_{abc}^m t_{deg}^n t_{mnf}^q, \tag{16}
 \end{aligned}$$

where the sum is over all permutations of  $(a, b, c, d, e, f)$ . The ternutator identities (11) then yield identities for the structure constants  $t_{ijk}^m$ ,

$$I(t; a, b, c, d, e, f; g, q) := (I_1 + I_2)(t; a, b, c, d, e, f; g, q) = 0. \tag{17}$$

Just as a set of structure constants satisfying the Jacobi identities converts a vector space into a Lie algebra, it would be interesting if the identities (17) similarly provide necessary and sufficient conditions for a set of structure constants  $\{t_{ijk}^m\}$  to define a ternary algebra with relations (15).

The associativity of the multiplication implies the quadratic Jacobi identity for the commutator (4) as well as the cubic identities (11) for the ternutator. If we lift associativity, both these identities do not hold. For the alternative multiplication for the basis elements of the imaginary octonions,  $e_i e_j = -\delta_{ij} + \psi_{ijk} e_k$ , the left-hand side of the Jacobi identity (4) is proportional to the associator  $(e_i, e_j, e_k) = (e_i e_j) e_k - e_i (e_j e_k) = \varphi_{ijkl} e_l$ , where both  $\psi_{ijk}$  and  $\varphi_{ijkl}$  are completely antisymmetric. The ternutator  $[e_i, e_j, e_k]$  is also proportional to the associator, and we have verified by direct calculation using REDUCE that the  $G_2$ -invariant structure constants of the associator algebra  $\varphi_{ijkl} = \frac{1}{6} \epsilon_{ijklmnp} \psi_{mnp}$  with nonzero components,

$$\varphi_{1234} = \varphi_{1256} = \varphi_{1357} = \varphi_{1476} = \varphi_{2376} = \varphi_{2475} = \varphi_{3456} = 1, \tag{18}$$

corresponding to the octonion structure constants  $\{\psi_{127} = \psi_{163} = \psi_{154} = \psi_{253} = \psi_{246} = \psi_{347} = \psi_{567} = 1\}$  indeed do not satisfy the identities (17). Instead, the two polynomials in (16) evaluated for  $\varphi$  are equal.

Indeed, in dimensions 6 and 7, for antisymmetric  $t$ 's, the values of the two polynomials in (16) are equal, so in these dimensions the identities (17) are equivalent to either  $I_1 = 0$  or  $I_2 = 0$ . We have checked explicitly that the equality  $I_1 = I_2 \neq 0$  also holds for the components of the Spin(7)-invariant 4-form [16, 17] defined by (18) together with the further nonzero components  $\varphi_{ijk8} = \psi_{ijk}$ , namely,  $\varphi_{1278} = \varphi_{1386} = \varphi_{1485} = \varphi_{2385} = \varphi_{2468} = \varphi_{3478} = \varphi_{5678} = 1$ , and hence this 4-form also does not satisfy the ternutator identity.

Let  $\pi$  be a permutation of the four indices  $(a, b, c, d)$ . The following quadratic equations in the structure constants

$$\sum_{\pi \in S(a,b,c,d)} \text{sgn}(\pi) t_{abc}^m t_{mdg}^q = 0 \tag{19}$$

are related to the above-mentioned Nambu identity and have been advocated [11] as defining conditions for algebras (1). These equations are by no means necessary conditions for the existence of the ternary algebra since the ternutator does not satisfy a quadratic identity (see theorem 1). However, it is readily seen that if (19) holds then our cubic necessary conditions (11) are satisfied. Hence the relations (19) could be sufficient (but by no means necessary) to guarantee the existence of the corresponding ternary algebra.

The Cartan–Killing metric plays a crucial role in the classification of Lie algebras. There are several tensors, constructed from the structure constants  $t_{ijk}^m$ , which possibly could play a similar role for ternary algebras. For Lie algebras, in terms of the adjoint map of element  $e_j$  with respect to an element  $e_i$  defined by  $\text{ad}_{e_i} e_j := [e_i, e_j]$ , the Cartan–Killing form is given

by  $g(e_i, e_j) := \text{Tr}(\text{ad}_{e_i} \circ \text{ad}_{e_j})$ . In terms of the structure constants of the Lie algebra we have  $g_{ij} = f_{im}^n f_{jn}^m$ . Using the ternutation relations (15) to define an analogous endomorphism of the ternary algebra,  $X_{e_i, e_j} e_k := [e_i, e_j, e_k]$ , yields the most natural candidate for a ‘metric’, namely  $\text{Tr}(X_{e_i, e_j} \circ X_{e_k, e_l})$ , which is clearly antisymmetric in  $ij$ , antisymmetric in  $kl$  and symmetric under interchange of these ordered pairs (in other words, it has the symmetries of the Riemann tensor). This object has components

$$g_{ij,kl} = t_{ijn}^m t_{klm}^n \tag{20}$$

and can serve as a metric on the  $N(N-1)/2$ -dimensional space of 2-forms indexed by  $\{ij\}$ ,  $i < j$ . If this metric is invertible, its inverse  $g^{ij,kl}$  then satisfies

$$\sum_{k < l} g_{ij,kl} g^{kl,mn} = \delta_i^m \delta_j^n \tag{21}$$

and affords the construction of the further symmetric tensor,

$$g^{pq} = \frac{1}{2} (t_{ijk}^p g^{ij,kq} + t_{ijk}^q g^{ij,kp}), \tag{22}$$

which is a candidate for a metric in the linear space of the operators.

If a Lie algebra is semi-simple, a particularly important subset of Lie algebras, it is well known that the metric  $g_{ij} = f_{ia}^b f_{jb}^a$  formed from the structure constants is nonsingular and that  $f_{ijk} = g_{ka} f_{ij}^a$  is completely antisymmetric in its indices. It would be remarkable if analogous statements hold for ternary algebras. In particular, we expect a special role for ternary algebras having the following two properties:

- (a) there exists a basis in the space of operators such that the structure constants are completely antisymmetric,

$$[e_i, e_j, e_k] = t_{ijkm} e_m, \quad \text{with } t_{ijkm} \text{ antisymmetric,} \tag{23}$$

- (b) in this basis, a metric appears in the form

$$\sum_{a < b < c} t_{abc} t_{jabc} = \lambda \delta_{ij}. \tag{24}$$

#### 4. Examples

The conditions (17) are so overdetermined that a complete classification of solutions, without imposing further conditions, seems to be a rather difficult task. However, isolated solutions in lower dimensions can certainly be found. We display four solutions in six dimensions with the structure constants being given by components of a 4-form  $t_{ijkm}$ , which satisfy the identities (17). It is convenient to use the Hodge-dual 2-form  $\tilde{t} = * t$  with components  $\tilde{t}_{a_1 a_2} = \frac{1}{4!} \epsilon_{a_1 a_2 a_3 a_4 a_5 a_6} t_{a_3 a_4 a_5 a_6}$ .

##### Example 1

$$\tilde{t}_{12} = -\tilde{t}_{13} = -\tilde{t}_{14} = \tilde{t}_{15} = -\tilde{t}_{23} = -\tilde{t}_{24} = \tilde{t}_{26} = \tilde{t}_{35} = -\tilde{t}_{36} = \tilde{t}_{45} = -\tilde{t}_{46} = \tilde{t}_{56} = 1. \tag{25}$$

##### Example 2

$$-\tilde{t}_{12} = \tilde{t}_{13} = \tilde{t}_{14} = -\tilde{t}_{15} = \tilde{t}_{26} = -\tilde{t}_{36} = -\tilde{t}_{46} = \tilde{t}_{56} = 1. \tag{26}$$

##### Example 3

$$\tilde{t}_{14} = \tilde{t}_{15} = -\tilde{t}_{16} = -\tilde{t}_{24} - \tilde{t}_{25} = \tilde{t}_{26} = \tilde{t}_{34} = \tilde{t}_{35} = -\tilde{t}_{36} = 1. \tag{27}$$

**Example 4**

$$\begin{aligned}
\tilde{t}_{46} &= -1, & \tilde{t}_{56} &= 1, \\
\tilde{t}_{36} &= x_1, & \tilde{t}_{26} &= -x_3, & \tilde{t}_{35} &= -x_2, & \tilde{t}_{34} &= x_2, \\
\tilde{t}_{25} &= x_4, & \tilde{t}_{24} &= -x_4, & \tilde{t}_{23} &= x_4x_1 - x_2x_3.
\end{aligned} \tag{28}$$

In the first three examples, the structure constants satisfy the calibration criterion

$$t_{abcd}^3 = t_{abcd} \implies \tilde{t}_{ab}^3 = \tilde{t}_{ab}. \tag{29}$$

Example 4 does not seem to be a calibration.

**5. Conclusions**

Ternutators and ternary algebras generalize the familiar commutators and Lie algebras and have recently appeared in various contexts in the literature (see e.g. [18]). In this paper, we have obtained explicit forms of the seven identities, cubic in ternutators, which are obeyed by any seven operators with an associative product. We have shown that these identities exhaust the space of identities for seven operators. It remains an open question whether higher order ( $q \geq 4$ ) identities involving  $2q + 1$  operators are trivial consequences of the seven basic identities. The seven independent identities for seven operators clearly imply identities (of fourth order in the ternutator) for nine operators. As is clear from (11), the former seven identities can be considered as operations  $X(B_1, \dots, B_6; B_7)$ , depending on seven operators, skew-symmetric in the first six. Amongst the identities for any given choice of nine operators  $A_i, i = 1 \dots 9$  there exist those having one of the following structures:  $A = [X(A_1, \dots, A_6; A_7), A_8, A_9]$ ,  $B = X(A_1, \dots, A_6; [A_7, A_8, A_9])$ ,  $C = X(A_1, \dots, A_5, [A_6, A_8, A_9]; A_7)$ . There are  $7 \cdot \binom{9}{2} = 252$  identities of type A, and  $\binom{9}{3} = 84$  identities of type B and type C. So altogether 420 identities amongst 9 operators have their origins in the identities for seven operators. Are there any further identities?

As a consequence of these identities, the structure constants of ternary algebras must obey cubic relations, which are highly overdetermined. We have given some examples of solutions of these identities in six dimensions.

**Acknowledgments**

Two of us (JN and GW) acknowledge partial funding from the SFB 647 ‘Raum-Zeit-Materie’ of the Deutsche Forschungsgemeinschaft for a visit to Potsdam. JN also thanks the Belgian FNRS for support and Winfried Neun of the Konrad-Zuse-Zentrum für Informationstechnik Berlin (ZIB) for help with REDUCE. GW is supported by the Mexican DGAPA Research Project IN115408 *Geometría Riemanniana Global* and CONACyT Project 82471.

**References**

- [1] Nambu Y 1973 Generalized Hamiltonian mechanics *Phys. Rev. D* **7** 2405–12
- [2] Bayen F and Flato M 1975 Remarks concerning Nambu’s generalized mechanics *Phys. Rev. D* **11** 3049–53
- [3] Takhtajan L 1994 On foundation of the generalized Nambu mechanics *Commun. Math. Phys.* **160** 295 (arXiv:hep-th/9301111)
- [4] Pioline B 2002 Comments on the topological open membrane *Phys. Rev. D* **66** 025010 (arXiv:hep-th/0201257)
- [5] Filippov V T 1985 n-Lie algebras *Siberian Math. J.* **26** 879–91
- [6] Curtright T and Zachos C 2003 Classical and quantum Nambu mechanics *Phys. Rev. D* **68** 085001
- [7] Okubo S 1995 Introduction to octonion and other non-associative algebras in physics (*Montroll Memorial Lecture Series in Mathematical Physics*) (Cambridge: Cambridge University Press)



- [8] Bagger J and Lambert N 2008 Gauge symmetry and supersymmetry of multiple M2-branes *Phys. Rev. D* **77** 065008 (arXiv:0711.0955)
- [9] Curtright T L, Fairlie D B and Zachos C K 2008 Ternary Virasoro–Witt algebra *Phys. Lett. B* **666** 386 (arXiv:0806.3515)
- [10] Curtright T L, Jin X, Mezincescu L, Fairlie D B and Zachos C K 2009 Classical and quantal ternary algebras *Phys. Lett. B* **675** 387 (arXiv:0903.4889)
- [11] Bagger J and Lambert N 2009 Three-algebras and  $\mathcal{N} = 6$  Chern–Simons gauge theories *Phys. Rev. D* **79** 025002 (arXiv:0807.0163)
- [12] Kerner R 2000 Ternary algebraic structures and their applications in physics arXiv:math-ph/0011023
- [13] Bremner M 1998 Identities for the ternary commutator *J. Algebra* **206** 615–23
- [14] Bremner M and Peresi L A 2006 Ternary analogues of Lie and Malcev algebras *Linear Algebra Appl.* **414** 1–18
- [15] Curtright T, Jin X and Mezincescu L 2009 Multi-operator brackets acting thrice arXiv:0905.2759
- [16] Corrigan E, Devchand C, Fairlie D B and Nuyts J 1983 First order equations for gauge fields in spaces of dimension greater than four *Nucl. Phys. B* **214** 452
- [17] Devchand C, Nuyts J and Weingart G 2009 Matryoshka of special democratic forms *Commun. Math. Phys.* (arXiv:0812.3012) at press (doi:10.1007/s00220-009-0939-5)
- [18] Ataguema H 2008 Classification et déformations des algèbres ternaires *PhD thesis* Université de Haute Alsace