

On the linearization of the first and second Painlevé equations

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2009 J. Phys. A: Math. Theor. 42 055208

(<http://iopscience.iop.org/1751-8121/42/5/055208>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.156

The article was downloaded on 03/06/2010 at 08:27

Please note that [terms and conditions apply](#).

On the linearization of the first and second Painlevé equations

N Joshi¹, A V Kitaev² and P A Treharne¹

¹ School of Mathematics and Statistics, The University of Sydney, New South Wales 2006, Australia

² Steklov Mathematical Institute, Fontanka 27, St Petersburg, 191023, Russia

E-mail: N.Joshi@maths.usyd.edu.au, kitaev@pdmi.ras.ru and P.Treharne@maths.usyd.edu.au

Received 2 June 2008, in final form 26 November 2008

Published 6 January 2009

Online at stacks.iop.org/JPhysA/42/055208

Abstract

We found Fuchs–Garnier pairs in 3×3 matrices for the first and second Painlevé equations which are linear in the spectral parameter. As an application of our pairs for the second Painlevé equation we use the generalized Laplace transform to derive an invertible integral transformation relating two of its Fuchs–Garnier pairs in 2×2 matrices with different singularity structures, namely, the pair due to Jimbo and Miwa and that found by Harnad, Tracy and Widom. Together with the certain other transformations it allows us to relate all known 2×2 matrix Fuchs–Garnier pairs for the second Painlevé equation to the original Garnier pair.

PACS numbers: 02.30.Gp, 02.30.Hq, 02.30.Ik

Mathematics Subject Classification: 33E17, 34M25, 34M55

(Some figures in this article are in colour only in the electronic version)

1. Introduction

The phrase ‘linearization of the Painlevé equations’ is widely understood to refer to the fact that the nonlinear Painlevé equations can be associated with certain overdetermined systems of linear differential equations in two complex variables. The linear systems may be written as scalar or matrix equations and are typically referred to as Lax pairs for the Painlevé equations. In our previous paper [10], we suggested calling these systems the ‘Fuchs–Garnier’ pairs for the Painlevé equations to pay tribute to the two scientists who first introduced these systems in the beginning of the twentieth century. In the period since Fuchs and Garnier first wrote their (scalar) pairs the list has expanded so that now, for most of the Painlevé equations, there are several different Fuchs–Garnier pairs associated with the same Painlevé equation³.

³ Actually most of the ‘new’ pairs appeared as similarity reductions of various Lax pairs for nonlinear partial differential equations integrable via the inverse scattering transform method which explains why they are often referred to as Lax pairs for the Painlevé equations.

The different Fuchs–Garnier pairs for a given Painlevé equation may differ from each other not only by simple gauge transformation but also by the matrix dimension and/or analytic structure (the number and type of singular points). Our general belief is that all Fuchs–Garnier pairs for a given Painlevé equation should be equivalent in the sense that there should exist explicit transformations that map these pairs to each other. In many cases such transformations are known; however, there are still several instances of different Fuchs–Garnier pairs whose equivalence is expected but not yet established. The main goal of this work is to construct such an explicit transformation between two Fuchs–Garnier pairs for the second Painlevé equation:

$$P_2 : \frac{d^2y}{dt^2} = 2y^3 + ty + \alpha, \tag{1.1}$$

where $\alpha \in \mathbb{C}$ is a complex parameter. Both pairs were obtained in 1979–1980: one pair is due to Flaschka and Newell [3] (FN-pair) and the other is due to Jimbo and Miwa [9] (JM₂-pair). The FN-pair was originally obtained as a similarity reduction of the Lax pair for the modified KdV equation; the JM₂-pair was originally obtained from the scalar pair of Garnier, although it can also be obtained as a similarity reduction of the Lax pair for the nonlinear Schrödinger equation.

The solution of the aforementioned problem associated with P_2 is intimately connected with the other central theme of this work, namely, the construction of the so-called secondary linearized Fuchs–Garnier pairs for P_2 and the first Painlevé equation:

$$P_1 : \frac{d^2y}{dt^2} = 6y^2 + t. \tag{1.2}$$

The notion of secondary linearized Fuchs–Garnier pairs for the Painlevé equations was introduced in our previous work [10] and refers to Fuchs–Garnier pairs which are *linear* in the spectral parameter λ ; see system (1.5) below. However, it is important to mention that the question concerning a relation between the FN- and JM₂- pairs was one of the main motivations for both our works.

We recall that the matrix Fuchs–Garnier pairs for the Painlevé equations have the following form:

$$\frac{dY}{d\lambda} = \mathcal{A}(\lambda, t)Y, \quad \frac{dY}{dt} = \mathcal{U}(\lambda, t)Y, \tag{1.3}$$

where $\lambda \in \mathbb{C}$ is an auxiliary variable called the spectral parameter, and $\mathcal{A}(\lambda, t), \mathcal{U}(\lambda, t) \in GL(N, \mathbb{C})$ are *rational* functions of λ and are analytic in t . Jimbo and Miwa [9] showed that for all Painlevé equations such pairs exist in 2×2 matrices. The Frobenius compatibility condition of system (1.3),

$$\mathcal{A}_t - \mathcal{U}_\lambda + [\mathcal{A}, \mathcal{U}] = 0, \tag{1.4}$$

where $[\cdot, \cdot]$ is the usual matrix commutator, being imposed identically for all values of λ , is equivalent to one of the Painlevé equations.

As is mentioned above Fuchs–Garnier pairs can be scalar and matrix. The scalar pairs for P_1 and P_2 were obtained by Garnier [5]. All other scalar Fuchs–Garnier pairs that we know in the literature are related to the Garnier pairs via simple transformations. The situation with the matrix Fuchs–Garnier pairs for P_1 is also fairly easy to summarize. The basic 2×2 matrix pair for P_1 (JM₁-pair) was found by Jimbo and Miwa [9]. It is straightforward to prove that the JM₁-pair is equivalent to the scalar pair obtained by Garnier. Moreover, the other 2×2 matrix pairs for P_1 that can be found in the literature are related to the JM₁-pair by the Fabri [2, 8] and Schlesinger transformations [9] (in the other terminology via quadratic *RS*-transformations [12]). As for the matrix Fuchs–Garnier pairs for P_2 the situation is more interesting. To the best of our knowledge all matrix pairs for P_2 that were discussed so far in

the literature are given in 2×2 matrices. There are three different matrix pairs: the FN- and JM_2 -pairs mentioned already in the previous paragraph, and that obtained by Harnad, Tracy and Widom [6] (HTW-pair). We give a detailed account of all these pairs in section 3. It is straightforward to establish that the Fabri transformation maps the HTW-pair into the FN-pair (see details in section 3) and that the JM_2 -pair is a matrix version of the scalar Garnier pair (see appendix A). However, a direct link between the FN- (equivalently HTW-) and the JM_2 -pairs is not that obvious; this link is one of the main matters of our paper.

Central to our investigation of this problem are the so-called secondary linearized Fuchs–Garnier pairs, the technique that we began to develop in our previous work [10]. In that work, we found the secondary linearized Fuchs–Garnier pairs in 3×3 matrices for the third, fourth and fifth Painlevé equations. In this paper, we complete the list of the secondary linearized pairs for the Painlevé equations by adding to it the pairs for P_1 and P_2 . The secondary linearized Fuchs–Garnier pair for the sixth Painlevé equation was known earlier due to Harnad [7] (see also [14]).

Secondary linearization is here taken to mean presenting each of the Painlevé equations in terms of a Fuchs–Garnier pair with an equation on the spectral parameter λ of the following form:

$$(\lambda B_1(t) + B_2(t)) \frac{d\Psi}{d\lambda} = (\lambda B_3(t) + B_4(t)) \Psi, \quad (1.5)$$

i.e., with coefficients *linear* with respect to λ . In [10], using the similarity reductions of the Lax pair for the three-wave resonant interaction (3WRI) system, we obtained secondary linearized Fuchs–Garnier pairs in 3×3 matrices for all the Painlevé equations except P_1 and P_2 . As there are no similarity reductions of the 3WRI system to P_1 and P_2 , this approach could not be used to obtain secondary linearized pairs for the latter equations. So in this paper we complete a list of the secondary linearized Fuchs–Garnier pairs for the Painlevé equations.

The main advantage of the secondary linearized pairs is that the Laplace transform maps them one into another without changing the matrix dimension of the corresponding Fuchs–Garnier pairs. This property is lost for Fuchs–Garnier pairs which are rational functions of the spectral parameter of degree more than 1. In this work, the fact that the matrix dimension is not altered is the key to constructing an explicit link between the JM_2 - and the FN-pairs for P_2 since the two secondary linearized 3×3 matrix pairs that are related by the generalized Laplace transform can be reduced independently to the different 2×2 matrix pairs. The reduction of our secondary linearized 3×3 matrix pairs to the 2×2 matrix Fuchs–Garnier pairs is done via two different mechanisms: (i) by a special normalization of equation (1.5) and (ii) from a degeneracy that occurs under an application of the generalized Laplace transform to equation (1.5). We call a secondary linearized Fuchs–Garnier pair *degenerate* iff $\det(\lambda B_1(t) + B_2(t)) \equiv 0$ for all $\lambda \in \mathbb{C}$. The reader will see that, in the case of P_1 and P_2 , the generalized Laplace transform maps nondegenerate Fuchs–Garnier pairs into degenerate ones.

In section 2, we present a nondegenerate secondary linearized Fuchs–Garnier pair for P_1 (JKT_1 -pair). We show that under the formal⁴ Laplace transform it maps to a degenerate Fuchs–Garnier pair ($dJKT_1$ -pair). We then show that the $dJKT_1$ -pair is equivalent to the JM_1 -pair, which is, in turn, a matrix form of the original Garnier pair, G_1 . Schematically, the content of section 2 can be described by the graph shown in figure 1, where the vertices represent the corresponding Fuchs–Garnier pairs, and the edges are invertible transformations relating them.

⁴ We call a Laplace transform formal in case we do not specify its contour of integration.



Figure 1. The diagram of the Fuchs–Garnier pairs for P_1 and mappings between them.




Figure 2. The commutative diagram of the Fuchs–Garnier pairs for P_2 and corresponding mappings.

The rest of the paper is devoted to P_2 . Schematically, its content can be presented by the graph shown in figure 2. The vertices and the edges of the graph are different Fuchs–Garnier pairs for P_2 and the mappings between them, respectively. The vertices abbreviated as JKT with sub- and superscripts denote 3×3 matrix Fuchs–Garnier pairs that we have constructed. The subscript {2} means the pair for P_2 , the prefix d , as above, says that the corresponding pair is degenerate, and the superscripts {1, 2, 3} label different degenerate Fuchs–Garnier pairs. The graph is commutative and all mappings are invertible. The edges without any name correspond to the reduction transformation from the 3×3 to 2×2 matrix and the 2×2 matrix to scalar Fuchs–Garnier pairs and their inverses. Our main result is the diagonal mapping indicated by the red edge. It is obtained in two ways: as the composition of transformations along the upper and lower roots connecting vertices JM_2 and HTW . These compositions coincide which proves the commutativity of our diagram.

This diagram is constructed in sections 3–5 and appendices appendix A and B.

In section 3, we recall the main subjects of our study—the 2×2 matrix Fuchs–Garnier pairs for P_2 due to Jimbo and Miwa (JM_2 -pair), Flaschka and Newell (FN-pair), and Harnad, Tracy and Widom (HTW -pair). We also show that the Fabri transformation [2, 8], which is natural to employ for the HTW -pair and is well known in asymptotic theory, maps the HTW -pair to the FN-pair. This fact was earlier noted in [11]. Finally, we conclude this section with a presentation of the main result of this paper, i.e., we give a direct invertible integral transformation mapping the HTW -pair to the JM_2 -pair. Appendix A completes the general overview of the 2×2 matrix pairs by showing a relation of the JM_2 -pair to the scalar Garnier pair (G_2 -pair). Here we also consider one more Fuchs–Garnier pair for P_2 in 2×2 matrices obtained by Conte and Musette [1] (CM_2 -pair) and show how it can be mapped directly to the JM_2 -pair without reference to the scalar G_2 -pair.

In section 4, we present two new 3×3 matrix secondary linearized Fuchs–Garnier pairs for P_2 , one of which is nondegenerate, together with the corresponding integral transformation between them and their reductions to the JM_2 - and HTW -pairs. Using these results we construct a formal integral transform between the JM_2 - and HTW -pairs.

In section 5, we present two other new 3×3 matrix secondary linearized Fuchs–Garnier pairs for P_2 , different from those in section 4. Both pairs are degenerate and are related via the generalized Laplace transform. As in the previous section, we construct transformations relating these pairs to the JM_2 - and HTW-pairs and on this basis obtain exactly the same formal integral transform between the latter pairs as in section 4.

Finally in appendix B, we show how to find a contour of integration in our integral transformations which completes the proof of our main result.

2. Fuchs–Garnier pairs for P_1

In subsection 2.1, we construct the Fuchs–Garnier pairs and corresponding mappings presented by the graph in figure 1. In subsection 2.2, we discuss the Fabri-transformed JM_1 -pairs that appeared in the literature and present a new and simpler version of the Fabri transformation.

2.1. Secondary linearization of P_1

Proposition 2.1. *Consider the following system of linear ODEs:*

$$\text{JKT}_1 : \begin{cases} \frac{d\Psi}{d\mu} = \left(\mu \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} -y & -z & -(4y^2 + 2t) \\ 0 & y & z \\ 1/4 & 0 & 0 \end{pmatrix} \right) \Psi \\ \frac{d\Psi}{dt} = 2 \left(\mu \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & -y & -z \\ -1/4 & 0 & -2y \\ 0 & -1/4 & 0 \end{pmatrix} \right) \Psi, \end{cases} \quad (2.1)$$

where $y = y(t)$ and $z = z(t)$ are analytic functions of t . Then the compatibility condition reads

$$\frac{dy}{dt} = z, \quad \frac{dz}{dt} = 6y^2 + t, \quad (2.2)$$

i.e. is equivalent to equation (1.2).

Proof. The straightforward check of the Frobenius compatibility condition (1.4) with $\lambda \rightarrow \mu$. \square

In our terminology this is a nondegenerate secondary linearized Fuchs–Garnier pair for P_1 .

Let us make the generalized Laplace transform of the Fuchs–Garnier pair (2.1) with respect to the variable μ ,

$$\Psi(\mu, t) = \int_L e^{\lambda\mu} \Phi(\lambda, t) d\lambda, \quad (2.3)$$

with such contour L chosen to make the certain off-integral terms vanish⁵. The result reads

$$\text{dJKT}_1 : \begin{cases} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \frac{d\Phi}{d\lambda} = \left(-\lambda \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \begin{pmatrix} -y & -z & -(4y^2 + 2t) \\ 0 & y & z \\ 1/4 & 0 & 0 \end{pmatrix} \right) \Phi, \\ \frac{d\Phi}{dt} = 2 \left(\lambda \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ -1/4 & 0 & -2y \\ 0 & -1/4 & 0 \end{pmatrix} \right) \Phi. \end{cases} \quad (2.4)$$

⁵ Here we do not discuss the choice of this contour, so that we leave this transformation at the formal level. The notation is explained in more detail in appendix B, where the appropriate choice of contour for the case of P_2 is given.

In our terminology this is a degenerate Fuchs–Garnier pair. To formulate our next result we recall the Fuchs–Garnier pair for P_1 found by Jimbo and Miwa [9]:

$$JM_1 : \begin{cases} \frac{dY}{d\lambda} = \left(\lambda^2 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 & y \\ 4 & 0 \end{pmatrix} + \begin{pmatrix} -z & y^2 + \frac{t}{2} \\ -4y & z \end{pmatrix} \right) Y, \\ \frac{dY}{dt} = \left(\frac{\lambda}{2} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & y \\ 2 & 0 \end{pmatrix} \right) Y. \end{cases} \quad (2.5)$$

Proposition 2.2. *The Fuchs–Garnier pair (2.4) is equivalent to the JM_1 -pair (2.5).*

Proof. The third row of the λ -equation in system (2.4) gives the following relation between the elements of the functions Φ :

$$\Phi_1 = 4\lambda\Phi_3.$$

Using this relation to eliminate Φ_1 from system (2.4) and defining $Y = (-\Phi_2/4, \Phi_3)^T$ we find that Y solves the JM_1 -pair (2.5). \square

We conclude this subsection by mentioning that if the vector solution to system (2.5) is written as $Y = (Y_1, Y_2)^T$, then the function $V = V(\lambda, t)$ defined as $Y_2 = \sqrt{\lambda - y}V$ satisfies the original Garnier pair for P_1 :

$$G_1 : \begin{cases} \frac{d^2V}{d\lambda^2} = \left(\frac{3}{4(\lambda - y)^2} - \frac{y'}{\lambda - y} + 4\lambda^3 + 2t\lambda + (y')^2 - 4y^3 - 2ty \right) V, \\ \frac{dV}{dt} = \frac{1}{2(\lambda - y)} \frac{dV}{d\lambda} + \frac{1}{4(\lambda - y)^2} V. \end{cases}$$

2.2. The Fabri-type transformation for the JM_1 -pair

Since the matrix coefficient of λ^2 in the right-hand side of the JM_1 -pair (2.5) has zero determinant and zero trace, it is standard in asymptotic theory to apply the Fabri-type transformation [2, 8] $\lambda = \zeta^2$ to ‘cure the defect’ at infinity. It was first applied to system (2.5) by Jimbo and Miwa in the same work [9] where the JM_1 -pair was obtained (see p 437) under the name of the ‘shearing’ transformation. As a result they obtain an equation (see (C.5) in [9]) with an additional apparent Fuchsian singularity at the origin. Although this does not cause any problems for the application of the isomonodromy deformation technique in studying, say, asymptotics of P_1 ,⁶ this form of the Fabri-transformed JM_1 -pair does create problems in the application of the Riemann–Hilbert approach. This is due to the fact that the corresponding connection matrix (the matrix connecting fundamental solutions at the singular points zero and infinity) for this pair now depends on the solution of P_1 . This problem was first identified by Fokas, Mugan and Zhou [4]. To correct this additional problem these authors introduced one more gauge transformation depending on a spectral parameter, and thus they produced another Fuchs–Garnier pair (FMZ-pair) for P_1 . The approach of [4] necessitated the introduction of an additional function, $v(t)$, which is related to the P_1 function $y(t)$ via the Riccati differential equation, $iv'(t) - 2v^2(t) = y(t)$. This function appeared in the FMZ-pair because in the Riemann–Hilbert setting there appears an additional parameter in the connection matrix which corresponds to the constant of integration in the Riccati equation.

Here we show that there exists a Fabri-type transformation which is free of both problems indicated in the previous paragraph: it does not have an additional Fuchsian singularity at

⁶ The monodromy matrix at the origin is just equal to $-I$, and thus does not depend on t .

the origin like the original Fabri-transformed JM_1 -pair and the FMZ-pair. Our Fabri-type transformation for the fundamental solutions of (2.5) reads

$$Y(\lambda, t) = \begin{pmatrix} 1 & -\zeta/2 \\ 0 & 1 \end{pmatrix} Z(\zeta, t), \quad \lambda = \zeta^2.$$

We find that our Fabri-type transformation maps the JM_1 -pair to the following one:

$$JM_{1/F} : \begin{cases} \frac{dZ}{d\zeta} = \left(4\zeta^4 \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \zeta^3 \begin{pmatrix} 0 & 4y \\ 8 & 0 \end{pmatrix} + \zeta^2 \begin{pmatrix} -4y & 2z \\ 0 & 4y \end{pmatrix} \right. \\ \qquad \left. + \zeta \begin{pmatrix} -2z & 2y^2 + t \\ -8y & 2z \end{pmatrix} + \begin{pmatrix} 0 & 1/2 \\ 0 & 0 \end{pmatrix} \right) Z, \\ \frac{dZ}{dt} = \left(\zeta \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \begin{pmatrix} 0 & y \\ 2 & 0 \end{pmatrix} \right) Z. \end{cases} \quad (2.6)$$

An important role in the study of the Fabri-transformed JM_1 -pairs is played by the so-called σ_1 -symmetry of the fundamental solutions related to the reflection, $\zeta \rightarrow -\zeta$. The latter symmetry is not easy to observe by looking directly at our pair (2.6); however, a method of its derivation suggests the following identity for the fundamental solutions:

$$\begin{pmatrix} 1 & -\zeta/2 \\ 0 & 1 \end{pmatrix} Z(\zeta) = Y(\lambda) = \begin{pmatrix} 1 & \zeta/2 \\ 0 & 1 \end{pmatrix} Z(-\zeta)C, \quad (2.7)$$

for some $C \in SL(2, \mathbb{C})$. Thus, the σ_1 -symmetry for our pair reads

$$Z(-\zeta) = \begin{pmatrix} 1 & -\zeta \\ 0 & 1 \end{pmatrix} Z(\zeta)\tilde{C},$$

where clearly $\tilde{C} \in SL(2, \mathbb{C})$, and thus has nothing to do any more with the Pauli matrix σ_1 !

3. The 2×2 matrix Fuchs–Garnier pairs for P_2

The rest of the paper is devoted to P_2 . Henceforth, the notation y and z means solutions of P_2 (1.1) and equation P_{34} (see equation (3.3), below), respectively. Subsections 3.1–3.4 are devoted to the review of the known results for the 2×2 matrix Fuchs–Garnier pairs for P_2 . In subsection 3.5 we formulate the main result of the paper.

3.1. The Jimbo–Miwa pair

Jimbo and Miwa [9] give the following matrix version of the Fuchs–Garnier pair for P_2 :

$$JM_2 : \begin{cases} \frac{dY}{d\lambda} = \left(\lambda^2 \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \lambda \begin{pmatrix} 0 & u \\ -2u^{-1}z & 0 \end{pmatrix} + \begin{pmatrix} z + t/2 & -uy \\ -2u^{-1}(yz + \theta) & -z - t/2 \end{pmatrix} \right) Y, \\ \frac{dY}{dt} = \left(\frac{\lambda}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 0 & u \\ -2u^{-1}z & 0 \end{pmatrix} \right) Y. \end{cases} \quad (3.1)$$

The compatibility condition (1.4) for the JM_2 -pair (3.1) reads

$$\frac{du}{dt} = -yu, \quad \frac{dy}{dt} = y^2 + z + \frac{t}{2}, \quad \frac{dz}{dt} = -2yz - \theta. \quad (3.2)$$

Excluding the functions u and z from (3.2) we find that the function y satisfies P_2 (1.1) with $\alpha = \frac{1}{2} - \theta$.

Excluding the functions u and y from (3.2) we find that the function z satisfies the following second-order equation:

$$P_{34} : \quad \frac{d^2z}{dt^2} = \frac{1}{2z} \left(\frac{dz}{dt} \right)^2 - 2z^2 - tz - \frac{\theta^2}{2z}, \quad (3.3)$$

which, up to a scaling change of z and t , coincides with the 34th equation in the classical Painlevé–Gambier list; see p 340 in [8].

For the convenience of the reader we present in appendix A a relation of the JM_2 -pair with the original scalar Garnier pair (G_2 -pair) and a direct mapping of another 2×2 matrix version of the G_2 -pair by Conte and Musette [1] (CM_2 -pair) to the JM_2 -pair.

3.2. The Flaschka–Newell pair

Flaschka and Newell [3] found the following Fuchs–Garnier pair for P_2 :

$$FN : \quad \begin{cases} \frac{dZ}{d\zeta} = \left(-4i\zeta^2\sigma_3 + 4y\zeta\sigma_1 - 2y'\sigma_2 - i(2y^2 + t)\sigma_3 - \frac{\alpha}{\zeta}\sigma_1 \right) Z, \\ \frac{dZ}{dt} = (-i\zeta\sigma_3 + y\sigma_1)Z, \end{cases} \quad (3.4)$$

where ζ is the spectral parameter, t is the dynamical variable, prime denotes differentiation with respect to t , and $\sigma_1, \sigma_2, \sigma_3$ are the standard notation for the Pauli matrices:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The compatibility condition (1.4) for the FN-pair (3.4) implies that y is a solution of P_2 (1.1).

We note that the equation in λ in the JM_2 -pair has one singularity only, an irregular singularity at infinity, while the equation in ζ in the FN-pair has two singularities: a regular singularity at $\zeta = 0$ and an irregular singularity at infinity. It follows that there does not exist an algebraic gauge transformation for generic values of the parameter α between these systems.

Let us also mention one more difference between the JM_2 - and FN-pairs, namely, the additional σ_1 -symmetry for solutions of the FN-pair,

$$Z(-\zeta) = i\sigma_1 Z(\zeta)C, \quad \text{where } C \in SL(2, \mathbb{C}). \quad (3.5)$$

3.3. The Harnad–Tracy–Widom pair

There also exists a third 2×2 matrix Fuchs–Garnier pair for P_2 , which was first given implicitly by Harnad, Tracy and Widom in [6] in connection with random matrix theory. Explicitly, this pair was presented by Kapaev and Hubert [11] and in connection with the symmetric form of P_2 by Noumi [13]. The HTW-pair may be written as

$$HTW : \quad \begin{cases} \frac{dW}{d\mu} = \left(\mu \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} -y & -(z + 2y^2 + t) \\ \frac{1}{2} & y \end{pmatrix} + \frac{1}{2\mu} \begin{pmatrix} \theta & 0 \\ z & -\theta \end{pmatrix} \right) W, \\ \frac{dW}{dt} = - \left(\mu \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} -y & 0 \\ \frac{1}{2} & y \end{pmatrix} \right) W, \end{cases} \quad (3.6)$$

where θ is a complex parameter. Compatibility of system (3.6) implies that the functions y and z satisfy the following system of nonlinear ODEs:

$$\frac{dy}{dt} = y^2 + z + \frac{t}{2}, \quad \frac{dz}{dt} = -2zy - \theta. \quad (3.7)$$

Eliminating z from this system we find that the function y satisfies the second Painlevé equation (1.1) with the parameter $\alpha = 1/2 - \theta$.

For completeness we recall the symmetric form of P_2 [13]:

$$f_0' = -2qf_0 + \alpha_0, \quad f_1' = 2qf_1 + \alpha_1, \quad q' = \frac{1}{2}(f_1 - f_0),$$

where $\alpha_0 = 1 - \theta, \alpha_1 = \theta$, and the functions $f_0 = f_0(t), f_1 = f_1(t)$ and $q = q(t)$ in our notation read

$$f_0 = z + 2y^2 + t, \quad f_1 = -z, \quad q = -y,$$

so that the HTW-pair gets a natural parametrization in terms of the ‘symmetric variables’.

3.4. The Fabri transformation

The HTW-pair (3.6) and the FN-pair (3.4) are related via a special Fabri-type transformation [11]:

$$Z(\zeta, t) = G(\zeta)W(\mu, t), \quad G(\zeta) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \left(\frac{i}{2\zeta} \right)^{\sigma_3/2}, \quad \mu = -2\zeta^2. \quad (3.8)$$

The function z in (3.6) is given by the equation $z = y' - y^2 - \frac{t}{2}$. The σ_1 -symmetry for Z (3.5) follows from the Fabri transformation because of the identity⁷,

$$G(-\zeta)G^{-1}(\zeta) = i\sigma_1.$$

3.5. Main result

Now we are ready to formulate the main result of this paper.

Theorem 3.1. *The fundamental solution $Y(\lambda, t)$ of the JM_2 -pair (3.1) is related to the fundamental solution $W(\mu, t)$ of the HTW-pair (3.6) via the following integral transform:*

$$\begin{pmatrix} -u/\mu & 0 \\ 0 & 2 \end{pmatrix} \mu^{\theta/2} W(\mu, t) = \int_L e^{-\lambda^3/3 + \lambda(\mu - t/2)} Y(\lambda, t) d\lambda, \quad (3.9)$$

where the contour L is specified in appendix B. The inverse transformation is given by the inverse Laplace transform.

The proof of this theorem is given in sections 4 and 5.

Corollary 3.2. *The relation between the JM_2 - and FN-pairs can be obtained as a composition of equations (3.8) and (3.9).*

4. A 3 × 3 Fuchs–Garnier pair for P_2

Proposition 4.1. *The compatibility condition for the linear system,*

$$JKT_2 : \begin{cases} \begin{pmatrix} -1 & \lambda + y & 0 \\ 0 & -\frac{1}{2} & \lambda \\ 0 & 0 & -1 \end{pmatrix} \frac{d\Phi}{d\lambda} = \begin{pmatrix} z + 2y^2 + t & -1 - \kappa_1 & 0 \\ -y & -\frac{1}{2}z & -1 - \kappa_2 \\ 1 & 0 & 0 \end{pmatrix} \Phi, \\ \frac{d\Phi}{dt} = \begin{pmatrix} -\lambda & \frac{1}{2}z & 1 + \kappa_2 \\ -1 & y & 0 \\ 0 & -\frac{1}{2} & -y \end{pmatrix} \Phi, \end{cases} \quad (4.1)$$

⁷ Compare with the analogous derivation for P_1 (2.7).

where $\kappa_j, j = 1, 2$ are parameters, is governed by the following system of nonlinear equations:

$$\frac{dy}{dt} = y^2 + z + \frac{t}{2}, \quad \frac{dz}{dt} = -2yz - (\kappa_1 - \kappa_2). \tag{4.2}$$

Eliminating z from system (4.2) we find that the function y satisfies the second Painlevé equation (1.1) with the parameter $\alpha = \frac{1}{2} - (\kappa_1 - \kappa_2)$.

Proof. The result follows from the Frobenius compatibility condition $\partial_t \partial_\lambda \Phi = \partial_\lambda \partial_t \Phi$. □

4.1. Reduction to the JM₂-pair

Since the compatibility condition (4.2) depends on the parameters κ_1 and κ_2 only through their difference, there is an additional degree of freedom in system (4.1). We will show that, by a special choice of the parameters κ_1, κ_2 , system (4.1) can be reduced to the JM₂-pair.

Proposition 4.2. *If $\kappa_j = -1$ for either $j = 1$ or $j = 2$ in system (4.1), then system (4.1) can be reduced to the JM₂-pair (3.1) plus a quadrature.*

Proof. To prove this statement, we first note that the coefficient matrix on the right-hand side of the λ equation in (4.1) has determinant $(1 + \kappa_1)(1 + \kappa_2)$. Setting $\kappa_j = -1$ for either $j = 1$ or $j = 2$ it follows that, upon diagonalizing the coefficient matrix, system (4.1) can be reduced to a 2×2 matrix system plus a quadrature. To simplify the following calculation we note that system (4.1) can be written in the following form:

$$\frac{d\Phi}{d\lambda} = - \begin{pmatrix} 2\lambda^2 + z + t & -\lambda z - yz - (1 + \kappa_1) & -2(1 + \kappa_2)(\lambda + y) \\ 2(\lambda - y) & -z & -2(1 + \kappa_2) \\ 1 & 0 & 0 \end{pmatrix} \Phi.$$

We present a proof only for a simpler case, $\kappa_2 = -1$, which does not require the diagonalizing procedure. It is the case we refer below in subsection 4.3. The case $\kappa_1 = -1$ is not employed in our work and left to the interested reader. So, we set $\kappa_2 = -1$ and note that the third component can be solved by the quadrature once the remaining two components are determined. The first two components of Φ satisfy the following linear 2×2 matrix system:

$$\begin{aligned} \frac{d\phi}{d\lambda} &= - \left(\lambda^2 \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 & -z \\ 2 & 0 \end{pmatrix} + \begin{pmatrix} z + t & -yz - (1 + \kappa_1) \\ -2y & -z \end{pmatrix} \right) \phi, \\ \frac{d\phi}{dt} &= -\frac{1}{2} \left(\lambda \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & -z \\ 2 & -2y \end{pmatrix} \right) \phi, \quad \phi = \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix}. \end{aligned} \tag{4.3}$$

We now make a gauge transformation in system (4.3),

$$\phi = \begin{pmatrix} 0 & \frac{1}{2} \\ -u^{-1} & 0 \end{pmatrix} \chi, \tag{4.4}$$

where the function $u(t)$ is defined by $u' = -yu$. The resulting system is given by

$$\begin{aligned} \frac{d\chi}{d\lambda} &= \left(\lambda^2 \begin{pmatrix} 0 & 0 \\ 0 & -2 \end{pmatrix} + \lambda \begin{pmatrix} 0 & u \\ -2u^{-1}z & 0 \end{pmatrix} + \begin{pmatrix} z & -uy \\ -2u^{-1}(yz + 1 + \kappa_1) & -z - t \end{pmatrix} \right) \chi, \\ \frac{d\chi}{dt} &= \frac{1}{2} \left(\lambda \begin{pmatrix} 0 & 0 \\ 0 & -2 \end{pmatrix} + \begin{pmatrix} 0 & u \\ -2u^{-1}z & 0 \end{pmatrix} \right) \chi. \end{aligned} \tag{4.5}$$

This system is gauge equivalent to the JM₂-pair given in (3.1). To see this, we define the parameter $\theta = (1 + \kappa_1)$ and make the change of variables:

$$\chi(\lambda, t) = e^{-(\lambda^3/3 + \lambda t/2)} Y(\lambda, t). \tag{4.6}$$

□

4.2. Reduction to the HTW-pair

Proposition 4.3. System (4.1) can be mapped to the HTW-pair (3.6) by the application of the generalized Laplace transform:

$$\Psi(\mu, t) = \int_L e^{\lambda\mu} \Phi(\lambda, t) d\lambda, \tag{4.6}$$

where the contour L is specified in appendix B.

Proof. We start the proof by writing system (4.1) in the following form:

$$\begin{aligned} \begin{pmatrix} -1 & \lambda + y & 0 \\ 0 & -\frac{1}{2} & \lambda \\ 0 & 0 & -1 \end{pmatrix} \frac{d\Phi}{d\lambda} &= \begin{pmatrix} z + 2y^2 + t & -1 - \kappa_1 & 0 \\ -y & -\frac{1}{2}z & -1 - \kappa_2 \\ 1 & 0 & 0 \end{pmatrix} \Phi, \\ \frac{d\Phi}{dt} &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \lambda \frac{d\Phi}{d\lambda} + \begin{pmatrix} 0 & \frac{1}{2}z & 1 + \kappa_2 \\ -1 & y & 0 \\ 0 & -\frac{1}{2} & -y \end{pmatrix} \Phi. \end{aligned} \tag{4.7}$$

Substituting (4.6) into (4.7), and assuming that the contour L can be chosen to eliminate any remainder terms that arise from integration by parts, we find

$$dJKT_2^3 : \begin{cases} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \mu \frac{d\Psi}{d\mu} = \begin{pmatrix} \mu - (z + 2y^2 + t) & -\mu y + \kappa_1 & 0 \\ y & \frac{1}{2}\mu + \frac{1}{2}z & \kappa_2 \\ -1 & 0 & \mu \end{pmatrix} \Psi, \\ \frac{d\Psi}{dt} = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \mu \frac{d\Psi}{d\mu} + \begin{pmatrix} 0 & \frac{1}{2}z & \kappa_2 \\ -1 & y & 0 \\ 0 & -\frac{1}{2} & -y \end{pmatrix} \Psi. \end{cases} \tag{4.8}$$

In our terminology, system (4.8) is a degenerate secondary linearized Fuchs–Garnier pair for P_2 . The third row of the μ equation in (4.8) gives the following relation between the elements of the function Ψ :

$$\Psi_1 = \mu \Psi_3.$$

Using this relation to eliminate Ψ_1 from the above equations, we find that the remaining components of Ψ satisfy the following linear 2×2 matrix system:

$$\begin{aligned} \frac{d\psi}{d\mu} &= \left(\mu \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} -y & -(z + 2y^2 + t) \\ \frac{1}{2} & y \end{pmatrix} + \frac{1}{\mu} \begin{pmatrix} \kappa_1 & 0 \\ \frac{1}{2}z & \kappa_2 \end{pmatrix} \right) \psi, \\ \frac{d\psi}{dt} &= - \left(\mu \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} -y & 0 \\ \frac{1}{2} & y \end{pmatrix} \right) \psi, \quad \psi = \begin{pmatrix} \Psi_2 \\ \Psi_3 \end{pmatrix}, \end{aligned} \tag{4.9}$$

which is gauge equivalent to the HTW-pair given in (3.6). □

4.3. Integral transform between the JM_2 - and the HTW-pairs

In this section, we construct explicitly the integral transform which maps the 2×2 system of Jimbo–Miwa into the 2×2 system of Harnad–Tracy–Widom.

Theorem 4.4. The function $W(\mu, t)$, which solves the HTW-pair given in (3.6), is related to the function $Y(\lambda, t)$, which solves the JM_2 -pair given in (3.1), via the integral transform (3.9).

Proof. From proposition 4.2, we note that the function $Y(\lambda, t)$ is related to the function $\phi(\lambda, t)$ in system (4.3) via the gauge transformation:

$$\phi(\lambda, t) = \begin{pmatrix} 0 & \frac{1}{2} \\ -u^{-1} & 0 \end{pmatrix} e^{-(\lambda^3/3+\lambda t/2)} Y(\lambda, t).$$

Similarly, if we set $\kappa_2 = -1$ and $\kappa_1 = \theta - 1$, then from proposition 4.3 we note that the function $W(\mu, t)$ is related to the function $\psi(\mu, t)$ in system (4.9) via the following change of variables:

$$\psi(\mu, t) = \mu^{-1+\theta/2} W(\mu, t).$$

Finally, $\psi(\mu, t)$ is related to the function $\phi(\lambda, t)$ in (4.3) via the integral transform given in (4.6) and a simple gauge transformation:

$$\begin{pmatrix} 0 & \mu \\ 1 & 0 \end{pmatrix} \psi(\mu, t) = \int_L e^{\lambda\mu} \phi(\lambda, t) d\lambda.$$

We then find

$$\begin{pmatrix} 0 & \mu \\ 1 & 0 \end{pmatrix} \mu^{-1+\theta/2} W(\mu, t) = \int_L e^{\lambda\mu} \begin{pmatrix} 0 & \frac{1}{2} \\ -u^{-1} & 0 \end{pmatrix} e^{-(\lambda^3/3+\lambda t/2)} Y(\lambda, t) d\lambda,$$

which simplifies to give (3.9). □

5. Alternate secondary linearization of the 2×2 Fuchs–Garnier pairs for P_2

In section 4, we introduced a novel 3×3 matrix Fuchs–Garnier pair for the second Painlevé equation P_2 ; see system (4.1). One of the principal advantages of this 3×3 matrix system was that it is linear with respect to the spectral parameter λ , i.e. it is a secondary linearization of P_2 . In this section, we present an alternate secondary linearization of the 2×2 matrix Fuchs–Garnier pairs for P_2 .

Proposition 5.1. *The degenerate linear system*

$$d\text{JKT}_2^1 : \begin{cases} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \frac{d\Phi}{d\lambda} = \begin{pmatrix} -(z+t) & \lambda z + (yz + \theta) & 2\lambda \\ 2y & z & 2 \\ \lambda & 0 & 1 \end{pmatrix} \Phi, \\ \frac{d\Phi}{dt} = \frac{1}{2} \begin{pmatrix} 0 & z & 2 \\ -2 & 2y & 0 \\ 0 & -\lambda z & -2\lambda \end{pmatrix} \Phi, \end{cases} \quad (5.1)$$

is reducible to the 2×2 matrix Fuchs–Garnier pair for P_2 of Jimbo–Miwa.

Proof. From the third row in the λ equation in (5.1) we have the following relation between elements of the function Φ :

$$\Phi_3 = \lambda\Phi_1.$$

Using this relation to eliminate Φ_3 from system (5.1) we find that the remaining two components satisfy the linear 2×2 matrix system given in (4.3) with $\theta = (1 + \kappa_1)$. It was shown in proposition 4.2 that this system is related to the JM_2 -pair via an elementary gauge transformation. □

System (5.1) can be mapped to the HTW-pair. This fact is proved in the following proposition.

Proposition 5.2. System (5.1) can be mapped to the HTW-pair (3.6) by the application of the generalized Laplace transform defined in (4.6).

Proof. System (5.1) is linear with respect to the spectral variable λ and so we can immediately apply the Laplace transform in (4.6). The resulting degenerate 3×3 matrix system is given by

$$dJKT_2^2 : \begin{cases} \begin{pmatrix} 0 & z & 2 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \frac{d\Psi}{d\mu} = \begin{pmatrix} -\mu + (z+t) & -(yz+\theta) & 0 \\ -2y & -\mu - z & -2 \\ 0 & 0 & -1 \end{pmatrix} \Psi, \\ \frac{d\Psi}{dt} = \frac{1}{2} \begin{pmatrix} 0 & z & 2 \\ -2 & 2y & 0 \\ \mu - (z+t) & yz+\theta & 0 \end{pmatrix} \Psi. \end{cases} \quad (5.2)$$

In order to simplify the following calculation, we make a gauge transformation in system (5.2) of the form

$$\Psi = \begin{pmatrix} 0 & 0 & 1 \\ z^{-1} & z^{-1} & 0 \\ 0 & -\frac{1}{2} & 0 \end{pmatrix} \chi. \quad (5.3)$$

The resulting system is given by

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \frac{d\chi}{d\mu} = \begin{pmatrix} -(y+\theta z^{-1}) & -(y+\theta z^{-1}) & -\mu + (z+t) \\ -\mu - z & -\mu & -2yz \\ 0 & \frac{1}{2} & 0 \end{pmatrix} \chi, \quad (5.4a)$$

$$\frac{d\chi}{dt} = \begin{pmatrix} 0 & 0 & \mu - 2z - t \\ -y - \theta z^{-1} & -y - \theta z^{-1} & -\mu + z + t \\ \frac{1}{2} & 0 & 0 \end{pmatrix} \chi. \quad (5.4b)$$

The second row in equation (5.4a) implies a relation between the elements of the function χ :

$$\chi_2 = -\chi_1 - \frac{1}{\mu}(z\chi_1 + 2yz\chi_3). \quad (5.5)$$

Using this relation to eliminate χ_2 from system (5.4) we find that the remaining two components satisfy the following linear 2×2 matrix system:

$$\begin{aligned} \frac{d\psi}{d\mu} &= \left(\mu \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & z+t \\ -\frac{1}{2} & 0 \end{pmatrix} + \frac{1}{\mu} \begin{pmatrix} yz+\theta & 2yz(y+\theta z^{-1}) \\ -\frac{1}{2}z & -yz \end{pmatrix} \right) \psi, \\ \frac{d\psi}{dt} &= \left(\mu \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & -2z-t \\ \frac{1}{2} & 0 \end{pmatrix} \right) \psi, \quad \psi = \begin{pmatrix} \chi_1 \\ \chi_2 \end{pmatrix}. \end{aligned} \quad (5.6)$$

Making the gauge transformation,

$$\psi = \begin{pmatrix} -1 & -2y \\ 0 & 1 \end{pmatrix} \mu^{\theta/2} W, \quad (5.7)$$

in system (5.6) we get system (3.6). □

5.1. Alternate proof of theorem 4.4

Proof. From proposition 5.1 we note that the degenerate system (5.1) is reducible to the JM₂-pair (3.1), while from proposition 5.2 the degenerate system (5.4) is reducible to the

HTW-pair (3.6). The function $\Phi(\lambda, t)$ in (5.1) is related to the function $\chi(\mu, t)$ in (5.4) via the following integral transform:

$$\begin{pmatrix} 0 & 0 & 1 \\ z^{-1} & z^{-1} & 0 \\ 0 & -\frac{1}{2} & 0 \end{pmatrix} \chi(\mu, t) = \int_L e^{\lambda\mu} \Phi(\lambda, t) d\lambda.$$

The first two components of this expression give

$$\begin{pmatrix} \chi_3(\mu, t) \\ z^{-1}(\chi_1(\mu, t) + \chi_2(\mu, t)) \end{pmatrix} = \int_L e^{\lambda\mu} \begin{pmatrix} \Phi_1(\lambda, t) \\ \Phi_2(\lambda, t) \end{pmatrix} d\lambda.$$

Using relation (5.5) to eliminate χ_2 from this expression, we find

$$\begin{pmatrix} 0 & 1 \\ -\frac{1}{\mu} & -\frac{2y}{\mu} \end{pmatrix} \psi(\mu, t) = \int_L e^{\lambda\mu} \phi(\lambda, t) d\lambda,$$

where $\psi = (\chi_1, \chi_3)^T$ and $\phi = (\Phi_1, \Phi_2)^T$. From proposition 5.1, the function $Y(\lambda, t)$ is related to the function $\phi(\lambda, t)$ via the gauge transformation:

$$\phi(\lambda, t) = \begin{pmatrix} 0 & \frac{1}{2} \\ -u^{-1} & 0 \end{pmatrix} e^{-(\lambda^3/3+\lambda t/2)} Y(\lambda, t).$$

Similarly, from proposition 5.2, the function $W(\mu, t)$ is related to the function $\psi(\lambda, t)$ in system (5.6) via the gauge transform given in (5.7). Combining these two expressions we find

$$\begin{pmatrix} 0 & 1 \\ -\frac{1}{\mu} & -\frac{2y}{\mu} \end{pmatrix} \begin{pmatrix} -1 & -2y \\ 0 & 1 \end{pmatrix} \mu^{\theta/2} W(\mu, t) = \int_L e^{\lambda\mu} \begin{pmatrix} 0 & \frac{1}{2} \\ -u^{-1} & 0 \end{pmatrix} e^{-(\lambda^3/3+\lambda t/2)} Y(\lambda, t) d\lambda,$$

which simplifies to give (3.9). □

Appendix A. On the matrix versions of the Garnier pair for P_2

There are two matrix versions of the original scalar G_2 -pair: the JM_2 -pair and the Fuchs–Garnier pair obtained by Conte and Musette [1], the CM_2 -pair. Since both pairs were obtained by some simple transformations from the G_2 -pair they are equivalent and should be related via a simple gauge transformation. However, in [9] there are no explicit details given of the relation between the JM_2 - and G_2 -pairs and so, to complete our diagram in figure 2, we give this relation here. In the work [1], one finds details of the derivation of the matrix CM_2 -pair from the scalar G_2 -pair; however, it looks very much different from the matrix JM_2 -pair. To establish their equivalence we present a direct transformation between the JM_2 - and CM_2 -pairs avoiding the original ‘intermediate’ object G_2 -pair.

We begin with the relation between the JM_2 - and G_2 -pairs. Consider any column of the fundamental matrix solution to system (3.1): $Y_k = (Y_{k1}, Y_{k2})^T$, $k = 1, 2$, then the function $V = V(\lambda, t)$ defined as $Y_{k1} = \sqrt{u(\lambda - y)}V$, for any k , satisfies the original Garnier pair for P_2 :

$$G_2 : \begin{cases} \frac{d^2 V}{d\lambda^2} = \left(\frac{3}{4(\lambda - y)^2} - \frac{y'}{\lambda - y} + (y')^2 + \lambda^4 - y^4 + t(\lambda^2 - y^2) + 2\alpha(\lambda - y) \right) V, \\ \frac{dV}{dt} = \frac{1}{2(\lambda - y)} \frac{dV}{d\lambda} + \frac{1}{4(\lambda - y)^2} V, \end{cases} \quad (A.1)$$

where $\alpha = \frac{1}{2} - \theta$.

Thus we have shown that the JM₂-pair gives a matrix representation of the scalar G₂-pair. Of course, different matrix representations are also possible and, in particular, we discussed above the CM₂-pair. Clearly, all these matrix representations are equivalent; however, sometimes a direct explicit mapping between them is not immediately obvious. The CM₂-pair reads

$$\text{CM}_2 : \begin{cases} \frac{dM}{d\lambda} = \left(\lambda^3 \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \lambda^2 \begin{pmatrix} 0 & y \\ 0 & 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 & y^2 + t \\ 1 & 0 \end{pmatrix} + \begin{pmatrix} -y' & y^3 + ty + 2\alpha \\ -y & y' \end{pmatrix} \right) M, \\ \frac{dM}{dt} = \left(\frac{\lambda^2}{2} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + \lambda \begin{pmatrix} 0 & y \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & (3y^2 + t)/2 \\ 1/2 & 0 \end{pmatrix} \right) M, \end{cases} \tag{A.2}$$

where $y = y(t)$ is any solution of P_2 (1.1) and $y' = dy/dt$.

To map this system directly into the JM₂-pair given in (3.1) we introduce the parameter θ and functions $z = z(t), u = u(t)$ as follows:

$$\theta = \frac{1}{2} - \alpha, \quad z = y' - y^2 - \frac{t}{2}, \quad \frac{du}{dt} = -yu.$$

Then a relation between the fundamental solutions of (3.1), $Y = Y(\lambda, t)$, and (A.2), $M = M(\lambda, t)$, is given by the following gauge transformation depending on the spectral parameter:

$$M(\lambda, t) = G(\lambda, t)Y(\lambda, t), \quad G(\lambda, t) = \begin{pmatrix} \lambda + y & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} u^{-1/2} & 0 \\ 0 & u^{1/2} \end{pmatrix}. \tag{A.3}$$

Appendix B. Contour of integration in the generalized Laplace transform

Here we explain how to define the contour of integration in equation (3.9). In an analogous way the reader can find the contour of integration in the Laplace transform for the Fuchs–Garnier pairs for P_1 considered in section 2 as well as contours of integration for the Laplace transforms of the Fuchs–Garnier pairs for some other Painlevé equations considered in our previous work [10].

We start with some remarks on notation. Let us denote the columns of the matrix X by $X^k = X^k, k = 1, 2$, i.e., $X = (X^1, X^2)$. We assume that $X(\lambda)$ is an integrable function of λ . The notation L in equation (3.9), as well as in the Laplace transform (2.3) considered in section 2, is understood to mean a pair of contours of integration for each column X^k :

$$L = (\mathcal{L}^1, \mathcal{L}^2), \quad \text{so that} \quad \int_L X(\lambda) d\lambda = \left(\int_{\mathcal{L}^1} X^1(\lambda) d\lambda, \int_{\mathcal{L}^2} X^2(\lambda) d\lambda \right). \tag{B.1}$$

For brevity we call L simply the contour of integration. Assume that, for $k = 1, 2$, the 2×2 matrices C_k and the 2×2 diagonal matrices D_k are independent of λ . Then for any two integrable 2×2 matrix functions $X_k(\lambda)$ we have the following identity:

$$\int_L (C_1 X_1(\lambda) D_1 + C_2 X_2(\lambda) D_2) d\lambda = C_1 \int_L X_1(\lambda) d\lambda D_1 + C_2 \int_L X_2(\lambda) d\lambda D_2.$$

Let us now commence our analysis with a discussion of some general issues related to the choice of the contour L in (3.9). On one hand, we need a closed contour of integration because in sections 4 and 5 we assume vanishing of the certain off-integral terms that appear due to the integration by parts. On the other hand, the integrand in (3.9) is an entire function of λ , thus the contour of integration cannot be closed in the finite domain of the complex λ plane

and should pass through the point at infinity, since we would like to get a nontrivial function $W(\mu, t)$. This brings us immediately to the issues of convergence and conditions on L which ensure that the integral does not vanish.

To cope with these two problems we have to consider in more detail the asymptotic behaviour of the fundamental solutions of the JM₂-pair. The main instruments for this are the canonical solutions $Y_n(\lambda, t)$, $n \in \mathbb{Z}$ of the JM₂-pair (3.1), which are defined (uniquely) by their asymptotic expansion as $\lambda \rightarrow \infty$:

$$Y_n(\lambda, t) \sim \left(I + \mathcal{O}\left(\frac{1}{\lambda}\right) \right) \exp((\lambda^3/3 + \lambda t/2 - \theta \log \lambda)\sigma_3), \tag{B.2}$$

where I is the identity matrix, and $\log \lambda = \log|\lambda| + i \arg \lambda$, in the corresponding sectors

$$S_n = \left\{ \lambda : \frac{\pi}{6} + \frac{\pi(n-2)}{3} < \arg \lambda < \frac{\pi}{6} + \frac{\pi n}{3} \right\}.$$

The canonical solutions are related to each other by the Stokes matrices, S_n :

$$Y_{n+1}(\lambda) = Y_n(\lambda)S_n, \quad S_{2k+1} = \begin{pmatrix} 1 & s_{2k+1} \\ 0 & 1 \end{pmatrix}, \quad S_{2k} = \begin{pmatrix} 1 & 0 \\ s_{2k} & 1 \end{pmatrix},$$

where the numbers $s_n \in \mathbb{C}$ are called the Stokes multipliers. Moreover,

$$Y_{n+6}(\lambda e^{2\pi i}) = Y_n(\lambda) e^{-2\pi i \theta \sigma_3}, \quad n \in \mathbb{Z}.$$

From the above definitions one can easily deduce that the Stokes matrices satisfy the so-called cyclic relation, although this is not important in the following. We now find the leading term of asymptotics of the integrand in (3.9) assuming that Y coincides with Y_n :

$$\hat{Y}_n \equiv e^{-\lambda^3/3 + (\mu-t/2)\lambda} Y_n(\lambda, t) \underset[\lambda \in S_n]{\lambda \rightarrow \infty} \sim \begin{pmatrix} \lambda^{-\theta} e^{\lambda\mu} & 0 \\ 0 & \lambda^\theta e^{\lambda\mu} e^{-2\lambda^3/3 - t\lambda} \end{pmatrix}. \tag{B.3}$$

In the above asymptotics we have two exponential functions, namely, $e^{\lambda\mu}$ and $e^{-2\lambda^3/3 - t\lambda}$. The contour of integration should be chosen such that both exponents should vanish as $|\lambda| \rightarrow \infty$. We make the ‘Laplace exponent’, $e^{\lambda\mu}$, small as $|\lambda| \rightarrow \infty$ by demanding that $\text{Re } \lambda\mu < 0$. Since the contour has two directions as $|\lambda| \rightarrow \infty$ we get two conditions on μ . To satisfy both conditions we must have the angle between these directions less than π . The directions themselves are chosen such that the second exponent, $e^{-2\lambda^3/3 - t\lambda}$, decays. It can be any directions inside of the following sectors: $S_{2k} \cap S_{2k+1}$, $k \in \mathbb{Z}$. To achieve a better convergence the contour can be chosen asymptotic to the rays $\{\lambda : \arg \lambda = 2\pi n/3, n \in \mathbb{Z}\}$.

We have to take care that integrating a fundamental solution $Y(\lambda, t)$ in equation (3.9) we arrive at some fundamental solution $W(\mu, t)$. This means that, between the asymptotic directions of our contour, an arch of circle centered at the origin and having a large radius should cross a Stokes ray where the corresponding column of the fundamental solution $Y(\lambda, t)$ is affected by the Stokes phenomenon. If this condition is not satisfied then the column vector will be exponentially vanishing on that circle (as the radius enlarges to infinity) and, by the Cauchy theorem, the integral of that column taken along the contour also vanishes. In fact, this condition actually means that the above-mentioned arch intersects *two* Stokes rays of the *matrix* solution. In the sector between these Stokes rays our column is unbounded as $|\lambda| \rightarrow \infty$, so that the Cauchy theorem does not apply. Strictly speaking, after we choose the contour we also have to prove that we obtain the fundamental solution $W(\mu, t)$ by, say, calculating its asymptotics as $\mu \rightarrow \infty$.⁸

⁸ This calculation should be obvious for the experienced reader, we omit it here.

Now we apply the above principle to construct the contour L in (3.9). Consider, for example, the canonical solution Y_{2k} , $k \in \mathbb{Z}$. The reader can check that defining contour $L = L_k$ with $\mathcal{L}_k^1 = \mathcal{L}_k^2 = \mathcal{L}_k$, where \mathcal{L}_k is defined as any smooth simple⁹ curve asymptotic to the rays R_{2k+1}^ε and R_{2k+2}^ε , where

$$\mathcal{R}_n^\varepsilon := \left\{ \lambda : \arg \lambda = -\frac{\pi}{6} + \frac{\pi n}{3} + (-1)^n \varepsilon \right\}, \quad n \in \mathbb{Z} \quad \text{and} \quad 0 < \varepsilon < \pi/3,$$

fits all the conditions indicated in the above paragraphs, provided that the following condition is imposed on μ :

$$\frac{\pi}{3} - \frac{2\pi k}{3} + \varepsilon < \arg \mu < \pi - \frac{2\pi k}{3} - \varepsilon. \tag{B.4}$$

This condition comes from imposing the exponential decay condition, $\pi/2 < \arg \lambda \mu < 3\pi/2$, for asymptotics of \hat{Y}_{2k} on the rays R_{2k+1}^ε and R_{2k+2}^ε . Note that the asymptotics on R_{2k+1}^ε is given by equation (B.3) for $n = 2k$, while on $\mathcal{R}_{2k+2}^\varepsilon$ the Stokes phenomenon dictates the following leading term of asymptotics for \hat{Y}_{2k} :

$$\hat{Y}_{2k} \underset{\lambda \rightarrow \infty}{\sim} \begin{pmatrix} \lambda^{-\theta} e^{\lambda \mu} & 0 \\ 0 & \lambda^\theta e^{\lambda \mu} e^{-2\lambda^3/3 - t\lambda} \end{pmatrix} \begin{pmatrix} 1 + s_{2k}s_{2k+1} & -s_{2k+1} \\ -s_{2k} & 1 \end{pmatrix}.$$

With this choice of the contour L in (3.9) we construct the function $W(\mu)$ for all values of μ except the rays, $\mu : \arg \mu \neq \pi(1 + 2k)/3, k \in \mathbb{Z}$. On the latter rays $W(\mu)$ can be obtained via the analytic continuation. We can, also, obtain $W(\mu)$ on these rays by a proper choice of the contour L . In the latter case, it splits into the two contours (B.1).

For example, assume $\arg \mu = \pi$. Construct the following solution $Y = (Y_2^2, Y_4^2)$, where Y_2^2 and Y_4^2 are the second columns of the canonical solutions Y_2 and Y_4 , respectively. The function Y is a fundamental solution of system (3.1), iff $s_3 \neq 0$. Consider now the following anti-Stokes rays¹⁰:

$$\mathcal{R}_n := \{ \lambda : \arg \lambda_0 = 2n\pi/3 \}, \quad n = 0, 1, 2.$$

Now we can define the contour $L = (\mathcal{L}_1, \mathcal{L}_2)$ in (3.9) where for $k = 1, 2$, \mathcal{L}_k is any smooth simple⁹ curve with two asymptotic rays R_k and R_0 .

References

- [1] Conte R and Musette M 2000 Towards second order Lax pairs to discrete Painlevé equations of first degree *Chaos Solitons Fractals* **11** 41–52
- [2] Fabry M E 1885 Sur les intégrales des equations différentielles linéaires à coefficient rationnels *Thèse, Paris* 103
- [3] Flaschka H and Newell A C 1980 Monodromy- and spectrum-preserving deformations I *Commun. Math. Phys.* **76** 65–116
- [4] Fokas A S, Mugaň U and Zhou X 1992 On the solvability of Painlevé I, III and V *Inverse Problems* **8** 757–85
- [5] Garnier R 1912 Sur les équations différentielles du troisième ordre dont l'intégrale générale est uniforme et sur une classe d'équations nouvelles d'ordre supérieur dont l'intégrale générale a ses points critique fixes *Ann. Sci. Ec. Norm. Super.* **29** 1–126
- [6] Harnad J, Tracy C A and Widom H 1993 Hamiltonian structure of equations appearing in random matrices *Low-Dimensional Topology and Quantum Field Theory (Cambridge, 1992) (NATO Adv. Sci. Inst. Ser. B Phys.)* vol 315 (New York: Plenum) pp 231–45
- [7] Harnad J 1994 Dual isomonodromic deformations and moment maps to loop algebras *Commun. Math. Phys.* **166** 337–65
- [8] Ince E L 1956 *Ordinary Differential Equations* (New York: Dover)

⁹ The absence of selfintersections is not actually important.

¹⁰ It is the rays that originally were called the Stokes rays.

- [9] Jimbo M and Miwa T 1981 Monodromy preserving deformation of linear ordinary differential equations with rational coefficients: II. *Phys. D* **2** 407–48
- [10] Joshi N, Kitaev A V and Treharne P A 2007 On the linearization of the Painlevé III-VI equations and reductions of the three-wave resonant system *J. Math. Phys.* **48** 103512
- [11] Kapaev A A and Hubert E 1999 A note on the Lax pairs for Painlevé equations *J. Phys. A: Math. Gen.* **32** 8145–56
- [12] Kitaev A V 2006 Quadratic transformations for the third and fifth Painlevé equations *J. Math. Sci.* **136** 3586–95
- [13] Noumi M 2004 Painlevé equations through symmetry *Translations of Mathematical Monographs* vol 223 (Providence, RI: American Mathematical Society)
- [14] Mazzocco M 2002 *Painlevé Sixth Equation as Isomonodromic Deformations Equation of An Irregular System (CRM Proc. Lect. Notes* vol 32) (Providence, RI: American Mathematical Society) pp 219–38
- [15] Noumi M and Yamada Y 1998 Affine Weyl groups, discrete dynamical systems and Painlevé equations *Commun. Math. Phys.* **199** 281–95