

## PSU(2, 2|4) transformations of IIB superstring in $\text{AdS}_5 \times \text{S}^5$

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2009 J. Phys. A: Math. Theor. 42 095401

(<http://iopscience.iop.org/1751-8121/42/9/095401>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.157

The article was downloaded on 03/06/2010 at 08:39

Please note that [terms and conditions apply](#).

# PSU(2, 2|4) transformations of IIB superstring in $\text{AdS}_5 \times \text{S}^5$

Madoka Nishimura<sup>1</sup> and Yoshiaki Tani<sup>2</sup>

<sup>1</sup> Department of Community Service and Science, Tohoku University of Community Service and Science, Imoriyama 3–5–1, Sakata 998–8580, Japan

<sup>2</sup> Division of Material Science, Graduate School of Science and Engineering, Saitama University, Saitama 338-8570, Japan

E-mail: [tani@phy.saitama-u.ac.jp](mailto:tani@phy.saitama-u.ac.jp)

Received 6 November 2008

Published 4 February 2009

Online at [stacks.iop.org/JPhysA/42/095401](http://stacks.iop.org/JPhysA/42/095401)

## Abstract

The PSU(2, 2|4) transformation laws of the IIB superstring theory in the  $\text{AdS}_5 \times \text{S}^5$  background are explicitly obtained for the light-cone gauge in the Green–Schwarz formalism.

PACS numbers: 11.25.Tq, 11.25.Hf

## 1. Introduction

The construction and quantization of the superstring theory in anti de Sitter (AdS) spacetime have been an important subject since the original AdS/CFT correspondence [1–3] was proposed. Metsaev and Tseytlin [4] constructed the Green–Schwarz-type action of the type-IIB superstring in  $\text{AdS}_5 \times \text{S}^5$  as a sigma model with a coset target space  $\text{PSU}(2, 2|4)/[\text{SO}(4,1) \times \text{SO}(5)]$ . Then the light-cone gauge-fixing of  $\kappa$  transformations and reparametrizations on the worldsheet were discussed in [5, 6].

In this paper we discuss a global symmetry of the type-IIB superstring in  $\text{AdS}_5 \times \text{S}^5$  by using a group theoretical method. The symmetry is represented by the supergroup PSU(2, 2|4). We use the worldsheet action of [5], where the  $\kappa$  symmetry is fixed by the light-cone gauge. We obtain explicit forms of the transformation laws for the symmetry PSU(2, 2|4) in the light-cone gauge. The transformation laws we obtain will be useful in constructing the Noether charges for this symmetry [6]. They are also useful in finding consistent truncations of the theory, which are needed in some recent investigations of the gauge/string correspondence [7–10].

## 2. IIB superstring in $\text{AdS}_5 \times \text{S}^5$

The type-IIB superstring in  $\text{AdS}_5 \times \text{S}^5$  can be described [4] as a sigma model with a target space  $\text{PSU}(2, 2|4)/[\text{SO}(4,1) \times \text{SO}(5)]$ . The supergroup  $\text{PSU}(2, 2|4)$  contains a bosonic subgroup  $\text{SO}(4,2) \times \text{SO}(6)$ , which is the isometry of  $\text{AdS}_5 \times \text{S}^5$ . Its generators are

$$T^{\hat{I}} = P^a, J^{ab}, D, K^a, J^i{}_j, Q^{\pm i}, S^{\pm i}, \quad (1)$$

where  $P^a, J^{ab}, D, K^a$  are  $\text{SO}(4,2)$  generators,  $J^i{}_j$  are  $\text{SU}(4) \sim \text{SO}(6)$  generators and  $Q^{\pm i}, S^{\pm i}$  are fermionic generators. Here,  $a, b, \dots = 0, 1, 2, 3$  and  $i, j, \dots = 1, 2, 3, 4$  denote  $\text{SO}(3,1)$  and  $\text{SU}(4)$  indices. The (anti-)commutation relations of these generators are given in [5], whose conventions we use throughout this paper. The generators of the subalgebra  $\text{SO}(4,1) \times \text{SO}(5)$  are

$$J^{ab}, \quad \hat{J}^{4a} = K^a + \frac{1}{2}P^a, \quad J^{A'B'} = -\frac{1}{2}(\gamma^{A'B'})^j{}_i J^i{}_j, \quad (2)$$

where  $A', B' = 1, 2, 3, 4, 5$  are  $\text{SO}(5)$  indices and  $\gamma^{A'}$  are  $\text{SO}(5)$  gamma matrices. We use the light-cone coordinates  $x^{\pm} = \frac{1}{\sqrt{2}}(x^3 \pm x^0)$ ,  $x = \frac{1}{\sqrt{2}}(x^1 + ix^2)$ ,  $\bar{x} = \frac{1}{\sqrt{2}}(x^1 - ix^2)$  and define  $P = P^x, \bar{P} = P^{\bar{x}}, K = K^x, \bar{K} = K^{\bar{x}}$ .

We choose a representative of the coset space  $\text{PSU}(2, 2|4)/[\text{SO}(4,1) \times \text{SO}(5)]$  as

$$G = \exp(x^a P^a) \exp(\theta^{-i} Q_i^+ + \theta_i^- Q^{+i} + \theta^{+i} Q_i^- + \theta_i^+ Q^{-i}) \\ \times \exp(\eta^{-i} S_i^+ + \eta_i^- S^{+i} + \eta^{+i} S_i^- + \eta_i^+ S^{-i}) \exp(\phi D) \exp\left(\frac{1}{2}iy^{A'}(\gamma^{A'})^j{}_i J^j{}_i\right), \quad (3)$$

where  $\theta_i^{\pm} = (\theta^{\pm i})^\dagger, \eta_i^{\pm} = (\eta^{\pm i})^\dagger, Q_i^{\pm} = (Q^{\pm i})^\dagger, S_i^{\pm} = (S^{\pm i})^\dagger$ . The variables  $x^a, \phi, y^{A'}, \theta^{\pm i}, \eta^{\pm i}$  are coordinates of the coset space. We then fix the  $\kappa$  symmetry by the light-cone gauge condition [5],

$$\theta^{+i} = \eta^{+i} = 0 \quad (4)$$

and put  $\theta^{-i} = \theta^i, \eta^{-i} = \eta^i$  for simplicity. The left-invariant Cartan one-forms  $L^{\hat{I}}$  are defined by

$$G^{-1} dG = L^{\hat{I}} T^{\hat{I}} \\ = L_P^a P^a + \frac{1}{2}L^{ab} J^{ab} + L_D D + L_K^a K^a + L^j{}_i J^i{}_j + L_Q^{-i} Q_i^+ + L_{Q_i}^- Q^{+i} \\ + L_Q^{+i} Q_i^- + L_{Q_i}^+ Q^{-i} + L_S^{-i} S_i^+ + L_{S_i}^- S^{+i} + L_S^{+i} S_i^- + L_{S_i}^+ S^{-i}. \quad (5)$$

Using the explicit forms of the Cartan one-forms the worldsheet action in the light-cone gauge was obtained in [5].

## 3. $\text{PSU}(2, 2|4)$ transformations

According to the general theory of the nonlinear realization [11, 12] the  $\text{PSU}(2, 2|4)$  transformation of the representative (3) is

$$G \rightarrow G' = g G h^{-1}(g), \quad (6)$$

where  $g$  is an arbitrary element of  $\text{PSU}(2, 2|4)$  and  $h(g)$  is a compensating  $\text{SO}(4,1) \times \text{SO}(5)$  transformation which is chosen such that  $G'$  has a form in equation (3). After the light-cone gauge fixing of the  $\kappa$  symmetry (4) we also need a compensating  $\kappa$  transformation. An infinitesimal  $\text{PSU}(2, 2|4)$  transformation is thus written as

$$G^{-1} \delta G = G^{-1} \epsilon G - \sigma(\epsilon) + G^{-1} \delta_\kappa G, \quad (7)$$

where  $\epsilon$  is an arbitrary element of the PSU(2, 2|4) algebra

$$\begin{aligned} \epsilon = & \xi^a P^a + \frac{1}{2} \lambda^{ab} J^{ab} + \Lambda D + \zeta^a K^a + v^j_i J^j_j + \epsilon^{-i} Q^+_i + \epsilon_i^- Q^{+i} \\ & + \epsilon^{+i} Q^-_i + \epsilon_i^+ Q^{-i} + \beta^{-i} S^+_i + \beta_i^- S^{+i} + \beta^{+i} S^-_i + \beta_i^+ S^{-i} \end{aligned} \quad (8)$$

and  $\sigma(\epsilon)$  is a compensating SO(4,1)  $\times$  SO(5) transformation

$$\sigma(\epsilon) = \frac{1}{2} \tilde{\lambda}^{ab} J^{ab} + \tilde{\xi}^a \hat{J}^{4a} + \frac{1}{2} \tilde{v}^{A'B'} J^{A'B'}. \quad (9)$$

The last term in equation (7) is a compensating  $\kappa$  transformation. The parameters  $\tilde{\lambda}^{ab}$ ,  $\tilde{\xi}^a$ ,  $\tilde{v}^{A'B'}$  and those of the  $\kappa$  transformation depend on  $\epsilon$ .

The general  $\kappa$  transformation has a form [4],

$$\begin{aligned} G^{-1} \delta_\kappa G = & \tilde{\kappa}_Q^{-i} Q^+_i + \tilde{\kappa}_Q^- Q^{+i} + \tilde{\kappa}_Q^{+i} Q^-_i + \tilde{\kappa}_Q^+ Q^{-i} \\ & + \tilde{\kappa}_S^{-i} S^+_i + \tilde{\kappa}_S^- S^{+i} + \tilde{\kappa}_S^{+i} S^-_i + \tilde{\kappa}_S^+ S^{-i} + (J^{ab}, J^{A'B'} \text{ terms}). \end{aligned} \quad (10)$$

The coefficients in the present convention are given by

$$\begin{aligned} \tilde{\kappa}_Q^{+i} = & 2i[\hat{L}_\mu^+ \kappa_S^{\mu-i} + \hat{L}_\mu^{\bar{x}} \kappa_S^{\mu+i} + i\hat{L}_\mu^4 \kappa_Q^{\mu+i} - L_\mu^{A'} (\gamma^{A'})^i_j \kappa_Q^{\mu+j}], \\ \tilde{\kappa}_Q^{-i} = & 2i[\hat{L}_\mu^- \kappa_S^{\mu+i} - \hat{L}_\mu^x \kappa_S^{\mu-i} + i\hat{L}_\mu^4 \kappa_Q^{\mu-i} - L_\mu^{A'} (\gamma^{A'})^i_j \kappa_Q^{\mu-j}], \\ \tilde{\kappa}_S^{+i} = & -2i[2\hat{L}_\mu^+ \kappa_Q^{\mu-i} + 2\hat{L}_\mu^x \kappa_Q^{\mu+i} + i\hat{L}_\mu^4 \kappa_S^{\mu+i} + L_\mu^{A'} (\gamma^{A'})^i_j \kappa_S^{\mu+j}], \\ \tilde{\kappa}_S^{-i} = & -2i[2\hat{L}_\mu^- \kappa_Q^{\mu+i} - 2\hat{L}_\mu^{\bar{x}} \kappa_Q^{\mu-i} + i\hat{L}_\mu^4 \kappa_S^{\mu-i} + L_\mu^{A'} (\gamma^{A'})^i_j \kappa_S^{\mu-j}], \end{aligned} \quad (11)$$

where  $\mu = 0, 1$  is a world index on the worldsheet and  $\kappa_Q^{\mu\pm i}$ ,  $\kappa_S^{\mu\pm i}$  on the right-hand sides are independent transformation parameters. The  $\hat{L}_\mu^a$ ,  $\hat{L}_\mu^4$ ,  $L_\mu^{A'}$  are the pullbacks of the following one-forms to the worldsheet

$$\hat{L}^a = L_p^a - \frac{1}{2} L_K^a, \quad \hat{L}^4 = -L_D, \quad L^{A'} = -\frac{1}{2} i(\gamma^{A'})^i_j L^j_i. \quad (12)$$

For a general variation of the variables  $X^M = (x^a, \phi, y^{A'}, \theta^i, \eta^i)$  the variation of  $G$  in equation (3) is given by

$$\begin{aligned} G^{-1} \delta G = & \delta X^M L_M^{\hat{I}} T^{\hat{I}} \\ = & e^\phi \delta x^+ P^- + e^\phi [\delta x^- - \frac{1}{2} i(\theta^i \delta \theta_i + \theta_i \delta \theta^i)] P^+ + e^\phi \delta x \bar{P} + e^\phi \delta \bar{x} P \\ & + e^{-\phi} [\frac{1}{4} (\eta^2)^2 \delta x^+ + \frac{1}{2} i(\eta^i \delta \eta_i + \eta_i \delta \eta^i)] K^+ + \delta \phi D \\ & + [(\delta U U^{-1})^i_j + i(\tilde{\eta}^i \tilde{\eta}_j - \frac{1}{4} \eta^2 \delta^i_j) \delta x^+] J^j_i + e^{\frac{1}{2} \phi} (\delta \tilde{\theta}^i + i\tilde{\eta}^i \delta x) Q^+_i \\ & + e^{\frac{1}{2} \phi} (\delta \tilde{\theta}_i - i\tilde{\eta}_i \delta \bar{x}) Q^{+i} - i e^{\frac{1}{2} \phi} \tilde{\eta}^i \delta x^+ Q^-_i + i e^{\frac{1}{2} \phi} \tilde{\eta}_i \delta x^+ Q^{-i} \\ & + e^{-\frac{1}{2} \phi} (\delta \tilde{\eta}^i + \frac{1}{2} i\eta^2 \tilde{\eta}^i \delta x^+) S^+_i + e^{-\frac{1}{2} \phi} (\delta \tilde{\eta}_i - \frac{1}{2} i\eta^2 \tilde{\eta}_i \delta x^+) S^{+i}, \end{aligned} \quad (13)$$

where we have used the explicit forms of the Cartan one-form in the light-cone gauge given in [5].  $U^i_j$  is the SU(4) matrix determined by the coordinates  $y^{A'}$

$$U = \cos \frac{|y|}{2} + i\gamma^{A'} n^{A'} \sin \frac{|y|}{2}, \quad (14)$$

where  $|y|^2 = y^{A'} y^{A'}$ ,  $n^{A'} = y^{A'} / |y|$  and  $\tilde{\theta}^i = U^i_j \theta^j$ ,  $\tilde{\theta}_i = \theta_j (U^{-1})^j_i$ , etc. The compensating transformations in equation (7) are chosen such that the total transformation (7) has this form.

We are now ready to obtain explicit forms of the PSU(2, 2|4) transformations. We first compute the first term in equation (7). Useful formulae for doing this are listed in appendix. Then, we choose compensating transformations in the second and third terms such that the total transformations take the form in equation (13). Comparing the results of these computations and equation (13) we obtain the PSU(2, 2|4) transformations of the variables  $X^M$ .

The transformations for  $P^a$ ,  $D$ ,  $J^{+-}$ ,  $J^{+x}$ ,  $J^{x\bar{x}}$ ,  $J^{A'B'}$  and  $Q^+$  do not need compensating  $\kappa$  transformations and are easy to obtain. They were already given in [6]. We give them here for completeness.

•  $P^a$  transformations:

$$\delta x^a = \xi^a, \quad \delta(\text{others}) = 0. \quad (15)$$

•  $D$  transformations:

$$\begin{aligned} \delta x^a &= -\Lambda x^a, & \delta\phi &= \Lambda, & (U^{-1}\delta U)^i_j &= 0, \\ \delta\theta^i &= -\frac{1}{2}\Lambda\theta^i, & \delta\eta^i &= \frac{1}{2}\Lambda\eta^i. \end{aligned} \quad (16)$$

•  $J^{+-}$  transformations:

$$\begin{aligned} \delta x^+ &= -\lambda^{-+}x^+, & \delta x^- &= \lambda^{-+}x^-, & \delta x &= \delta\phi = (U^{-1}\delta U)^i_j = 0, \\ \delta\theta^i &= \frac{1}{2}\lambda^{-+}\theta^i, & \delta\eta^i &= \frac{1}{2}\lambda^{-+}\eta^i. \end{aligned} \quad (17)$$

•  $J^{+x}$  and  $J^{x\bar{x}}$  transformations:

$$\delta x^- = \lambda^{-\bar{x}}x + \lambda^{-x}\bar{x}, \quad \delta x = -\lambda^{-x}x^+, \quad \delta(\text{others}) = 0. \quad (18)$$

•  $J^{x\bar{x}}$  transformations:

$$\delta x = -\lambda^{\bar{x}x}x, \quad \delta\theta^i = -\frac{1}{2}\lambda^{\bar{x}x}\theta^i, \quad \delta\eta^i = \frac{1}{2}\lambda^{\bar{x}x}\eta^i, \quad \delta(\text{others}) = 0. \quad (19)$$

•  $J^i_j$  transformations:

$$\begin{aligned} \delta x^a &= \delta\phi = 0, & \delta\theta^i &= -v^i_j\theta^j, & \delta\eta^i &= -v^i_j\eta^j, \\ (U^{-1}\delta U)^i_j &= v^i_j + \frac{1}{4}\tilde{v}^{A'B'}(U^{-1}\gamma^{A'B'}U)^i_j. \end{aligned} \quad (20)$$

•  $Q^{+i}$  and  $Q_i^+$  transformations:

$$\delta x^- = \frac{1}{2}i\epsilon_i^-\theta^i + \frac{1}{2}i\epsilon^{-i}\theta_i, \quad \delta\theta^i = \epsilon^{-i}, \quad \delta(\text{others}) = 0. \quad (21)$$

The transformations for  $K^+$  do not need a compensating  $\kappa$  transformation either.

•  $K^+$  transformations:

$$\begin{aligned} \delta x^a &= \zeta^-(x^+x^a - \frac{1}{2}x \cdot x\eta^{a+} - \frac{1}{2}e^{-2\phi}\eta^{a+}), & \delta\phi &= -\zeta^-x^+, \\ \delta\eta^i &= -\zeta^-x^+\eta^i, & \delta(\text{others}) &= 0. \end{aligned} \quad (22)$$

The compensating  $SO(5)$  transformation with the parameter  $\tilde{v}^{A'B'}$  in equation (20) is not yet fixed. We will determine it and obtain the  $PSU(2, 2|4)$  transformation of the independent variables  $y^{A'}$  in section 4.

Other transformations need compensating  $\kappa$  transformations and are more involved.

•  $J^{-x}$  and  $J^{-\bar{x}}$  transformations:

$$\begin{aligned} \delta x^+ &= \lambda^{+\bar{x}}x + \lambda^{+x}\bar{x}, \\ \delta x^- &= \frac{1}{4}i e^{-\frac{3}{2}\phi}(\eta^i\hat{\kappa}_{Si}^- + \eta_i\hat{\kappa}_S^{-i}) + \frac{1}{2}i e^{-\frac{1}{2}\phi}(\theta^i\hat{\kappa}_{Qi}^- + \theta_i\hat{\kappa}_Q^{-i}) - \frac{1}{4}i e^{-2\phi}(\lambda^{+x}\theta_i\eta^i + \lambda^{+\bar{x}}\theta^i\eta_i)\eta^2, \\ \delta x &= -\lambda^{+x}(x^- - \frac{1}{2}i\theta^2 + \frac{1}{4}i e^{-2\phi}\eta^2), \\ \delta\phi &= -\frac{1}{2}(\lambda^{+\bar{x}}\theta^i\eta_i - \lambda^{+x}\theta_i\eta^i), \\ (U^{-1}\delta U)^i_j &= \lambda^{+\bar{x}}\theta^i\eta_j + \lambda^{+x}\theta_j\eta^i - (\text{trace part}) + \frac{1}{4}\tilde{v}^{A'B'}(U^{-1}\gamma^{A'B'}U)^i_j, \\ \delta\theta^i &= e^{-\frac{1}{2}\phi}\hat{\kappa}_Q^{-i} - \frac{1}{4}\lambda^{+x}e^{-2\phi}\eta^2\eta^i, \\ \delta\eta^i &= e^{\frac{1}{2}\phi}\hat{\kappa}_S^{-i} + \frac{1}{2}\lambda^{+\bar{x}}\eta^2\theta^i + \lambda^{+x}\theta_j\eta^j\eta^i, \end{aligned} \quad (23)$$

where we have defined

$$\hat{\kappa}_Q^{\pm i} = (U^{-1})^i_j\tilde{\kappa}_Q^{\pm j}, \quad \hat{\kappa}_S^{\pm i} = (U^{-1})^i_j\tilde{\kappa}_S^{\pm j}. \quad (24)$$

To obtain the form (13) we need to choose the parameters of the  $\kappa$  transformation as

$$\hat{\kappa}_Q^{+i} = \lambda^{+\bar{x}} e^{\frac{1}{2}\phi}\theta^i, \quad \hat{\kappa}_S^{+i} = \lambda^{+x} e^{-\frac{1}{2}\phi}\eta^i. \quad (25)$$

As we will see in section 4  $\hat{\kappa}_Q^{-i}, \hat{\kappa}_S^{-i}$  in the transformation (23) are determined from these conditions. Similarly, we obtain other transformations and conditions on the parameters of the  $\kappa$  transformations as follows.

•  $K$  and  $\bar{K}$  transformations:

$$\begin{aligned} \delta x^+ &= (\bar{\zeta}x + \zeta\bar{x})x^+, \\ \delta x^- &= \frac{1}{4}i e^{-\frac{3}{2}\phi}(\eta^i \hat{\kappa}_{S_i}^- + \eta_i \hat{\kappa}_S^{-i}) + \frac{1}{2}i e^{-\frac{1}{2}\phi}(\theta^i \hat{\kappa}_{Q_i}^- + \theta_i \hat{\kappa}_Q^{-i}) + \bar{\zeta}x(x^- + \frac{1}{2}i\theta^2) + \zeta\bar{x}(x^- - \frac{1}{2}i\theta^2) \\ &\quad + \frac{1}{4}i e^{-2\phi}x^+(\bar{\zeta}\theta^i\eta_i - \zeta\theta_i\eta^i)\eta^2, \\ \delta x &= \bar{\zeta}x^2 - \zeta x^+(x^- - \frac{1}{2}i\theta^2) - \frac{1}{2}\zeta e^{-2\phi}(1 + \frac{1}{2}ix^+\eta^2), \end{aligned} \quad (26)$$

$$\begin{aligned} \delta\phi &= -(\bar{\zeta}x + \zeta\bar{x}) - \frac{1}{2}x^+(\bar{\zeta}\theta^i\eta_i - \zeta\theta_i\eta^i), \\ (U^{-1}\delta U)^i_j &= x^+(\bar{\zeta}\theta^i\eta_j + \zeta\theta_j\eta^i) - (\text{trace part}) + \frac{1}{4}\bar{v}^{A'B'}(U^{-1}\gamma^{A'B'}U)^i_j, \\ \delta\theta^i &= e^{-\frac{1}{2}\phi}\hat{\kappa}_Q^{-i} + (\bar{\zeta}x + \zeta\bar{x})\theta^i + \frac{1}{2}i\zeta e^{-2\phi}(1 + \frac{1}{2}ix^+\eta^2)\eta^i, \\ \delta\eta^i &= e^{\frac{1}{2}\phi}\hat{\kappa}_S^{-i} + i\bar{\zeta}(1 - \frac{1}{2}ix^+\eta^2)\theta^i - \bar{\zeta}x\eta^i + \zeta x^+\theta_j\eta^j\eta^i, \\ \hat{\kappa}_Q^{+i} &= e^{\frac{1}{2}\phi}\bar{\zeta}x^+\theta^i, \quad \hat{\kappa}_S^{+i} = e^{-\frac{1}{2}\phi}\zeta x^+\eta^i. \end{aligned} \quad (27)$$

•  $K^-$  transformations:

$$\begin{aligned} \delta x^+ &= \zeta^+(x^-x^+ - \frac{1}{2}x \cdot x - \frac{1}{2}e^{-2\phi}), \\ \delta x^- &= \zeta^+[\frac{1}{2}i e^{-\frac{1}{2}\phi}(\theta^i \hat{\kappa}_{Q_i}^- + \theta_i \hat{\kappa}_Q^{-i}) + \frac{1}{4}i e^{-\frac{3}{2}\phi}(\eta^i \hat{\kappa}_{S_i}^- + \eta_i \hat{\kappa}_S^{-i}) + (x^-)^2 - \frac{1}{4}(\theta^2)^2 \\ &\quad + \frac{1}{2}e^{-2\phi}\theta^i\eta_i\theta_j\eta^j + \frac{1}{4}i e^{-2\phi}\eta^2(x\theta_i\eta^i + \bar{x}\theta^i\eta_i) + \frac{1}{16}e^{-4\phi}(\eta^2)^2], \\ \delta x &= \zeta^+[(x^- - \frac{1}{2}i\theta^2)x + \frac{1}{2}e^{-2\phi}(\theta^i\eta_i + \frac{1}{2}ix\eta^2)], \\ \delta\phi &= \zeta^+(-x^- - \frac{1}{2}x\theta_i\eta^i + \frac{1}{2}\bar{x}\theta^i\eta_i), \\ (U^{-1}\delta U)^i_j &= -i\zeta^+(\theta^i\theta_j + ix\eta^i\theta_j - i\bar{x}\theta^i\eta_j - \frac{1}{2}e^{-2\phi}\eta^i\eta_j) - (\text{trace part}) \\ &\quad + \frac{1}{4}\bar{v}^{A'B'}(U^{-1}\gamma^{A'B'}U)^i_j, \\ \delta\theta^i &= \zeta^+[e^{-\frac{1}{2}\phi}\hat{\kappa}_Q^{-i} + (x^- - \frac{1}{2}i\theta^2)\theta^i - \frac{1}{2}i e^{-2\phi}(\theta^j\eta_j + \frac{1}{2}ix\eta^2)\eta^i], \\ \delta\eta^i &= \zeta^+[e^{\frac{1}{2}\phi}\hat{\kappa}_S^{-i} + i\theta_j\eta^j(\theta^i + ix\eta^i) - \frac{1}{2}\bar{x}\eta^2\theta^i + \frac{1}{4}i e^{-2\phi}\eta^2\eta^i], \\ \hat{\kappa}_Q^{+i} &= \zeta^+(-\bar{x}e^{\frac{1}{2}\phi}\theta^i + \frac{1}{2}i e^{-\frac{3}{2}\phi}\eta^i), \quad \hat{\kappa}_S^{+i} = \zeta^+e^{-\frac{1}{2}\phi}i(\theta^i + ix\eta^i). \end{aligned} \quad (28)$$

•  $Q^-$  and  $Q_i^-$  transformations:

$$\begin{aligned} \delta x^+ &= 0, \quad \delta x = i\epsilon_i^+\theta^i, \quad \delta\phi = -\frac{1}{2}(\epsilon_i^+\eta^i - \epsilon^{+i}\eta_i), \\ \delta x^- &= \frac{1}{2}i e^{-\frac{1}{2}\phi}(\theta^i \hat{\kappa}_{Q_i}^- + \theta_i \hat{\kappa}_Q^{-i}) + \frac{1}{4}i e^{-\frac{3}{2}\phi}(\eta^i \hat{\kappa}_{S_i}^- + \eta_i \hat{\kappa}_S^{-i}) + \frac{1}{8}i e^{-2\phi}\eta^2(\epsilon_i^+\eta^i + \epsilon^{+i}\eta_i), \\ (U^{-1}\delta U)^i_j &= -(\epsilon_j^+\eta^i + \epsilon^{+i}\eta_j) - (\text{trace part}) + \frac{1}{4}\bar{v}^{A'B'}(U^{-1}\gamma^{A'B'}U)^i_j, \end{aligned} \quad (30)$$

$$\begin{aligned} \delta\theta^i &= e^{-\frac{1}{2}\phi}\hat{\kappa}_Q^{-i}, \quad \delta\eta^i = e^{\frac{1}{2}\phi}\hat{\kappa}_S^{-i} - \epsilon^{+i}\eta^i - \frac{1}{2}\eta^2\epsilon^{+i}, \\ \hat{\kappa}_Q^{+i} &= -e^{\frac{1}{2}\phi}\epsilon^{+i}, \quad \hat{\kappa}_S^{+i} = 0. \end{aligned} \quad (31)$$

•  $S^+$  and  $S_i^+$  transformations:

$$\begin{aligned} \delta x^+ &= 0, \quad \delta x = x^+\beta_i^-\theta^i, \quad \delta\phi = \frac{1}{2}ix^+(\beta_i^-\eta^i + \beta^{-i}\eta_i), \\ \delta x^- &= \frac{1}{2}i\theta^i(e^{-\frac{1}{2}\phi}\hat{\kappa}_{Q_i}^- - i\bar{x}\beta_i^-) + \frac{1}{2}i\theta_i(e^{-\frac{1}{2}\phi}\hat{\kappa}_Q^{-i} + ix\beta^{-i}) \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{4}i e^{-2\phi} \eta^i [e^{\frac{1}{2}\phi} \hat{\kappa}_{S_i}^- - (1 - \frac{1}{2}ix^+ \eta^2) \beta_i^-] \\
& + \frac{1}{4}i e^{-2\phi} \eta_i [e^{\frac{1}{2}\phi} \hat{\kappa}_S^{-i} - (1 + \frac{1}{2}ix^+ \eta^2) \beta^{-i}], \quad (32) \\
(U^{-1}\delta U)^i_j &= ix^+ (\beta_j^- \eta^i - \beta^{-i} \eta_j) - (\text{trace part}) + \frac{1}{4} \tilde{v}^{A'B'} (U^{-1} \gamma^{A'B'} U)^i_j, \\
\delta\theta^i &= e^{-\frac{1}{2}\phi} \hat{\kappa}_Q^{-i} - ix \beta^{-i}, \\
\delta\eta^i &= e^{\frac{1}{2}\phi} \hat{\kappa}_S^{-i} + (1 - \frac{1}{2}ix^+ \eta^2) \beta^{-i} + ix^+ \beta_j^- \eta^j \eta^i, \\
\hat{\kappa}_Q^{+i} &= -ix^+ e^{\frac{1}{2}\phi} \beta^{-i}, \quad \hat{\kappa}_S^{+i} = 0. \quad (33)
\end{aligned}$$

•  $S^-$  and  $S_i^-$  transformations:

$$\begin{aligned}
\delta x^+ &= 0, \quad \delta x = x \beta_i^+ \theta^i + \frac{1}{2}i e^{-2\phi} \beta^{+i} \eta_i, \\
\delta x^- &= \frac{1}{4}i e^{-\frac{3}{2}\phi} (\eta^i \hat{\kappa}_{S_i}^- + \eta_i \hat{\kappa}_S^{-i}) + \frac{1}{2}i e^{-\frac{1}{2}\phi} (\theta^i \hat{\kappa}_Q^- + \theta_i \hat{\kappa}_Q^{-i}) \\
& + \frac{1}{4}i e^{-2\phi} [(\theta_j - \frac{1}{2}i\bar{x}\eta_j) \eta^j \beta^{+i} \eta_i - (\theta^j + \frac{1}{2}ix\eta^j) \eta_j \beta^+ \eta^i] \\
& + \frac{1}{2} (x^- - \frac{1}{2}i\theta^2) \beta^+ \theta^i - \frac{1}{2} (x^- + \frac{1}{2}i\theta^2) \beta^+ \theta_i, \quad (34) \\
\delta\phi &= -\frac{1}{2}\beta_i^+ (\theta^i - ix\eta^i) + \frac{1}{2}\beta^{+i} (\theta_i + i\bar{x}\eta_i), \\
(U^{-1}\delta U)^i_j &= \beta_j^+ (\theta^i + ix\eta^i) + \beta^{+i} (\theta_j - i\bar{x}\eta_j) - (\text{trace part}) + \frac{1}{4} \tilde{v}^{A'B'} (U^{-1} \gamma^{A'B'} U)^i_j, \\
\delta\theta^i &= e^{-\frac{1}{2}\phi} \hat{\kappa}_Q^{-i} + \beta_j^+ \theta^j \theta^i + i (x^- - \frac{1}{2}i\theta^2) \beta^{+i} + \frac{1}{2} e^{-2\phi} \beta^{+j} \eta_j \eta^i, \\
\delta\eta^i &= e^{\frac{1}{2}\phi} \hat{\kappa}_S^{-i} + \beta^+ \eta^j (\theta^i + ix\eta^i) - (\theta_j - \frac{1}{2}i\bar{x}\eta_j) \eta^j \beta^{+i}, \\
\hat{\kappa}_Q^{+i} &= -i\bar{x} e^{\frac{1}{2}\phi} \beta^{+i}, \quad \hat{\kappa}_S^{+i} = -e^{-\frac{1}{2}\phi} \beta^{+i}. \quad (35)
\end{aligned}$$

#### 4. Compensating $SO(5)$ and $\kappa$ transformations

Here we fix the compensating  $SO(5)$  and  $\kappa$  transformations left undetermined above. Let us first consider the  $SO(5)$  transformations. The  $PSU(2, 2|4)$  transformations of  $U$  obtained in equations (20), (23), (26), (28), (30), (32), (34) have a form

$$(U^{-1}\delta U)^i_j = v^i_j + \frac{1}{4} \tilde{v}^{A'B'} (U^{-1} \gamma^{A'B'} U)^i_j, \quad (36)$$

where  $v^i_j$  is a given function of the variables  $X^M$  and the transformation parameters, and  $\tilde{v}^{A'B'}$  represents a compensating  $SO(5)$  transformation. On the other hand, a variation of the independent variables  $y^{A'}$  in equation (14) gives

$$U^{-1}\delta U = \frac{1}{2}i \gamma^{A'} n^{A'} n^{B'} \delta y^{B'} + \frac{1}{2}i \frac{\sin|y|}{|y|} \gamma^{A'} (\delta^{A'B'} - n^{A'} n^{B'}) \delta y^{B'} + \frac{1}{|y|} \sin^2 \frac{|y|}{2} \gamma^{A'B'} n^{A'} \delta y^{B'}. \quad (37)$$

We choose the compensating  $SO(5)$  transformations such that equation (36) has the form (37). Decomposing  $v^i_j$  into equation (36) as

$$v^i_j = \frac{1}{2}i v^{A'} (\gamma^{A'})^i_j - \frac{1}{4} v^{A'B'} (\gamma^{A'B'})^i_j, \quad (38)$$

we find that  $\tilde{v}^{A'B'}$  and the  $PSU(2, 2|4)$  transformations of  $y^{A'}$  are given by

$$\begin{aligned}
\tilde{v}^{A'B'} &= v^{A'B'} + 2 \tan \frac{|y|}{2} n^{[A'} v^{B']}, \\
\delta y^{A'} &= v^{A'B'} y^{B'} + \left[ n^{A'} n^{B'} + \frac{|y|}{\tan|y|} (\delta^{A'B'} - n^{A'} n^{B'}) \right] v^{B'}. \quad (39)
\end{aligned}$$

Next, we shall obtain  $\hat{\kappa}_Q^{-i}, \hat{\kappa}_S^{-i}$  from the conditions on  $\hat{\kappa}_Q^{+i}, \hat{\kappa}_S^{+i}$  in equations (25), (27), (29), (31), (33), (35), which we write as

$$\hat{\kappa}_Q^{+i} = \tau_Q^i, \quad \hat{\kappa}_S^{+i} = \tau_S^i. \quad (40)$$

From equation (11) these conditions are satisfied if we choose the independent  $\kappa$  transformation parameters as

$$\kappa_S^{\mu-i} = -\frac{1}{4}i \frac{\tau_Q^i}{\hat{L}_{\mu^+}}, \quad \kappa_Q^{\mu-i} = \frac{1}{8}i \frac{\tau_S^i}{\hat{L}_{\mu^+}}, \quad \kappa_S^{\mu+i} = \kappa_Q^{\mu+i} = 0, \quad (41)$$

where  $\mu = +, -$  are indices of the worldsheet light-cone coordinates. Substituting these equations into  $\hat{\kappa}_Q^{-i}, \hat{\kappa}_S^{-i}$  in equation (11) we obtain

$$\begin{aligned} \hat{\kappa}_Q^{-i} &= -\frac{1}{2} \left( \frac{\partial_+ x}{\partial_+ x^+} + \frac{\partial_- x}{\partial_- x^+} \right) \tau_Q^i + \frac{1}{4} i e^{-\phi} \left( \frac{\partial_+ \phi}{\partial_+ x^+} + \frac{\partial_- \phi}{\partial_- x^+} \right) \tau_S^i \\ &\quad + \frac{1}{4} e^{-\phi} \left( \frac{L_+^{A'}}{\partial_+ x^+} + \frac{L_-^{A'}}{\partial_- x^+} \right) (\gamma^{A'})^i_j \tau_S^j, \\ \hat{\kappa}_S^{-i} &= -\frac{1}{2} \left( \frac{\partial_+ \bar{x}}{\partial_+ x^+} + \frac{\partial_- \bar{x}}{\partial_- x^+} \right) \tau_S^i + \frac{1}{2} i e^{-\phi} \left( \frac{\partial_+ \phi}{\partial_+ x^+} + \frac{\partial_- \phi}{\partial_- x^+} \right) \tau_Q^i \\ &\quad - \frac{1}{2} e^{-\phi} \left( n \frac{L_+^{A'}}{\partial_+ x^+} + \frac{L_-^{A'}}{\partial_- x^+} \right) (\gamma^{A'})^i_j \tau_Q^j, \end{aligned} \quad (42)$$

where we have used the explicit forms of equation (12) given in [5]

$$\begin{aligned} \hat{L}_{\mu^+} &= e^\phi \partial_\mu x^+, \quad \hat{L}_{\mu^-} = e^\phi \partial_\mu x, \quad \hat{L}_{\mu^4} = -\partial_\mu \phi, \\ L_{\mu^+}^{A'} &= -\frac{1}{2} i (\gamma^{A'})^j_i [(\partial_\mu U U^{-1})^i_j + i(\tilde{\eta}^i \tilde{\eta}_j - \frac{1}{4} \eta^2 \delta_j^i) \partial_\mu x^+]. \end{aligned} \quad (43)$$

Using these  $\hat{\kappa}^-$ 's in equations (23), (26), (28), (30), (32), (34) we obtain explicit transformation laws.

From equation (30) we see that the  $Q^-$  transformation of  $x^+$  vanishes. This means in particular that the commutator of two  $Q^-$  transformations is zero on  $x^+$ , which at first sight looks inconsistent with the PSU(2, 2|4) algebra

$$\{Q^{-i}, Q_j^-\} = iP^- \delta_j^i. \quad (44)$$

This apparent inconsistency can be resolved as follows. Since we have not fixed a gauge for reparametrizations on the worldsheet, the commutator algebra closes up to a reparametrization. From equations (30), (31), (42) the commutator of two  $Q^-$  transformations on  $x$ , which should vanish according to the PSU(2, 2|4) algebra (44), becomes

$$[\delta_{Q^-}(\epsilon_1^+), \delta_{Q^-}(\epsilon_2^+)]x = (\xi^+ \partial_+ + \xi^- \partial_-)x, \quad (45)$$

where

$$\xi^\pm = \frac{1}{2\partial_\pm x^+} i(\epsilon_{2i}^+ \epsilon_1^{+i} - \epsilon_{1i}^+ \epsilon_2^{+i}). \quad (46)$$

This is a reparametrization with the parameters  $\xi^\pm$ . As the reparametrization of  $x^+$  with these parameters is

$$(\xi^+ \partial_+ + \xi^- \partial_-)x^+ = i(\epsilon_{2i}^+ \epsilon_1^{+i} - \epsilon_{1i}^+ \epsilon_2^{+i}), \quad (47)$$

the commutator on  $x^+$  can be written as

$$[\delta_{Q^-}(\epsilon_1^+), \delta_{Q^-}(\epsilon_2^+)]x^+ = -i(\epsilon_{2i}^+ \epsilon_1^{+i} - \epsilon_{1i}^+ \epsilon_2^{+i}) + (\xi^- \partial_- + \xi^+ \partial_+)x^+. \quad (48)$$

The first term on the right-hand side is a  $P^-$  transformation of  $x^+$  expected from the PSU(2, 2|4) algebra (44). Thus, the algebra (44) is satisfied up to a reparametrization.



## Appendix

We summarize formulae useful in computing  $G^{-1} \in G$ . From the formula

$$e^A B e^{-A} = B + [A, B] + \frac{1}{2!}[A, [A, B]] + \frac{1}{3!}[A, [A, [A, B]]] + \dots \quad (\text{A.1})$$

we obtain the following identities.

$$\begin{aligned} e^{-x \cdot P} J^{ab} e^{x \cdot P} &= J^{ab} - x^a P^b + x^b P^a, \\ e^{-x \cdot P} D e^{x \cdot P} &= D - x \cdot P, \\ e^{-x \cdot P} K^a e^{x \cdot P} &= K^a - x^a D + x^b J^{ba} + x^a x \cdot P - \frac{1}{2} x \cdot x P^a, \\ e^{-x \cdot P} S^{+i} e^{x \cdot P} &= S^{+i} - i x^+ Q^{-i} + i \bar{x} Q^{+i}, \\ e^{-x \cdot P} S^{-i} e^{x \cdot P} &= S^{-i} - i x^- Q^{+i} - i x Q^{-i}, \\ e^{-\theta \cdot Q^+} J^{+-} e^{\theta \cdot Q^+} &= J^{+-} + \frac{1}{2} \theta \cdot Q^+, \\ e^{-\theta \cdot Q^+} J^{x\bar{x}} e^{\theta \cdot Q^+} &= J^{x\bar{x}} - \frac{1}{2} (\theta^i Q_i^+ - \theta_i Q^{+i}) - \frac{1}{2} i \theta^2 P^+, \\ e^{-\theta \cdot Q^+} J^{-x} e^{\theta \cdot Q^+} &= J^{-x} - \theta^i Q_i^- - \frac{1}{2} i \theta^2 P, \\ e^{-\theta \cdot Q^+} D e^{\theta \cdot Q^+} &= D - \frac{1}{2} \theta \cdot Q^+, \\ e^{-\theta \cdot Q^+} K e^{\theta \cdot Q^+} &= K + i \theta^i S_i^+ + \frac{1}{2} i \theta^2 J^{+x}, \\ e^{-\theta \cdot Q^+} K^- e^{\theta \cdot Q^+} &= K^- - i (\theta^i S_i^- - \theta_i S^{-i}) + \frac{1}{2} i \theta^2 J^{x\bar{x}} - i \theta^j \theta_j J^j_j \\ &\quad - \frac{1}{2} i \theta^2 (\theta^i Q_i^+ - \theta_i Q^{+i}) + \frac{1}{4} (\theta^2)^2 P^+, \\ e^{-\eta \cdot Q^+} J^i_j e^{\eta \cdot Q^+} &= J^i_j - \theta^i Q_j^+ + \theta_j Q^{+i} - i \theta^i \theta_j P^+ - (\text{trace part}), \\ e^{-\theta \cdot Q^+} Q^{+i} e^{\theta \cdot Q^+} &= Q^{+i} + i \theta^i P^+, \\ e^{-\theta \cdot Q^+} Q^{-i} e^{\theta \cdot Q^+} &= Q^{-i} + i \theta^i \bar{P}, \\ e^{-\theta \cdot Q^+} S^{+i} e^{\theta \cdot Q^+} &= S^{+i} - \theta^i J^{+x}, \\ e^{-\theta \cdot Q^+} S^{-i} e^{\theta \cdot Q^+} &= S^{-i} + \frac{1}{2} \theta^i (J^{+-} - J^{x\bar{x}} - D) + \theta^j J^j_j + \frac{1}{2} \theta^2 Q^{+i} + \theta^i \theta^j Q_j^+ + \frac{1}{2} i \theta^i \theta^2 P^+, \\ e^{-\eta \cdot S^+} P e^{\eta \cdot S^+} &= P - i \eta_i Q^{+i} + \frac{1}{2} i \eta^2 J^{+x}, \\ e^{-\eta \cdot S^+} P^- e^{\eta \cdot S^+} &= P^- - i (\eta^i Q_i^- - \eta_i Q^{-i}) + \frac{1}{2} i \eta^2 J^{x\bar{x}} + i \eta^i \eta_j J^j_j \\ &\quad + \frac{1}{2} i \eta^2 (\eta^i S_i^+ - \eta_i S^{+i}) + \frac{1}{4} (\eta^2)^2 K^+, \\ e^{-\eta \cdot S^+} D e^{\eta \cdot S^+} &= D + \frac{1}{2} \eta \cdot S^+, \\ e^{-\eta \cdot S^+} J^{+-} e^{\eta \cdot S^+} &= J^{+-} + \frac{1}{2} \eta \cdot S^+, \\ e^{-\eta \cdot S^+} J^{x\bar{x}} e^{\eta \cdot S^+} &= J^{x\bar{x}} + \frac{1}{2} (\eta^i S_i^+ - \eta_i S^{+i}) - \frac{1}{2} i \eta^2 K^+, \\ e^{-\eta \cdot S^+} J^{-x} e^{\eta \cdot S^+} &= J^{-x} - \eta_i S^{-i} - \frac{1}{2} i \eta^2 K, \\ e^{-\eta \cdot S^+} J^i_j e^{\eta \cdot S^+} &= J^i_j - \eta^i S_j^+ + \eta_j S^{+i} + i \eta^i \eta_j K^+ - (\text{trace part}), \\ e^{-\eta \cdot S^+} Q^{+i} e^{\eta \cdot S^+} &= Q^{+i} + \eta^i J^{+x}, \\ e^{-\eta \cdot S^+} Q^{-i} e^{\eta \cdot S^+} &= Q^{-i} - \frac{1}{2} \eta^i (J^{+-} + J^{x\bar{x}} + D) - \eta^j J^j_j - \eta^i \eta^j S_j^+ - \frac{1}{2} \eta^2 S^{+i} + \frac{1}{2} i \eta^i \eta^2 K^+, \\ e^{-\eta \cdot S^+} S^{+i} e^{\eta \cdot S^+} &= S^{+i} - i \eta^i K^+, \\ e^{-\eta \cdot S^+} S^{-i} e^{\eta \cdot S^+} &= S^{-i} - i \eta^i K. \end{aligned}$$

## References

- [1] Maldacena J 1998 The large  $N$  limit of superconformal field theories and supergravity *Adv. Theor. Math. Phys.* **2** 231 (arXiv:[hep-th/9711200](#))
- [2] Gubser S S, Klebanov I R and Polyakov A M 1998 Gauge theory correlators from non-critical string theory *Phys. Lett. B* **428** 105 (arXiv:[hep-th/9802109](#))
- [3] Witten E 1998 Anti de Sitter space and holography *Adv. Theor. Math. Phys.* **2** 253 (arXiv:[hep-th/9802150](#))
- [4] Metsaev R R and Tseytlin A A 1998 Type IIB superstring action in  $AdS_5 \times S^5$  background *Nucl. Phys. B* **533** 109 (arXiv:[hep-th/9805028](#))
- [5] Metsaev R R and Tseytlin A A 2001 Superstring action in  $AdS_5 \times S^5$ :  $\kappa$ -symmetry light cone gauge *Phys. Rev. D* **63** 046002 (arXiv:[hep-th/0007036](#))
- [6] Metsaev R R, Thorn C B and Tseytlin A A 2001 Light-cone superstring in AdS space-time *Nucl. Phys. B* **596** 151 (arXiv:[hep-th/0009171](#))
- [7] Beisert N 2005 The dilatation operator of  $\mathcal{N} = 4$  super Yang-Mills theory and integrability *Phys. Rept.* **405** 1 (arXiv:[hep-th/0407277](#))
- [8] Tseytlin A A 2004 Semiclassical strings and AdS/CFT arXiv:[hep-th/0409296](#)
- [9] Alday L F, Arutyunov G and Tseytlin A A 2005 On integrability of classical superstrings in  $AdS_5 \times S^5$  *J. High Energy Phys.* **JHEP07(2005)002** (arXiv:[hep-th/0502240](#))
- [10] Alday L F, Arutyunov G and Frolov S 2006 New integrable system of 2dim fermions from strings on  $AdS_5 \times S^5$  *J. High Energy Phys.* **JHEP01(2006)078** (arXiv:[hep-th/0508140](#))
- [11] Coleman S, Wess J and Zumino B 1969 Structure of phenomenological Lagrangians: 1 *Phys. Rev.* **177** 2239
- [12] Callan C G, Coleman S, Wess J and Zumino B 1969 Structure of phenomenological Lagrangians: 2 *Phys. Rev.* **177** 2247