

## A generalised Dirac-Kronig-Penney model

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1989 J. Phys.: Condens. Matter 1 109

(<http://iopscience.iop.org/0953-8984/1/1/009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.89

The article was downloaded on 10/05/2010 at 15:46

Please note that [terms and conditions apply](#).

## A generalised Dirac–Kronig–Penney model

F Domínguez-Adame

Departamento de Física de Materiales, Facultad de Ciencias Físicas,  
 Universidad Complutense de Madrid, 28040-Madrid, Spain

Received 25 April 1988, in final form 17 June 1988

**Abstract.** In the present work, a generalisation of the Dirac–Kronig–Penney model for polyatomic crystals has been developed. The crystal potential is taken as a periodic array of  $\delta$ -function potentials, with several  $\delta$ -functions of different strength in each unit cell. The dispersion law of Dirac electrons in such a potential is found in a closed form.

In a recent paper (Eldib *et al* 1987), the well known Kronig–Penney model has been generalised to be applicable for polyatomic crystals, containing  $M$  different atoms in the unit cell. Nevertheless, relativistic effects in the band structure have not been considered in the above-mentioned model. These effects can be important at high energies on the electrons in solids consisting of heavy atoms (for a review on relativistic electrons in one-dimensional systems, see Roy (1986)). Relativistic electronic states in diatomic (equally spaced atoms) crystals have been discussed in an earlier paper by Sen Gupta (1974). The aim of the present work is to generalise the Dirac–Kronig–Penney model for polyatomic crystals, taking into account relativistic effects.

A number of researchers have treated  $\delta$ -function potentials with the Dirac equation (Sutherland and Mattis (1981) and McKellar and Stephenson (1987), and references therein). It has also been noted that the solution obtained by solving the Dirac equation for a  $\delta$ -function potential, and the expression attained when considering the limiting case of a square well (or barrier) are different. Fairbairn *et al* (1973) argued that this disagreement is related to the Klein paradox. In the above-mentioned work, McKellar and Stephenson have shown a reasonable way to use  $\delta$ -function potentials with the one-dimensional Dirac equation. For electrostatic-type potentials which approach a  $\delta$ -function limit (located at  $x_0$ ), they found the following boundary condition for the two-component electron wavefunction (in the standard representation):

$$\psi(x_0^+) = \begin{bmatrix} \cos(A/\hbar c) & -i \sin(A/\hbar c) \\ -i \sin(A/\hbar c) & \cos(A/\hbar c) \end{bmatrix} \psi(x_0^-) \quad (1)$$

$A$  being the strength of the potential.

We shall use the boundary condition (1) to find the dispersion law of Dirac electrons under the action of the crystal potential

$$V_c(x) = \sum_{\mu=1}^M \sum_{n=-\infty}^{\infty} V_{\mu}(x - R_{\mu} - nL) \quad (2)$$

where the ‘atomic’ potentials are taken as  $V_\mu(x) \rightarrow A_\mu \delta(x)$ , with  $A_\mu > 0$ . The position of each ion in the unit cell is denoted by  $R_\mu$  and  $L$  is the lattice parameter. The Bloch theorem is as follows:

$$\psi(x) = \exp(ikx)\varphi(x) \quad (3)$$

$\varphi(x)$  being a two-component periodic function, with the same period as the crystal lattice. Therefore, it suffices to find  $\varphi(x)$  within the unit cell  $[0, L]$ . The function  $\varphi(x)$  satisfies the equation (Domínguez-Adame 1987)

$$[-i\hbar c \sigma_x (\partial_x + ik) + \sigma_z mc^2 + V_c(x) - E(k)]\varphi(x) = 0 \quad (4)$$

$\sigma_x$  and  $\sigma_z$  being the usual  $2 \times 2$  Pauli matrices. The solution of this equation is readily found

$$\varphi_\mu(x) = \exp(-ikx) \begin{bmatrix} \exp(i\eta x) & \exp(-i\eta x) \\ \xi^{-1} \exp(i\eta x) & -\xi^{-1} \exp(-i\eta x) \end{bmatrix} \mathbf{P}_\mu$$

$$R_\mu < x < R_{\mu+1} \quad \mu = 0, 1, \dots, M \quad (5)$$

with  $R_0 \equiv 0$  and  $R_{M+1} \equiv L$ . For simplicity, we have introduced the notation  $\eta\hbar c = (E^2 - m^2c^4)^{1/2}$  and  $\xi^2 = (E + mc^2)/(E - mc^2)$ .  $\mathbf{P}_\mu$  denotes here a two-component constant vector. After applying the boundary condition (1) at each  $R_\mu$ , we obtain

$$\mathbf{P}_{\mu+1} = \mathbf{D}_\mu(\eta)\mathbf{P}_\mu \quad \mu = 1, 2, \dots, M \quad (6)$$

where

$$\mathbf{D}_\mu(\eta) = \begin{bmatrix} \alpha_\mu(\eta) & \beta_\mu(\eta) \\ \beta_\mu^*(\eta) & \alpha_\mu^*(\eta) \end{bmatrix} \quad (7)$$

with

$$\alpha_\mu(\eta) = \cos(A_\mu/\hbar c) - i(E/\eta\hbar c) \sin(A_\mu/\hbar c)$$

$$\beta_\mu(\eta) = -i(mc^2/\eta\hbar c) \sin(A_\mu/\hbar c) \exp(-2i\eta R_\mu). \quad (8)$$

It is interesting to note that  $\det\|\mathbf{D}_\mu(\eta)\| = 1$ ; we shall use this property below. The periodicity of  $\varphi(x)$  leads to the condition

$$\mathbf{P}_{M+1} = \exp(ikL) \begin{bmatrix} \exp(-i\eta L) & 0 \\ 0 & \exp(i\eta L) \end{bmatrix} \mathbf{P}_1 \quad (9)$$

and by successive use of equation (6) we immediately infer that

$$\exp(ikL) \begin{bmatrix} \exp(-i\eta L) & 0 \\ 0 & \exp(i\eta L) \end{bmatrix} \mathbf{P}_1 = \mathbf{G}\mathbf{P}_1 \quad (10)$$

where we have defined  $\mathbf{G} = \mathbf{G}(\eta) = \mathbf{D}_M(\eta)\mathbf{D}_{M-1}(\eta) \dots \mathbf{D}_1(\eta)$ . This  $2 \times 2$  matrix has interesting properties, namely  $\det\|\mathbf{G}\| = 1$ ,  $G_{11}^* = G_{22}$  and  $G_{21}^* = G_{12}$ , as can be readily verified from the properties of  $\mathbf{D}_\mu(\eta)$ . The consistency of equation (10) and the afore-

mentioned properties of  $\mathbf{G}$  give us the required dispersion law of Dirac electrons in the crystal potential:

$$\cos(\kappa L) = \text{Re} G_{11} \cos(\eta L) - \text{Im} G_{11} \sin(\eta L) \quad (11)$$

where  $\text{Re } z$  and  $\text{Im } z$  indicate real and imaginary parts, respectively, of the complex number  $z$ . Note that we must evaluate only one element of the matrix  $\mathbf{G}$ . The solution of the dispersion relation  $E(\kappa)$  has to be found numerically, for any arbitrary value of  $M$ .

If only a few atoms are placed in the unit cell of the crystal, the dispersion law (11) can be written in a more simplified form. For monatomic crystals, we have  $G_{11} = \alpha(\eta)$  so that

$$\cos(\kappa L) = \cos(\eta L) \cos(A/\hbar c) + (E/\eta\hbar c) \sin(\eta L) \sin(A/\hbar c). \quad (12)$$

This expression has formerly been proposed by McKellar and Stephenson for the Dirac–Kronig–Penney model. It should be observed that, as  $c \rightarrow \infty$ , equation (12) approaches the non-relativistic Kronig–Penney result.

If we now place two different atoms in each unit cell, we have  $G_{11} = \alpha_1(\eta)\alpha_2(\eta) + \beta_1^*(\eta)\beta_2(\eta)$ . Therefore, the relativistic dispersion law for diatomic crystals is found to be

$$\begin{aligned} \cos(\kappa L) = & \cos(\eta L) \cos[(A_1 + A_2)/\hbar c] + (E/\eta\hbar c) \sin(\eta L) \sin[(A_1 + A_2)/\hbar c] \\ & + 2(mc^2/\eta\hbar c)^2 \sin(A_1/\hbar c) \sin(A_2/\hbar c) \\ & \times \sin[\eta(R_2 - R_1)] \sin[\eta(L - R_2 + R_1)] \end{aligned} \quad (13)$$

which reduces to equation (12) as  $A_2 = 0$ . The non-relativistic limit can be written as

$$\begin{aligned} \cos(\kappa L) = & \cos(\eta L) + [m(A_1 + A_2)/\eta\hbar^2] \sin(\eta L) \\ & + 2A_1A_2(m/\eta\hbar^2)^2 \sin[\eta(R_2 - R_1)] \sin[\eta(L - R_2 + R_1)]. \end{aligned} \quad (14)$$

This is the expression obtained by Eldib *et al* who solved directly the Schrödinger equation for the crystal potential (2).

In a similar way, the dispersion relation for triatomic crystals ( $M = 3$ ) is found to be

$$\begin{aligned} \cos(\kappa L) = & \cos(\eta L) \cos[(A_1 + A_2 + A_3)/\hbar c] + (E/\eta\hbar c) \sin(\eta L) \\ & \times \sin[(A_1 + A_2 + A_3)/\hbar c] + 2(mc^2/\eta\hbar c)^2 \{ \cos(A_1/\hbar c) \sin(A_2/\hbar c) \\ & \times \sin(A_3/\hbar c) \sin[\eta(R_3 - R_2)] \sin[\eta(L - R_3 + R_2)] + \sin(A_1/\hbar c) \\ & \times \cos(A_2/\hbar c) \sin(A_3/\hbar c) \sin[\eta(R_3 - R_1)] \sin[\eta(L - R_3 + R_1)] \\ & + \sin(A_1/\hbar c) \sin(A_2/\hbar c) \cos(A_3/\hbar c) \sin[\eta(R_2 - R_1)] \\ & \times \sin[\eta(L - R_2 + R_1)] + 2(E/\eta\hbar c) \sin(A_1/\hbar c) \sin(A_2/\hbar c) \\ & \times \sin(A_3/\hbar c) \sin[\eta(R_3 - R_2)] \sin[\eta(R_2 - R_1)] \\ & \times \sin[\eta(L - R_3 + R_1)] \} \end{aligned} \quad (15)$$

which reduces again to the results of Eldib *et al* in the non-relativistic limit.

In summary, we have generalised the Dirac–Kronig–Penney model for polyatomic crystals containing  $M$  atoms in each unit cell. The treatment given above may be regarded only as a simple but very instructive way to study relativistic effects in polyatomic crystals.

**Acknowledgment**

The author thanks Dr N de Diego for a critical reading of the manuscript.

**References**

- Domínguez-Adame F 1987 *Am. J. Phys.* **55** 1003  
Eldib A M, Hassan H F and Mohamed M A 1987 *J. Phys. C: Solid State Phys.* **20** 3011  
Fairbairn W M, Glasser M L and Steslicka M 1973 *Surf. Sci.* **36** 462  
McKellar B H J and Stephenson G J 1987 *Phys. Rev. A* **36** 2566; 1987 *Phys. Rev. C* **35** 2262  
Roy C L 1986 *Proc. 4th Int. School Condensed Matter Physics (Varna, Bulgaria)* p 130  
Sen Gupta N D 1974 *Phys. Status Solidi b* **65** 351  
Sutherland B and Mattis D C 1981 *Phys. Rev. A* **24** 1194