

The two-channel Kondo model as a generalized one-dimensional inverse square long-range Haldane-Shastry spin model

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2000 J. Phys.: Condens. Matter 12 L217

(<http://iopscience.iop.org/0953-8984/12/12/102>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.218

The article was downloaded on 15/05/2010 at 20:31

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

The two-channel Kondo model as a generalized one-dimensional inverse square long-range Haldane–Shastry spin model

Guang-Ming Zhang[†] and Lu Yu^{‡§}[†]Center for Advanced Study, Tsinghua University, Beijing 100084, People's Republic of China[‡]International Center for Theoretical Physics, PO Box 586, Trieste 34100, Italy[§]Institute of Theoretical Physics, CAS, Beijing 100080, People's Republic of China

Received 5 January 2000, in final form 27 January 2000

Abstract. Majorana fermion representations of the algebra associated with spin, charge, and flavour currents have been used to transform the two-channel Kondo Hamiltonian. Using a path integral formulation, we derive a reduced effective action with long-range impurity spin–spin interactions at different imaginary times. In the semiclassical limit, it is equivalent to a one-dimensional Heisenberg spin chain with two-spin, three-spin, etc, long-range interactions, as a generalization of the inverse-square long-range Haldane–Shastry model. In this representation the elementary excitations are ‘semions’, and the non-Fermi liquid low energy properties of the two-channel Kondo model are recovered.

The two-channel Kondo model is known to have a non-Fermi liquid (non-FL) low energy fixed point in the overscreening case, and has been put forward as a model to explain non-FL behaviour observed in several quite different physical systems at low temperatures, such as certain heavy fermion alloys and two-level systems [1]. There are exact solutions for the ground state and thermodynamics of this model derived from the Bethe–Ansatz equations [2, 3], but there are still continuing efforts to find an intuitive understanding of the nature of excitations in the neighbourhood of the low energy fixed point. The numerical renormalization group [4], conformal field theory (CFT) [5], and the bosonization approach [6–8] made a number of predictions for many-body excitations at the fixed point, but they do not provide an intuitive interpretation of these excitations as is possible at the FL fixed point of the single channel Kondo model [9].

It was shown a long time ago that the single channel Kondo model can be reduced to an inverse-square one-dimensional *Ising* model [10], which is a prototype classical model in statistical physics [11, 12]. Moreover, such a reduced effective model helped Anderson and co-workers to establish the correct FL behaviour of the low energy fixed point for the one-channel model in the early 1970s [10, 13]; namely, the *Ising* spin–spin correlation function should behave as $1/\tau^2$, where τ is the imaginary time. On the other hand, the non-FL behaviour of the two-channel Kondo model is characterized by the dynamic correlation function of the impurity spin, $\langle T_\tau \vec{S}_d(\tau) \cdot \vec{S}_d(0) \rangle \sim 1/|\tau|$, but its physical meaning, unfortunately, has not been fully understood yet. To our knowledge, the possible connection of the overcrowd two-channel Kondo model with quantum spin models has not been explored so far.

In this Letter, the algebra of total spin, charge, and flavour currents of the two-channel Kondo model are represented in terms of Majorana fermions. In the path integral formulation we derive a reduced equivalent quantum Heisenberg spin model, in which the impurity spins at different imaginary times are strongly correlated, including two-spin, three-spin, etc, long-range exchange interactions. In particular, the two-spin interaction has the exact the same form as the integrable inverse-square one-dimensional Heisenberg spin chain—the so-called Haldane–Shastry (HS) model [14]—while the three-spin and four-spin interaction parts, etc, are natural generalizations of the HS model. The non-FL fixed point action of the two-channel Kondo model is identified with the two-spin interaction part (the HS model), and the elementary excitations of this low energy non-FL fixed point are spinons ($S = 1/2$ objects) obeying semion (half-fractional) statistics intermediate between those for bosons and fermions [15]. Actually, the two-spin long-range interaction part forms *an ideal semion gas*, while the three-spin long-range term is a *dangerous irrelevant interaction*, leading to important corrections to the thermodynamic properties. The fourth and higher order terms are irrelevant variables. The non-FL properties of the two-channel Kondo model are thus recovered.

We start with the Hamiltonian of the two-channel Kondo model in the form

$$\begin{aligned} H &= H_0 + H_I \\ H_0 &= \frac{v_f}{2\pi} \sum_{j=1}^2 \sum_{\sigma=\uparrow,\downarrow} \int_{-\infty}^{+\infty} dx \psi_{j,\sigma}^\dagger(x) (i\partial_x) \psi_{j,\sigma}(x) \\ H_I &= \sum_{a=x,y,z} J_a S_d^a J_s^a(0) \end{aligned} \quad (1)$$

where we have retained only the s-wave scattering, linearized the fermion spectrum, and replaced the incoming and outgoing waves with two left-moving electron fields $\psi_{j,\sigma}(x)$. $J_s^a(x)$ are the conduction electron spin current operators. $J_s^a(x) = \sum_{j,\sigma,\sigma'} : \psi_{j,\sigma}^\dagger(x) s_{\sigma,\sigma'}^a \psi_{j,\sigma'}(x) :$, where s^a are spin-1/2 matrices and $::$ means normal ordering. We introduce charge and flavour currents $J_c(x) = \sum_{j,\sigma} : \psi_{j,\sigma}^\dagger(x) \psi_{j,\sigma}(x) :$ and $J_f^a(x) = \sum_{j,j',\sigma} : \psi_{j,\sigma}^\dagger(x) t_{j,j'}^a \psi_{j',\sigma}(x) :$, where $t_{j,j'}^a$ are generators of an $SU(2)$ symmetry group. Following Affleck and Ludwig [5] the free part of the Hamiltonian can be rewritten as a sum of three commuting terms by the usual point splitting procedure (Sugawara construction):

$$H_0 = \frac{v_f}{2\pi} \int_{-\infty}^{+\infty} dx \left[\frac{1}{8} : J_c(x) J_c(x) : + \frac{1}{4} : \vec{J}_f(x) \cdot \vec{J}_f(x) : + \frac{1}{4} : \vec{J}_s(x) \cdot \vec{J}_s(x) : \right] \quad (2)$$

while the interaction term is expressed only in terms of the electron spin currents and the impurity spin. The information about the number of channels is contained in the commutation relations obeyed by the spin currents

$$[J_s^a(x), J_s^b(x')] = i\epsilon^{abc} J_s^c(x) \delta(x-x') + \frac{ki}{4\pi} \delta_{a,b} \delta'(x-x')$$

indicating that $J_s^a(x)$ form an $SU(2)$ level $k = 2$ Kac–Moody algebraic equation. Meanwhile, the charge and flavour currents satisfy

$$\begin{aligned} [J_c(x), J_c(x')] &= 2ki\delta'(x-x') \\ [J_f^a(x), J_f^b(x')] &= i\epsilon^{abc} J_f^c(x) \delta(x-x') + \frac{ki}{4\pi} \delta_{a,b} \delta'(x-x'). \end{aligned}$$

They form a $U(1)$ Kac–Moody and another $SU(k = 2)$ level-2 Kac–Moody algebraic equation, respectively.

It is now quite natural to introduce a Majorana representation of the spin current operators,

$$\begin{aligned} J_s^x(x) &= -i\chi_2(x)\chi_3(x) \\ J_s^y(x) &= -i\chi_3(x)\chi_1(x) \\ J_s^z(x) &= -i\chi_1(x)\chi_2(x) \end{aligned} \quad (3)$$

where $\chi_1(x)$, $\chi_2(x)$, and $\chi_3(x)$ are left-moving free Majorana fermion fields, and it can be shown to reproduce the SU(2) level-2 Kac–Moody commutation relations. It is important to note that this representation is only appropriate for the two-channel model, as it leads to a level-2 algebraic equation. It would be *inappropriate* for the single-channel Kondo model where the corresponding spin current generates a level-1 algebraic equation.

In a similar way, we can also introduce Majorana representations for the flavour currents

$$\begin{aligned} J_f^x(x) &= -i\chi_2'(x)\chi_3'(x) \\ J_f^y(x) &= -i\chi_3'(x)\chi_1'(x) \\ J_f^z(x) &= -i\chi_1'(x)\chi_2'(x) \end{aligned} \quad (4)$$

which reproduce the commutation relations satisfied by the flavour currents, and $J_c(x) = -2i\chi_4'(x)\chi_5'(x)$ representing the charge current operator. Note that χ_α' with $\alpha = 1, 2, 3, 4, 5$ are also left-moving free Majorana fermion fields. It is well known that the dynamics of charge, flavour and spin are completely determined by the commutation relations of these current operators. Though the spin currents of the two-channel Kondo model can be represented in terms of three Majorana fermion fields, $\chi_\alpha(x)$ ($\alpha = 1, 2, 3$), they cannot be given any simple physical interpretation in terms of the original conduction electrons $\psi_{j,\sigma}(x)$.

Now, using these current operators, the Hamiltonian (2) is presented as a quartic form in the Majorana fields. This form is convenient if one pursues a purely algebraic approach as in the CFT [5]. However, for our purposes it is more convenient to perform an *inverse* Sugawara construction, using again the point splitting procedure, and rewrite the terms quartic in the Majorana fermions as kinetic energy terms which are quadratic [16, 17]:

$$\begin{aligned} : J_c(x)J_c(x) : &:= 4 \sum_{\alpha=4}^5 \chi_\alpha'(i\partial_x)\chi_\alpha'(x) \\ : \vec{J}_f(x) \cdot \vec{J}_f(x) : &:= 2 \sum_{\alpha=1}^3 \chi_\alpha'(i\partial_x)\chi_\alpha'(x) \\ : \vec{J}_s(x) \cdot \vec{J}_s(x) : &:= 2 \sum_{\alpha=1}^3 \chi_\alpha(i\partial_x)\chi_\alpha(x). \end{aligned} \quad (5)$$

The model Hamiltonian is thus transformed into the following two parts [18]

$$\begin{aligned} H_c + H_f &= \frac{v_f}{4\pi} \sum_{\alpha=1}^5 \int_{-\infty}^{+\infty} dx \chi_\alpha'(x)(i\partial_x)\chi_\alpha'(x) \\ H_s &= \frac{v_f}{4\pi} \sum_{\alpha=1}^3 \int_{-\infty}^{+\infty} dx \chi_\alpha(x)(i\partial_x)\chi_\alpha(x) - \frac{iJ}{2} \int_{-\infty}^{+\infty} dx \delta(x) \vec{S}_d \cdot \vec{\chi}(x) \times \vec{\chi}(x). \end{aligned} \quad (6)$$

$H_c + H_f$ describes the non-interacting charge and flavour degrees of freedom. It is invariant under the symmetry group $U(1) \otimes SU(2)_2$ described by the five free Majorana fermion fields $\chi_\alpha'(x)$ ($\alpha = 1, 2, 3, 4, 5$). H_s is the main part of the model and it describes the spin degrees of freedom with three left-moving Majorana fermion fields χ_α ($\alpha = 1, 2, 3$) interacting with the impurity spin. It has the $SU(2)_2$ symmetry so that the full Hamiltonian is described by eight different Majorana fermion fields.

In the two-channel model Hamiltonian, H_s will give rise to the essential low energy physics of the model because it is the only part which includes the interaction. The reduced partition function of the model Hamiltonian with H_s can be written in the form of a functional integral:

$$Z = \oint D\widehat{\Omega} \int \prod_{\alpha=1}^3 D\chi_\alpha \exp \left\{ iS_d \omega(\widehat{\Omega}) - \int_0^\beta d\tau \left(\int_{-\infty}^{+\infty} dx \sum_{\alpha=1}^3 \frac{1}{2} \chi_\alpha \partial_\tau \chi_\alpha + H_s[\widehat{\Omega}, \chi_\alpha] \right) \right\} \quad (7)$$

where the impurity spin part has been expressed in terms of a spin coherent state path integral [19], $\widehat{\Omega} = (\theta, \phi)$ is a unit vector describing the family of spin states $|S_d, m_d\rangle$, the eigenstates of S_d^2 and S_d^z have eigenvalues $S_d(S_d+1)$ and m , respectively, and the periodic boundary condition is assumed for the spin vector variable.

$$iS_d \omega(\widehat{\Omega}) = iS_d \int_0^\beta d\tau (1 - \cos \theta) \dot{\phi} \quad (8)$$

is known as the Berry phase of the spin history, which is a purely geometric factor and will play no essential role here. In the path integral representation, the reduced Hamiltonian H_s is expressed as

$$H_s[\widehat{\Omega}, \chi_\alpha] = \frac{v_f}{4\pi} \sum_{\alpha=1}^3 \int_{-\infty}^{+\infty} dx \chi_\alpha(x, \tau) (i\partial_x) \chi_\alpha(x, \tau) - \frac{iJS_d}{2} \int_{-\infty}^{+\infty} dx \delta(x) \widehat{\Omega}(\tau) \cdot \vec{\chi}(x, \tau) \times \vec{\chi}(x, \tau) \quad (9)$$

where $\chi_\alpha(x, \tau)$ ($\alpha = 1, 2, 3$) are real Grassmann variables corresponding to the three Majorana fermion fields and $J\delta(x)\widehat{\Omega}(\tau)$ corresponds to a local non-obliant gauge field. As far as χ_α are concerned, the path integrals over them are bilinear. The partition function can thus be rewritten in the following form

$$Z = \oint D\widehat{\Omega} \exp(iS_d \omega(\widehat{\Omega})) \int \prod_{\alpha=1}^3 D\chi_\alpha \exp \left\{ -\frac{1}{2} \int_0^\beta d\tau \int_{-\infty}^{+\infty} dx \Psi^\dagger(x, \tau) \widehat{M} \Psi(x, \tau) \right\} \quad (10)$$

with a three-component vector $\Psi^\dagger(x, \tau) = (\chi_1(x, \tau), \chi_2(x, \tau), \chi_3(x, \tau))$ and its transposition $\Psi(x, \tau)$. The matrix is denoted by $\widehat{M} = (\partial_\tau - \bar{v}_f i\partial_x)I + iJS_d\delta(x)\widehat{M}'(\tau)$, where

$$\widehat{M}'(\tau) = \begin{bmatrix} 0 & -\Omega^z(\tau) & \Omega^y(\tau) \\ \Omega^z(\tau) & 0 & -\Omega^x(\tau) \\ -\Omega^y(\tau) & \Omega^x(\tau) & 0 \end{bmatrix} \quad (11)$$

$\bar{v}_f = v_f/2\pi$, and \widehat{I} is a 3×3 unit matrix. Then we integrate out the variables χ_α and obtain an effective action which only contains the spin vector variables:

$$Z = Z_0 \oint D\widehat{\Omega} \exp(iS_d \omega(\widehat{\Omega}) - S_{eff}) \quad (12)$$

$$S_{eff} = -\frac{1}{2} \text{Tr} \ln [1 + iJS_d\delta(x)\widehat{G}\widehat{M}'(\tau)]$$

where $Z_0 = \frac{1}{2} \det [(\partial_\tau - \bar{v}_f i\partial_x)\widehat{I}]$ is the partition function of the non-interacting limit of H_s , and its free Majorana fermion propagator is given by $\widehat{G} = (\partial_\tau - \bar{v}_f i\partial_x)^{-1}$. Tracing here is taken over space, imaginary time, and the matrix indices. Using the identity

$$\text{Tr} \ln(1 + \widehat{A}) = -\sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{Tr}(\widehat{A})^n$$

we obtain the general expression for the effective action

$$S_{eff} = \sum_{n=1}^{\infty} \frac{(-iJS_d)^n}{2n} \int_0^{\beta} d\tau_1 \dots \int_0^{\beta} d\tau_n G(\tau_{12})G(\tau_{23}) \dots \times G(\tau_{n1}) \text{Tr} [\widehat{M}'(\tau_1)\widehat{M}'(\tau_2) \dots \widehat{M}'(\tau_n)] \quad (13)$$

where the space integration has been easily carried out due to the presence of the delta function so that the free Majorana fermion propagators are replaced by the local one

$$G(\tau) = \frac{1}{\bar{v}_f} \frac{\pi/\beta}{\sin(\pi\tau/\beta)}$$

and τ_{ij} denotes $\tau_i - \tau_j$. Note that $\tau_1, \tau_2, \dots, \tau_n$ are *unequal* imaginary times. In fact $S_{eff}^{(n)}$ is the contribution of a one loop diagram made of n local propagators $G(\tau_{ij})$ and n local vertices $iJS_d\widehat{M}'(\tau_i)$. The factor $1/(2n)$ in front of each term is due to symmetry of the corresponding diagram. In the above derivation, the impurity spin is not specified to be $1/2$ and only the spin of the conduction electrons has been assumed to be $1/2$. However, the following discussion will focus on the overscreening case ($S_d = 1/2$).

Up to the second order of JS_d , the effective action is worked out as

$$S_{eff}^{(2)} = \frac{1}{2}(JS_d)^2 \int_0^{\beta} d\tau_i \int_0^{\beta} d\tau_j G^2(\tau_{ij})\widehat{\Omega}(\tau_i) \cdot \widehat{\Omega}(\tau_j) \quad (14)$$

which describes a Heisenberg spin chain with an inverse-square long-range *antiferromagnetic* interaction between the impurity spins at two different imaginary times. The sign of the original Kondo exchange coupling (ferromagnetic or antiferromagnetic) is not distinguishable here. Note that the spin coherent state path integral has assumed a periodic boundary condition for the spin variable so that the Heisenberg spins in fact sit on a circle of length β with exchange inversely proportional to the square of the distance between spins. This has the same form of the path integral functional as the one-dimensional Heisenberg spin chain with an inverse-square long-range interaction, the HS model [14] in the *semiclassical* limit. Therefore, all the *static* properties of the HS model can be readily translated to the present model.

- (a) The low energy states of $S_{eff}^{(2)}$ in the large- β limit are described by the chiral-SU(2) invariant $k = 1$ Wess–Zumino–Witten model, which is a conformally invariant Gaussian field theory. $S_{eff}^{(2)}$ thus can represent a fixed point action of the two-channel Kondo model, and the elementary excitations are spinons, i.e., the $S = 1/2$ particles instead of spin waves which are the elementary excitations of an ordered antiferromagnet. The spinons satisfy semion statistics intermediate between bosons and fermions, being an example of the *exclusion* statistics interpretation of fractional statistics [15].
- (b) $S_{eff}^{(2)}$ describes a free gas of spinons, being a fundamental model for gapless spin-1/2 antiferromagnetic spin model, and the dominant asymptotic spin–spin correlation function of $S_{eff}^{(2)}$ is algebraic with an universal exponent $\eta = 1$ without logarithmic corrections [15].

$$\langle T_i \widehat{\Omega}(\tau_i) \cdot \widehat{\Omega}(\tau_j) \rangle \sim \frac{1}{|\tau_i - \tau_j|} \quad (15)$$

and the impurity spin variable $\widehat{\Omega}(\tau)$ thus acquires a *dynamic* scaling dimension $1/2$. Such behaviour is also the universal spin–spin correlation function of the low energy non-FL behaviour of the two-channel Kondo model [5–8], leading to a marginal FL form [20] of the impurity spin spectrum: $\text{Im}\chi_d(\omega + i0^+) \sim \tanh(\frac{\omega}{2T})$.

- (c) We can thus conclude that $S_{eff}^{(2)}$ represents the low energy non-FL fixed point action of the spin part of the two-channel Kondo model.

The third order of JS_d can be viewed as a correction to the fixed point action, which has been derived as

$$S_{eff}^{(3)} = \frac{-i}{6} (JS_d)^3 \int_0^\beta d\tau_i \int_0^\beta d\tau_j \int_0^\beta d\tau_k G(\tau_{ij}) G(\tau_{jk}) G(\tau_{ki}) \widehat{\Omega}(\tau_i) \cdot (\widehat{\Omega}(\tau_j) \times \widehat{\Omega}(\tau_k)). \quad (16)$$

This term describes a Heisenberg spin chain with a long-range interaction of the impurity spins at three different imaginary times, and is completely antisymmetric in its indices (i, j, k) . In accordance with the low energy non-FL fixed point action $S_{eff}^{(2)}$, this interaction term is irrelevant because it has a dynamic scaling dimension $3/2$. However, it is this interaction that distinguishes the sign of the original Kondo exchange coupling, so that it is a *dangerous* irrelevant operator. It is conceivable that the non-FL thermodynamic properties of the two-channel Kondo model around the low energy fixed point are derived from a perturbation theory of $S_{eff}^{(3)}$. For instance, the second order perturbation calculation of $S_{eff}^{(3)}$ gives rise to the extra low temperature specific heat due to the presence of the impurity spin as $T \ln T$. Detailed calculations of the thermodynamic properties will be given in a future publication.

When the expansion is carried out to the fourth order, we obtain a four spin long-range interaction of the impurity spins at four different times

$$S_{eff}^{(4)} = \frac{1}{4} (JS_d)^4 \int_0^\beta d\tau_i \int_0^\beta d\tau_j \int_0^\beta d\tau_k \int_0^\beta d\tau_l \times G(\tau_{ij}) G(\tau_{jk}) G(\tau_{kl}) G(\tau_{li}) \widehat{\Omega}(\tau_i) \cdot \widehat{\Omega}(\tau_j) \widehat{\Omega}(\tau_k) \cdot \widehat{\Omega}(\tau_l) \quad (17)$$

which is clearly irrelevant as far as the low energy fixed point is concerned, as its dynamic scaling dimension is 2. All higher order terms thus contain no essential physics and can be neglected completely. Therefore, the reduced effective action will be given by $S_{eff} = S_{eff}^{(2)} + S_{eff}^{(3)}$, which is a natural generalization of the inverse-square long-range HS spin exchange model.

In summary, we use the Majorana fermion representation of the algebra of spin, charge, and flavour currents to transform the two-channel Kondo model. In a path integral formulation, we derive a reduced effective action of the two-channel Kondo model, which is a one-dimensional Heisenberg spin chains with two-spin, three-spin, etc, long-range interactions, as a natural generalization of the inverse-square long-range HS spin model. It is argued that the nontrivial two-channel Kondo physics in the low energy regime can be reproduced from the first two terms of the spin action, and the other relevant issues are under investigation. As pointed out in [21], the infinite set of multiplicative degeneracy of HS model [15] is due to the hidden SU(2) Yangian symmetry. Comparing our present formulation with the CFT treatment of the two-channel Kondo problem [5], this statement becomes apparent. After completing the present work, we became aware of a general review article [22] on exact results for highly correlated electron systems in one dimension, where some analogies of the inverse square long-range models to other interacting models are discussed within the framework of the Bethe–Ansatz equations.

One of the authors (G-M Zhang) would like to thank A C Hewson for his helpful discussions on the two-channel Kondo model.

References

- [1] Cox D L and Zawadowski A 1998 *Adv. Phys.* **47** 599
- [2] Andrei N and Destri C 1984 *Phys. Rev. Lett.* **52** 364
Tsvetlik A M and Wiegmann P B 1984 *J. Stat. Phys.* **38** 125
- [3] Schlottmann P and Sacramento P D 1993 *Adv. Phys.* **42** 641

- [4] Cragg D M, Lloyd P and Nozières Ph 1980 *J. Phys. C: Solid State Phys.* **13** 803
Pang H B and Cox D L 1991 *Phys. Rev. B* **44** 9454
- [5] Affleck I and Ludwig A W W 1991 *Nucl. Phys. B* **360** 641
Affleck I and Ludwig A W W 1993 *Phys. Rev. B* **48** 7297
Affleck I, Ludwig A W W, Pang H-B and Cox D L 1992 *Phys. Rev. B* **45** 7918
- [6] Emery V J and Kivelson S 1992 *Phys. Rev. B* **46** 10812
- [7] Maldacena J M and Ludwig A W W 1997 *Nucl. Phys. B* **506** 565
- [8] Coleman P, Ioffe L and Tselik A M 1995 *Phys. Rev. B* **52** 6611
- [9] Wilson K G 1974 *Collective Properties of Physical Systems (Nobel Symposia vol 24)* (New York: Academic)
Wilson K G 1975 *Rev. Mod. Phys.* **47** 773
- [10] Anderson P W and Yuval G 1971 *J. Phys. C: Solid State Phys.* **4** 607
- [11] Dyson F J 1969 *Commun. Math. Phys.* **12** 91
Dyson F J 1969 *Commun. Math. Phys.* **12** 215
Thouless D J 1969 *Phys. Rev.* **187** 732
Nagel J F and Bonner J C 1969 *J. Phys. C: Solid State Phys.* **3** 352
- [12] Cardy J L 1981 *J. Phys. A: Math. Gen.* **14** 1407
- [13] Anderson P W, Yuval G and Hamann D R 1970 *Phys. Rev. B* **1** 4464
- [14] Haldane F D M 1988 *Phys. Rev. Lett.* **60** 635
Shastry B S 1988 *Phys. Rev. Lett.* **60** 639
- [15] Haldane F D M 1991 *Phys. Rev. Lett.* **66** 1529
Haldane F D M 1991 *Phys. Rev. Lett.* **67** 937
- [16] Knizhnik V G and Zamolodchikov A B 1984 *Nucl. Phys. B* **247** 83
- [17] Ginsparg P 1990 *Fields, Strings and Critical Phenomena* ed Les Houches XLIX, E Brèzin and J Zinn-Justin
(Amsterdam: North Holland)
- [18] Zhang G-M, Hewson A C and Bulla R 1999 *Solid State Commun.* **112** 105
- [19] For the references of the spin coherent-states path integral see
Auerbach A 1994 *Interacting Electrons and Quantum Magnetism* (Springer)
- [20] Varma C M *et al* 1989 *Phys. Rev. Lett.* **63** 1996
- [21] Haldane F D M *et al* 1992 *Phys. Rev. Lett.* **69** 2021
- [22] Schlottmann P 1997 *Int. J. Mod. Phys. B* **11** 355