

On the analyticity of solutions in the dynamical mean-field theory

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2001 J. Phys.: Condens. Matter 13 9455

(<http://iopscience.iop.org/0953-8984/13/42/306>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.226

The article was downloaded on 16/05/2010 at 15:00

Please note that [terms and conditions apply](#).

On the analyticity of solutions in the dynamical mean-field theory

Th Pruschke^{1,4}, W Metzner² and D Vollhardt³

¹ Institut für Theoretische Physik, Universität Regensburg, 93040 Regensburg, Germany

² Institut für Theoretische Physik C, Technische Hochschule Aachen, 52056 Aachen, Germany

³ Theoretische Physik III, Elektronische Korrelationen und Magnetismus, Universität Augsburg, 86135 Augsburg, Germany

E-mail: thomas.pruschke@physik.uni-augsburg.de

Received 6 June 2001, in final form 15 August 2001

Published 5 October 2001

Online at stacks.iop.org/JPhysCM/13/9455

Abstract

The unphysical solutions of the periodic Anderson model obtained by Keiter and Leuders (Keiter H and Leuders T 2000 *Europhys. Lett.* **49** 801) in dynamical mean-field theory (DMFT) are shown to result from the authors' restricted choice of the functional form of the solution, leading to a violation of the analytic properties of the exact solution. By contrast, iterative solutions of the self-consistency condition within the DMFT obtained by techniques which preserve the correct analytic properties of the exact solution (e.g., quantum Monte Carlo simulations and the numerical renormalization group technique) always lead to physical solutions.

1. Introduction

In a recent paper, Keiter and Leuders [1] discussed the structure and the solutions of the self-consistency condition of the dynamical mean-field theory (DMFT) for the periodic Anderson model at $U = \infty$. They raised two independent points:

- (i) they showed analytically that on certain assumptions for the structure of the solution the self-consistency condition can have multiple solutions, which merge in the upper half of the complex plane, and
- (ii) they argued that calculations of $(1/d)$ -corrections (d : spatial dimension of the lattice) to the self-consistency condition of the DMFT are problematic, since the limit $d \rightarrow \infty$ does not commute with the limit $n \rightarrow \infty$ of the length n of loops embedded in the lattice.

⁴ Permanent address, as of 1 June 2001: Theoretische Physik III, Elektronische Korrelationen und Magnetismus, Universität Augsburg, 86135 Augsburg, Germany.

In the following, both points will be addressed. The main part of the paper will be concerned with a detailed analysis of the first issue. In particular, we will show that the conclusions of Keiter and Leuders [1] relating to this point are unwarranted, i.e., are due to the deficiency of the method used by these authors to solve (or simulate the solution of) the single-impurity Anderson model (SIAM).

2. Structure of the solution of the self-consistency condition

Multiple solutions of a non-linear set of equations are not unusual. In particular, most multiple solutions of non-linear equations for propagators are unphysical, i.e., have poles in the upper half-plane or lead to an incorrect asymptotic behaviour (or both). In this respect the finding of Keiter and Leuders [1], that under certain conditions there exists a branch point in the upper half-plane where an unphysical and the physical solution to the self-consistency condition *merge*, is disturbing. If true in general, it would raise doubts about the uniqueness of numerically obtained iterative solutions to the self-consistency condition, and would make the physical relevance of results obtained within the DMFT questionable.

Here we will show that, in fact, the self-consistency condition for the periodic Anderson model (PAM)⁵ *always leads to a unique physical result unless the functional space for the solutions is artificially restricted* as was done in reference [1]. In standard notation, the PAM reads

$$H_{\text{PAM}} = \sum_{k\sigma} \epsilon_k c_{k\sigma}^\dagger c_{k\sigma} + \sum_{i\sigma} \left(\epsilon_f + \frac{U}{2} f_{i\bar{\sigma}}^\dagger f_{i\bar{\sigma}} \right) f_{i\sigma}^\dagger f_{i\sigma} + V \sum_{i\sigma} (c_{i\sigma}^\dagger f_{i\sigma} + \text{h.c.}). \quad (1)$$

Within the DMFT, the non-trivial part of the partition function is represented by an effective local action [2]:

$$S_{\text{eff}} = - \sum_{\sigma} \int d\tau d\tau' f_{\sigma}^\dagger(\tau) \tilde{\mathcal{G}}^{-1}(\tau - \tau') f_{\sigma}(\tau') + U \int d\tau f_{\uparrow}^\dagger(\tau) f_{\uparrow}(\tau) f_{\downarrow}^\dagger(\tau) f_{\downarrow}(\tau) \quad (2)$$

where⁶

$$\tilde{\mathcal{G}}(z) = [z - \epsilon_f - V^2 \mathcal{G}(z)]^{-1} \quad (3)$$

and the auxiliary propagator $\mathcal{G}(z)$ describes an artificial system where the f states at the site under consideration have been removed. Note that the action (2) defines an effective SIAM with band-electron Green function $\mathcal{G}(z)$ which has to be determined self-consistently. For the PAM, equation (1), Keiter and Leuders derived the following expression [1]:

$$\mathcal{G}(z) = \frac{1}{N} \sum_k \frac{g_k(z)}{1 - g_k(z)T(z) + T(z)\mathcal{G}(z)}. \quad (4)$$

Here $g_k(z) = 1/(z - \epsilon_k)$ is the propagator of the non-interacting conduction band system and $T(z) = V^2 F(z)$, with $F(z)$ as the local f-electron Green function. The latter relation implies that $T(z)$ cannot be chosen at liberty. Equation (4) is not sufficient to solve the problem, since $F(z)$ is unknown. The quantity $F(z)$ has to be determined from an effective SIAM, where the propagator of the host electrons is given by $\mathcal{G}(z)$. It should be noted that the relation (4) is not only valid for $U = \infty$, but for *all* values of U , including $U = 0$. Moreover, it is important to mention that in reference [1] the artificial host propagator $\mathcal{G}(z)$ appearing in (3) is denoted by

⁵ Here we shall focus on the PAM in order to make direct contact with the discussion in reference [1]. However, the arguments presented are valid also for general one-band models.

⁶ The appearance of a momentum-independent hybridization V in (1) implies that the representation of $\tilde{\mathcal{G}}(z)$ is not unique. Therefore the form of $\tilde{\mathcal{G}}(z)$ used in reference [1] and here differs from the one used in reference [2], but can be derived from it using simple algebra.

$G(z)$, while conventionally, and also in this paper, $G(z)$ refers to the propagator of the *physical band states*.

Let us briefly comment on the validity of equation (4) for $U = 0$. At first glance it seems rather strange that the solution to a non-interacting system should be given by a complicated, non-linear self-consistency condition. The reason, however, is quite simple. First, equation (4) is set up in *real space*, while the solution to (1) for $U = 0$ is naturally obtained in reciprocal space. Second, as mentioned before, the quantity $\mathcal{G}(z)$ is not a physical propagator but an artificial auxiliary quantity. It is perfectly valid, although not necessarily reasonable, to rewrite the solution of the PAM (1) in terms of this auxiliary local propagator even for $U = 0$, thus obtaining the complicated expression (4). For the *physical* propagator $G(z)$, on the other hand, one eventually finds, inserting $T(z) = V^2[z - \epsilon_f - V^2\mathcal{G}(z)]^{-1}$, a simple and sensible solution (cf. equation (7) below). It must be noted that for $U = 0$ this procedure is exact, independently of dimension, provided that the hybridization in (1) is local. The DMFT or limit $d = \infty$ enters at a different stage, namely through the locality of $T(z)$ *even in the presence of the two-particle interaction* U .

To examine the analytical properties of (4) under the condition that $T(z)$ is given by the solution to an effective SIAM, some simple algebraic transformations have to be performed. Let us assume that $T(z)$ and $\mathcal{G}(z)$ are solutions to (4). Then we may rewrite (4) as

$$\mathcal{G}(z) = \frac{1}{1 + T(z)\mathcal{G}(z)} \frac{1}{N} \sum_k \frac{1}{z - \epsilon_k - T(z)/[1 + T(z)\mathcal{G}(z)]}. \tag{5}$$

For the SIAM, the host-electron propagator of the interacting system is obtained from the host propagator of the system without f states by using

$$G(z) = \mathcal{G}(z) + \mathcal{G}(z)T(z)\mathcal{G}(z). \tag{6}$$

We can rewrite (4) in terms of the physical host-electron propagator as

$$G(z) = \frac{1}{N} \sum_k \frac{1}{z - \epsilon_k - T(z)/[1 + T(z)\mathcal{G}(z)]} \tag{7}$$

and thereby identify the self-energy

$$\Sigma^{(c)}(z) = \frac{T(z)}{1 + T(z)\mathcal{G}(z)} \tag{8}$$

of the host electrons. Note that for $U = 0$ the self-energy (8) reduces to the well-known expression $\Sigma^{(c)}(z) = V^2/[z - \epsilon_f]$. Since the k -dependence in (7) is only due to ϵ_k , the k -sum can be converted to an energy integral:

$$\frac{1}{N} \sum_k \dots \rightarrow \int d\epsilon \rho^{(0)}(\epsilon) \dots \tag{9}$$

with a positive-definite density of states (DOS) $\rho^{(0)}(\epsilon)$. Relation (7) is simply a Hilbert transform; therefore one can make use of its analytical properties to show [3] that

$$G(z) = \frac{1}{z - \Sigma^{(c)}(z) - \Gamma(z - \Sigma^{(c)}(z))} \tag{10}$$

with a Herglotz-analytic function $\Gamma(\zeta)$, i.e.

$$\text{sgn}(\text{Im } \Gamma(\zeta)) = -\text{sgn}(\text{Im } \zeta)$$

and $\Gamma(\zeta) \sim 1/\zeta$ for $|\zeta| \rightarrow \infty$. Furthermore, $\Gamma(\zeta)$ is, in principle, a known function which is determined by the structure of the DOS alone. Finally, $\mathcal{G}(z)$ can be obtained from the definition of $G(z)^{-1} = \mathcal{G}(z)^{-1} - \Sigma^{(c)}(z)$ as

$$\mathcal{G}(z)^{-1} = G(z)^{-1} + \Sigma^{(c)}(z) = z - \Gamma\left(z - \frac{T(z)}{1 + T(z)\mathcal{G}(z)}\right). \tag{11}$$

It should be stressed that (11) is not merely a non-linear *equation* determining $\mathcal{G}(z)$ but, rather, a *functional equation*, since $T(z)$ is a functional of $\mathcal{G}(z)$. The functional dependence is determined by the solution of a non-trivial problem, namely the SIAM. It is thus not sufficient merely to study the possible roots of the non-linear equation (11); one also has to discuss the structure of this equation under the constraint that $T(z)$ must obey the basic properties of the solution to the SIAM (2). Since there does not exist an analytic solution to the SIAM in closed form, this constraint can be incorporated currently only within an iterative scheme, even if $\Gamma(\zeta)$ is known in closed form as, for example, for the Bethe lattice or the toy DOS used in [1].

Let us therefore assume that we knew the exact solution of the SIAM, and study the consequences of using an iterative procedure to solve the self-consistency condition. Starting in the n th step of this iterative procedure with a physical $\mathcal{G}^{(n)}(z)$ with $\text{sgn}(\text{Im } \mathcal{G}^{(n)}(z)) = -\text{sgn}(\text{Im } z)$ and $\mathcal{G}^{(n)}(z) \sim 1/z$ ($|z| \rightarrow \infty$), the corresponding propagator of the f states, $F^{(n)}(z)$, and hence $T^{(n)}(z)$, will have the same properties, i.e., the right-hand side of (11) will definitely have the correct asymptotics, because $T^{(n)}(z)\mathcal{G}^{(n)}(z) \sim 1/z^2$ and thus

$$\frac{T^{(n)}(z)}{1 + T^{(n)}(z)\mathcal{G}^{(n)}(z)} \sim 1/z$$

as $|z| \rightarrow \infty$. More importantly, however, the exact solution to the effective SIAM (2) has to satisfy the condition⁷

$$\text{sgn} [\text{Im}(T^{(n)}(z)^{-1} + \mathcal{G}^{(n)}(z))] = +\text{sgn}(\text{Im } z) \quad (12)$$

which is necessary to ensure that $\Sigma^{(c)}(z)$, equation (8), has the correct analytic properties, i.e., $\text{sgn}(\text{Im } \Sigma^{(c)}(z)) = -\text{sgn}(\text{Im } z)$. Under those conditions, the iterative procedure always stays within the subspace of physical solutions. Here it must be stressed once more that one is not free to *choose* $T^{(n)}(z) = V^2 F^{(n)}(z)$, or even choose a particular functional form for $T^{(n)}(z)$, since this form is entirely and uniquely determined by the solution of the effective SIAM with a fixed host Green function $\mathcal{G}^{(n)}(z)$. In particular, condition (12) turns out to be very susceptible to approximations and is easily violated. Such a violation would manifest itself in an indefinite sign of $\text{Im } \Sigma(z)$ and would thus lead to the crossing of a branch point into the upper half-plane. Note also that in the course of the iteration, the quantities entering on the right-hand side of (11) are those of the n th step, producing the host Green function $\mathcal{G}^{(n+1)}(z)$ for the $(n+1)$ th step and, unless self-consistency has been reached, $\mathcal{G}^{(n+1)}(z) \neq \mathcal{G}^{(n)}(z)$ in general⁸.

Hence we find that the branch point in the upper half-plane observed by Keiter and Leuders [1] is merely due to the authors' restricted choice of the functional form of $T(z)$ which leads to a violation of (12). That is, motivated by results for the standard SIAM, these authors *chose* for $T(z)$ the following *ansatz* [5]:

$$T(z) = V^2 \left(\frac{a_{\epsilon_f}}{\omega - \epsilon_f + i\Gamma_{\epsilon_f}} + \frac{a_K}{\omega + i\Gamma_K} \right). \quad (13)$$

This form represents the strong-coupling regime of the SIAM, i.e. $-\epsilon_f, \epsilon_f + U \gg \pi\rho^{(0)}(0)V^2$. The original single-particle level at ϵ_f is broadened by the coupling to the conduction bath. In addition, the sharp Abrikosov–Suhl resonance which characterizes the virtual bound state generated by spin-flip scattering appears at the Fermi energy. A typical result will have the form shown in figure 1(a). In figure 1 we assumed a Gaussian free DOS $\rho^{(0)}(\epsilon) = \exp(-\epsilon^2)/\sqrt{\pi}$ and set $V^2 = 0.2$, $\epsilon_f = -1$. Let us compare this result with a fully self-consistent DMFT calculation for the PAM with the same parameters [4]. The result is shown in figure 1(b) (full

⁷ This relation follows quite simply from the fact that $T(z) = V^2 F(z)$, where the local propagator $F(z)$ has the form $F(z) = [z - \epsilon_f - V^2\mathcal{G}(z) - \Sigma^{(f)}(z)]^{-1}$ with a Herglotz-analytic self-energy $\Sigma^{(f)}(z)$.

⁸ One consequence is that, for an iterative solution, equation (5) is not a valid reformulation of (11), because it only holds when self-consistency has been reached.




Figure 1. The imaginary part of $T(\omega + i\delta)$ for the PAM. (a) The result according to (13) as used in reference [1]; (b) the result from a full DMFT calculation with the NRG (numerical renormalization group) [4]. In the inset in (b) the two curves are compared for energies close to the Fermi energy.

line). It is obvious from the inset to figure 1(b) that the true solution of the PAM does not have the simple structure given by (13) assumed in reference [1]. In particular, one does not observe a single peak at the Fermi energy, but two peaks separated by a gap.

Consequently, the conclusion following equation (10) in reference [1] is unwarranted: the observed breakdown, i.e. the appearance of a branch point in the upper half-plane, does not result from the fact that in this regime physically interpretable single-valued solutions cannot be found, but merely from the deficiency of the method used to solve (or simulate the solutions of) the SIAM. In particular, one of the authors [6] and Anders [7] observed that approximations based on the resolvent perturbation theory [8], like the non-crossing approximation (NCA) or improved versions of it, always lead to spectra similar to (13) and are not able to reproduce more complicated features like the double-peak structure at the Fermi energy appearing in figure 1(b). It is therefore not surprising that the application of these techniques in solving the DMFT for the PAM leads to the breakdown described in reference [1]: it is a problem generated by the method used to solve the self-consistency condition and not of the self-consistency condition itself.

Our arguments do not imply that the self-consistency condition (11) must necessarily have only one unique solution for a given set of model parameters. Such a conclusion would require a more detailed knowledge of the solutions of the SIAM (in particular, whether there exists a one-to-one correspondence between $\mathcal{G}(z)$ and $F(z)$), and of the extent of the basin of attraction for a particular solution. Since there does not exist an exact analytical solution to the SIAM for an arbitrary host Green function $\mathcal{G}(z)$, statements about these properties of the self-consistency condition are out of reach. Indeed for the Hubbard model such a situation (namely, two physically valid, coexisting solutions) occurs close to the Mott–Hubbard metal–insulator transition [2].

3. $(1/d)$ -corrections to the DMFT

In DMFT a fermionic lattice model is mapped onto a SIAM, complemented by a self-consistency condition [2]. This mapping is rigorous for $d = \infty$, a fact which is not disputed in reference [1]. The possibilities and limitations of systematic $(1/d)$ -expansions around $d = \infty$ have already been explored by a number of authors [9–15]. Whether $(1/d)$ -corrections to DMFT are able to improve the theory depends on the quantity under investigation, on the type of $(1/d)$ -expansion, and also on the physics that one would like to describe. For example, while $(1/d)$ -corrections include non-local short-range correlations, they are unable to capture certain long-range correlations. Furthermore, $(1/d)$ -corrections, when truncated at finite order, may violate basic analytical properties of dynamical quantities [12]. All this is not unexpected. Indeed, analogous problems arise in $(1/d)$ -expansions around the Weiss mean-field theory for the Heisenberg model and the coherent potential approximation for the Anderson disorder model [10]. We do not see why the peculiarities of large loops in the large- d limit emphasized by Keiter and Leuders [1] should make expansions around DMFT particularly difficult or ambiguous. Since even the Green function of the *non-interacting* system can be expanded in loops, it is clear that these peculiarities are in one-to-one correspondence with the special behaviour of the *bare* DOS in the limit $d \rightarrow \infty$ (e.g., the absence of van Hove singularities) [16]⁹. Potential problems with the $(1/d)$ -expansion of the bare DOS can be avoided by inserting the actual DOS of the finite-dimensional system into the DMFT equations. Then only corrections to the *local approximation* of irreducible vertices underlying the DMFT need to be considered, as has been done already by several authors [9, 14, 17, 18].

4. Conclusions

In summary, we showed that the iterative solution of the self-consistency condition within the DMFT always leads to solutions that stay in the physical subspace. Therefore, the unphysical branch cuts observed by Keiter and Leuders [1] are due to the method used by these authors to solve the SIAM which violates fundamental analytic properties of the exact solution. Similar violations are caused by the NCA and other approximations based on the resolvent perturbation theory [8]. In fact, this also appears to be why it has not been possible up to now to obtain meaningful results for the PAM using the NCA and related methods, while techniques which preserve the correct analytic properties of the exact solution (e.g., iterated perturbation theory, quantum Monte Carlo simulations, and use of the numerical renormalization group) do not encounter similar difficulties.

⁹ Indeed, for the Bethe lattice, where self-avoiding loops do not exist, the large-loop peculiarities discussed in reference [1] and the unphysical band tails typical for hypercubic lattices for $d \rightarrow \infty$ are absent. The DMFT and (10), however, are well defined also for a Bethe lattice.

Acknowledgments

We thank F Anders, V Janiš, M Jarrell, H Keiter and D Otto for useful communication.

References

- [1] Keiter H and Leuders T 2000 *Europhys. Lett.* **49** 801
- [2] Georges A, Kotliar G, Krauth W and Rozenberg M 1996 *Rev. Mod. Phys.* **68** 13
- [3] Maier Th, Jarrell M, Pruschke Th and Keller J 2000 *Eur. Phys. J. B* **13** 613
- [4] Pruschke Th, Bulla R and Jarrell M 2000 *Phys. Rev. B* **61** 12 799
- [5] Leuders T 1998 *PhD Thesis* University of Dortmund
- [6] Pruschke T, unpublished
- [7] Anders F, private communication
- [8] Keiter H and Kimball J C 1970 *Phys. Rev. Lett.* **25** 672
- [9] Gebhard F 1990 *Phys. Rev. B* **41** 9452
- [10] Vlaming R and Vollhardt D 1992 *Phys. Rev. B* **45** 4637
- [11] Vlaming R, Uhrig G and Vollhardt D 1992 *J. Phys.: Condens. Matter* **4** 7773
Halvorsen E, Uhrig G S and Czycholl G 1994 *Z. Phys. B* **94** 291
Uhrig G S 1996 *Phys. Rev. B* **54** 10 436
- [12] van Dongen P J G 1994 *Phys. Rev. B* **50** 14 016
It follows immediately from (A9) therein that the spectral weight has an acausal part of order $\exp(-2/\sqrt{d})$:
van Dongen P J G private communication
Also see
van Dongen P J G and Schauerte T 2001 to be published
- [13] Schiller A and Ingersent K 1995 *Phys. Rev. Lett.* **75** 113
Zaránd G, Cox D L and Schiller A 2000 *Phys. Rev. B* **62** R16227
- [14] Hettler M, Tahvildar-Zadeh A N, Jarrell M, Pruschke Th and Krishnamurthy H R 1998 *Phys. Rev. B* **58** 7475
Hettler M H, Mukerjee M, Jarrell M and Krishnamurthy H R 2000 *Phys. Rev. B* **61** 12 739
- [15] Janiš V and Vollhardt D 2001 *Phys. Rev. B* **63** 125112
- [16] Müller-Hartmann E 1989 *Z. Phys. B* **74** 507
- [17] Metzner W 1989 *Z. Phys. B* **77** 253
- [18] Maier Th, Jarrell M, Pruschke Th and Keller J 2000 *Phys. Rev. Lett.* **85** 1524