

Driving a first order quantum phase transition by coupling a quantum dot to a 1D charge density wave

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2007 J. Phys.: Condens. Matter 19 086215

(<http://iopscience.iop.org/0953-8984/19/8/086215>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 129.252.86.83

The article was downloaded on 28/05/2010 at 16:19

Please note that [terms and conditions apply](#).

Driving a first order quantum phase transition by coupling a quantum dot to a 1D charge density wave

Y Weiss, M Goldstein and R Berkovits

The Minerva Center, Department of Physics, Bar-Ilan University, Ramat-Gan 52900, Israel

Received 20 October 2006, in final form 10 January 2007

Published 9 February 2007

Online at stacks.iop.org/JPhysCM/19/086215

Abstract

The ground state properties of a one-dimensional system with particle-hole symmetry, consisting of a gate controlled dot coupled to an interacting reservoir, are explored using the numerical DMRG method. It has previously been shown that the system's thermodynamic properties as a function of the gate voltage in the Luttinger liquid phase are qualitatively similar to the behaviour of a non-interacting wire with an effective (renormalized) dot-lead coupling. Here we examine the thermodynamic properties of the wire in the charge density wave phase, and show that these properties behave quite differently. The number of electrons in the system remains constant as a function of the gate voltage, while the total energy becomes linear. Moreover, by tuning the gate voltage on the dot in the charge density wave phase it is possible to drive the wire through a first order quantum phase transition in which the population of each site in the wire is inverted.

The properties of one-dimensional (1D) interacting systems have attracted much interest going back half a century [1, 2]. Much recent effort has concentrated on understanding the conductivity and I - V characteristics of a Luttinger liquid (LL) coupled to an impurity [3]. These properties are probed experimentally by measurements of the temperature dependent conduction through 1D systems [4], and tunnelling spectroscopy into 1D wires [5]. The measurements of [5] also indicate a localization transition in the wire for low densities, which might be associated with a charge density wave (CDW). Recently it has been realized that the occupation of an impurity level or a quantum dot coupled to an interacting 1D sample may be used as a probe of its properties [6, 7]. The occupation is experimentally accessible, for example by the charging effect of the impurity on a quantum point contact in its vicinity [8]. A generic model for such a situation is a dot coupled to a lead. If such a dot is controlled by a gate one can change its orbital energy. In this paper we shall demonstrate that by following the occupation of the dot as a function of the gate voltage it is possible to identify a transition in the wire from an LL to a CDW phase. In the CDW regime of the wire the occupation of the dot switches abruptly once the dot orbital crosses the Fermi energy. Moreover, by changing the

gate voltage of the quantum dot the wire is driven through a first order quantum phase transition (QPT), in which the ground state of the entire system switches.

The difference between measuring the conductivity through the dot–lead system or the local density of states at the impurity [9], and probing the dot occupation using e.g. a quantum point contact, must be emphasized. It is commonly known that the special behaviour of the LL spectrum near the Fermi energy causes the conductivity to vanish, even in the presence of a single impurity [10]. Thus measuring the conductivity of a system containing an LL lead and a dot is not expected to show any level broadening at zero temperature.

Nevertheless, connecting the dot to the LL lead, while measuring its occupation using a quantum point contact, results in a different situation. In this case the LL acts as a reservoir for the dot, and a level broadening can be seen. This difference results from the fact that such a level is coupled to all the lead states, even those far from the Fermi energy, and thus the special behaviour of the LL near E_F is not expected to dominate. It was indeed shown that the dot’s level width follows the regular Breit–Wigner line shape, even in the presence of interactions, although the strength of coupling between the dot and the 1D wire is renormalized as a function of the interactions in the wire [7].

In this paper we show that a qualitatively different behaviour of the dot’s population is exhibited once the wire is in the CDW phase. It is known that a one-dimensional system of spinless electrons with nearest neighbour repulsive interactions undergoes a QPT between an LL phase for weak interactions and a CDW phase for strong interactions [11–13]. The different behaviour of the dot population between the LL and the CDW phases stems from the difference between the ground states of the two phases in the wire. It is argued that, while the LL is only locally influenced by the coupling to the dot, a semi-infinite CDW coupled to the dot undergoes a QPT when the orbital’s level crosses the wire’s chemical potential. We show that a sharp jump in the dot’s population occurring at this point is accompanied by an abrupt switch of the ground state of the entire system. This switch is not a simple level crossing, but rather a true QPT, since it becomes a sharp transition only once the wire is semi-infinite [14]. Other hallmarks of a first order QPT, such as a discontinuity in the first derivative of the grand potential, as well as an inversion of the CDW order parameter as a result of the inversion of the wire’s occupation for each site, are seen.

The Hamiltonian describing a spinless fermionic system composed of a single orbital (‘dot’) coupled to a 1D nearest neighbour interacting wire (‘lead’) is given by $\hat{H} = \hat{H}_{\text{dot}} + \hat{H}_{\text{dot-lead}} + \hat{H}_{\text{lead}}$, where

$$\begin{aligned}\hat{H}_{\text{dot}} &= \epsilon_0 \hat{a}^\dagger \hat{a}, \\ \hat{H}_{\text{dot-lead}} &= -V(\hat{a}^\dagger \hat{c}_1 + \hat{c}_1^\dagger \hat{a}), \\ \hat{H}_{\text{lead}} &= -t \sum_{j=1}^{L-1} (\hat{c}_j^\dagger \hat{c}_{j+1} + \text{h.c.}) + I \sum_{j=1}^{L-1} (\hat{c}_j^\dagger \hat{c}_j - \frac{1}{2})(\hat{c}_{j+1}^\dagger \hat{c}_{j+1} - \frac{1}{2}).\end{aligned}\tag{1}$$

We denote the dot’s energy level by ϵ_0 , $V(t)$ is the dot–lead (lead) hopping matrix element, and I is the nearest-neighbour interaction strength in the lead. \hat{a}^\dagger (\hat{a}) is the creation (annihilation) operator of an electron in the dot, and \hat{c}_j^\dagger (\hat{c}_j) is the creation (annihilation) operator of an electron at site j in the lead. In the interaction term, a positive background is assumed. The lead hopping matrix element, t , is taken as unity, in order to set the energy scale.

In order not to increase the number of free parameters, the model discussed here does not contain a dot–lead interaction term. Nevertheless, we have checked that such a term does not influence qualitatively the results of this paper. We refer the reader to a previous work [7], in which it was explicitly shown that addition of such an interaction term influences only the effective position and width of the dot level.




Figure 1. The dot population n_{dot} as a function of the level energy ϵ_0 . For $I < 2$ (LL phase) the curves fit the non-interacting formula with an effective coupling constant V_{eff} (lines—DMRG results, symbols—fit to equation (2)), while for $I > 2$ (CDW phase) the width is zero. Exactly at the transition point ($I = 2$) n_{dot} is continuous but does not fit the non-interacting formula.

As was previously mentioned, while the lead is in the LL phase, the Breit–Wigner formula with an effective dot–lead coupling describes the dot population $n(\epsilon_0)$ quite well [7]. For a dot coupled to an LL lead in which the interaction parameter $g < 1/2$, a jump in the dot population was predicted by Furusaki and Matveev [6]. However, for the spinless model with nearest neighbour interaction the LL parameter is always in the range of $1/2 \leq g \leq 1$, and the Furusaki–Matveev jump is not expected.

The grand potential $\hat{\Omega} = \hat{H} - \mu \hat{N}$ (\hat{N} is the particle number operator) was diagonalized using a finite-size DMRG calculation [7, 15] for different values of V and I , with a lead of up to $L = 300$ sites. In order to obtain particle–hole symmetry, for which the LL to CDW transition was widely discussed [11–13], μ is set to zero. The total number of particles in the system is not fixed during the renormalization process, so that the results describe the experimental situation of a finite section of a 1D wire which is coupled to a dot and to an external electron reservoir. In particular, the total occupation obtained for such a system can be non-integral. As a function of the dot’s level (ϵ_0), the following ground state properties were calculated: the system’s grand potential Ω , the total number of electrons N and the dot population n_{dot} . The population of lead sites was also calculated in order to differentiate between local effects of the dot population and global phenomena in the lead.

We begin by comparing the behaviour of n_{dot} in the two phases of the interacting lead. For spinless fermions with nearest-neighbour interactions, it is known that the system experiences a phase transition (of a Kosterlitz–Thouless type [16]), between the LL and the CDW phases, at $I = 2t$. Indeed, the results presented in figure 1 show a qualitative difference between $n_{\text{dot}}(\epsilon_0)$ in these two regimes. In the LL phase the curves fit quite well the Breit–Wigner formula [7] with an effective coupling V_{eff}

$$n_{\text{dot}}(\mu, \epsilon_0) = \frac{1}{\pi} \int_{-2t}^{\mu} \frac{\frac{V_{\text{eff}}^2}{t} \sqrt{1 - \frac{\epsilon^2}{4t^2}}}{\frac{V_{\text{eff}}^4}{t^2} (1 - \frac{\epsilon^2}{4t^2}) + ((1 - \frac{V_{\text{eff}}^2}{2t^2})\epsilon - \epsilon_0)^2} d\epsilon, \quad (2)$$




Figure 2. The CDW order parameter P as a function of the interaction I for a 300-site lead (DMRG—symbols, theory—lines). For the CDW phase P is inverted between the cases $\epsilon_0 \rightarrow 0^+$ and $\epsilon_0 \rightarrow 0^-$, while in the LL phase both cases result in $P = 0$. Inset: P as a function of ϵ_0 for $I = 1.5$ (triangles), 2.5 (squares) and 3.5 (circles). The LL case results in a constant $P = 0$, while an inversion of P occurs at $\epsilon_0 = 0$ for the two CDW cases.

while in the CDW phase the width becomes zero, and a jump in n_{dot} occurs at $\epsilon_0 = 0$. This jump is associated with the degeneracy of the CDW ground state.

In order to support this argument, we first calculate the properties of the ground state for a lead not coupled to a dot, by taking $V = t$ and the limit $\epsilon_0 \rightarrow 0$. The CDW order parameter can be defined as [17]

$$P(i) = \frac{1}{2}(-1)^i[2n(i) - n(i - 1) - n(i + 1)], \quad (3)$$

where $n(i) = \langle \hat{c}_i^\dagger \hat{c}_i \rangle$ is the occupation of the i th lead site in the ground state. The value of $P(i)$ does not change much as a function of the spatial coordinate, except in the vicinity of the lead's edges. We thus define $P = P(L/2)$.

In figure 2 the dependence of P on the interaction strength I , as obtained by the DMRG method, is shown, and compared to the exact results [18]. For $I < 2$, the system is in the LL phase and indeed $P = 0 \pm 10^{-4}$. In this case the population of each lead site is $1/2$ and the lead is half filled; i.e. $N = L/2$.

As expected, the metal–insulator transition occurs at $I = 2$. For $I > 2$, in which the system is in the CDW phase, the value of P is finite. For values of I which are far from the transition point ($I > 2.5$), we get a very good fit to the theory. Near the transition point the numerics tend to emphasize the charge oscillations, resulting in too large values of $|P|$. This tendency, however, does not affect the following qualitative conclusions.

As can be seen, the CDW ground state is different for the cases $\epsilon_0 \rightarrow 0^+$ and $\epsilon_0 \rightarrow 0^-$, resulting in two values of P (i.e. $\pm|P|$) for any value of I in the CDW regime. For $\epsilon_0 = 0$ the ground state is twofold degenerate in the thermodynamic limit.

In the CDW case, special care should be devoted to the number of electrons in the system, and to the difference between even and odd lead length L . For $\epsilon_0 \rightarrow 0$ and $V = t$, the lead length is effectively $L + 1$. Denoting the ground state for $\epsilon_0 \rightarrow 0^\pm$ by $\psi_0^{(\pm)}$, and its population in each site as $n_j^{(\pm)}$, where $j = 0$ for the dot and $1 \leq j \leq L$ for the lead sites,




Figure 3. The occupation of the dot and the first lead sites of the CDW states $\psi_0^{(+)}$ (circles) and $\psi_0^{(-)}$ (squares), of a lead with 300 sites and $I = 4$ which is coupled to the dot (filled symbols) with $V = 0.8$. For $\epsilon_0 = 0$ these two states are degenerate in the thermodynamic limit. Lines are guides to the eye.

the particle-hole symmetry implies that $n_j^{(-)} = 1 - n_j^{(+)}$ everywhere. For odd L (even $L + 1$) one has the requirement $n_j^{(-)} = n_{L-j}^{(+)}$, and the number of electrons in both $\psi_0^{(+)}$ and $\psi_0^{(-)}$ is $N = (L + 1)/2$. When L is even ($L + 1$ is odd), however, one has an additional symmetry requirement, $n_j = n_{L-j}$, for both $\psi_0^{(+)}$ and $\psi_0^{(-)}$, and the states differ in their electron number: $\psi_0^{(+)}$ contains $N = L/2$ electrons, for any $I > 2$, while $\psi_0^{(-)}$ has $N = L/2 + 1$. It is worth noting that $n_1^{(+)} > 1/2$, while $n_1^{(-)} < 1/2$.

When $\epsilon_0 \neq 0$, i.e. a dot with a finite on-site energy is connected to the lead, it has some local influence on the ground state in its vicinity. For a lead in the CDW phase, the two states described above are slightly modified, and can be denoted by $\psi^{(+)}$ and $\psi^{(-)}$. Nevertheless, the main influence of the dot is to lift the degeneracy between these states. If $\epsilon_0 < 0$, the dot state population is high and $\psi^{(-)}$ is energetically preferable due to the dot-lead hopping term. For $\epsilon_0 > 0$ the opposite happens, resulting in a preference of $\psi^{(+)}$. The occupation of the dot-lead system in the CDW phase is shown in figure 3. When the dot's orbital energy changes from a negative to a positive value, the system switches from $\psi^{(-)}$ to $\psi^{(+)}$. In the following we show, by a calculation of the free energy, that this is indeed a first order QPT. The resulting phase diagram is drawn in the inset of figure 4.

Before discussing the free energy in the CDW phase, we turn to an exact calculation of the energy and of the total number of electrons in a non-interacting case. Using the Green function technique [19], one can find that the change in the total density of states in the system, due to the presence of the dot, is

$$\Delta\nu(\epsilon) = \frac{1}{\pi} \text{Im} \frac{\partial}{\partial\epsilon} \ln(\epsilon - \epsilon_0 - \Sigma(\epsilon)), \quad (4)$$

where $\Sigma(\epsilon) = (V/t)^2\epsilon/2 + i(V^2/t)\sqrt{1 - (\epsilon/2t)^2}$ is the self-energy of an electron in the dot [7].




Figure 4. The total number of electrons in the system, as a function of ϵ_0 , for $L = 299$, $\mu = 0$, $V = 0.3$ and different values of I . Symbols—DMRG results. Lines—fit to equation (5). Inset: the phase diagram; a Kosterlitz–Thouless transition (solid line) occurs at $I = 2$, and a first order phase transition (dashed line) at $\epsilon_0 = 0$ for $I > 2$.

Therefore, the change in the number of electrons in the entire system is

$$\Delta N(\mu, \epsilon_0) = \int_{-2t}^{\mu} \Delta v(\epsilon) d\epsilon = \frac{1}{2} + \frac{1}{\pi} \tan^{-1} \frac{\mu - \epsilon_0 - \text{Re } \Sigma(\mu)}{\text{Im } \Sigma(\mu)}, \quad (5)$$

and the change in the free energy of the system is

$$\Delta \Omega(\mu, \epsilon_0) = \Delta E - \mu \Delta N(\mu) = \int_{-2t}^{\mu} (\epsilon - \mu) \Delta v(\epsilon) d\epsilon = - \int_{-2t}^{\mu} \Delta N(\epsilon, \epsilon_0) d\epsilon. \quad (6)$$

In figure 4 the total number of electrons in the system is presented. For $I = 0$, N fits the predicted formula (equation (5)) well. As noted above, the agreement between the numerical results for a finite lead and the exact result for a semi-infinite lead is obtained due to the fact that the number of particles can vary during the DMRG process. For the LL phase ($0 < I < 2$), $N(\epsilon_0)$ looks quite similar, varying between $L/2 + 1$ (at $\epsilon_0 \rightarrow -\infty$) and $L/2$ (at $\epsilon_0 \rightarrow \infty$), taking the average value $(L + 1)/2$ at $\epsilon_0 = 0$. As in the case of n_{dot} , the results fit equation (5) with a renormalized dot–lead coupling for small values of I . Increasing I towards the transition point ($I > 1.5$) results in a less accurate fit.

The values of $V_{\text{eff}}(I)$ obtained by the fit of the N curves to equation (5) are in good agreement with the values obtained by the fit of n_{dot} to equation (2). We find that V_{eff} decreases monotonically with increasing I , exhibiting the RPA-like behaviour described in [7].

The CDW phase ($I > 2$), however, is qualitatively different: for an odd lead length, $N(\epsilon_0)$ remains a constant integer ($N = (L + 1)/2$), which does not depend on ϵ_0 at all. For even L , $N = L/2 + 1$ for $\epsilon_0 < 0$, and $N = L/2$ for $\epsilon_0 > 0$. This is a direct result from the switch of the system ground state from $\psi^{(-)}$ to $\psi^{(+)}$ at $\epsilon_0 = 0$. For an odd lead length, the total number of sites in the system ($L + 1$) is even, so that N is equal for both states. For an even lead length, $L + 1$ is odd, and the total number of electrons is changed by one when ϵ_0 passes 0. Except for the decrease of one electron at $\epsilon_0 = 0$ for an even lead, N remains constant in the CDW phase, independent of ϵ_0 . Thus even with the continuous change in the population of the dot as a function of the dot’s level for $\epsilon_0 \neq 0$, the number of electrons in the entire system remains




Figure 5. The ground state free energy as a function of ϵ_0 for different interaction strengths. The lead-dot coupling was taken as $V = 0.3$, and the interaction strength I takes values between 0 and 3.5, in jumps of 0.5. The lines represent the DMRG results (dashed line— $I = 2$), while the symbols show the fit to equation (6) for values of I lower than 2. Inset: $\Omega(\epsilon_0) - \Omega(-\epsilon_0)$ as a function of ϵ_0 for $I = 0, 1, 2, 3$ (symbols) and $\Omega(\epsilon_0) - \Omega(-\epsilon_0) = \epsilon_0$ (line). For all values of interaction we get $|\Omega(\epsilon_0) - \Omega(-\epsilon_0) - \epsilon_0| < 10^{-6}$.

constant. The change in the occupation of the dot as a function of ϵ_0 is compensated by the lead.

This difference in the behaviour of N as a function of ϵ_0 between the two phases (i.e. constant for the CDW phase, compared to a continuous decrease for the LL phase) is a direct manifestation of their transport properties. The LL phase is metallic, and therefore compressible. Hence, infinitesimal changes of the electron number are possible. On the other hand, the CDW phase is insulating and thus incompressible, which results in a constant N .

In figure 5, typical numerical results for the free energy $\Omega(\epsilon_0) - \Omega(0)$ as a function of ϵ_0 are shown. A perfect fit between the DMRG results and the exact formula equation (6) for $I = 0$ is obtained. In the LL phase ($0 < I < 2$) our DMRG calculations show that the effect of interactions on Ω can be fitted by replacing V in equation (6) by the same effective coupling V_{eff} as obtained for the behaviour of n_{dot} discussed in [7], and for the behaviour of N discussed above. For the CDW phase ($I > 2$), however, there is obviously a qualitative change in the energy curve: the dependence of Ω on ϵ_0 is linear both below and above $\epsilon_0 = 0$, with an abrupt change of $\frac{d\Omega}{d\epsilon_0}$ at $\epsilon_0 = 0$.

These results point out that the single impurity, connected at one end of a long interacting lead, has a well defined influence on the ground state of the entire coupled system. As discussed above, when the dot level passes through $\epsilon_0 = 0$, the lead's population is inverted at every site, leading to an inversion of the CDW order parameter, as presented in figure 2 (inset). The dot population is inverted as well. As a result, a dramatic change in the dependence of the free energy of the system on ϵ_0 occurs.

Since Ω is the free energy of the system, the jump in its first derivative might be a sign of a first order QPT. In order to see whether this non-analyticity of the free energy is just a trivial level crossing (LC) of two levels in the system, the dependence of the transition shape on the system size L is explored. For the case where the sharp transition in the energy results from the




Figure 6. The ground state free energy as a function of ϵ_0 for different CDW system sizes, with $V = 0.3$ and $I = 2.5$ (dotted, dashed-dotted and dashed lines). Inset: the results for $L = 29$ are compared to the case of $I = -2.5$, which is inside the ferromagnetic phase, and to the CDW case with $L = 30$.

fact that the external field (in our case the gate voltage) commutes with the Hamiltonian, and thus an LC is possible, no size dependence of the sharpness of the transition is expected. On the other hand, a real QPT will become sharp only in the thermodynamic limit (i.e. semi-infinite lead).

In order to compare the two scenarios (LC versus QPT) we solve the same Hamiltonian (equation (1)) with strong attractive interactions, i.e. $I < -2t$. In this case, transforming the Hamiltonian to an XXZ spin chain, the strong attractive interaction regime corresponds to the ferromagnetic phase [16] (while the CDW is equivalent to the Néel phase). In the fermionic case these attractive interactions yield a twofold degenerate ground state, composed of states in which the sites are either entirely occupied or empty (if there is an explicit restriction to half filling, a ‘phase separation’ evolves). Pinning by the dot lifts the degeneracy, because an occupied dot causes a preference of an empty lead, and vice versa. Thus the influence of the dot on the coupled wire is superficially similar to the CDW case in so far as in both cases the ground state of the entire system is determined by the dot orbital. Nevertheless, the ferromagnetic system is clearly an LC system, and no dependence on the system size is expected.

The dependence of Ω on ϵ_0 was calculated for $L = 29, 99, 299$ in the CDW case, and for $L = 29$ in the ferromagnetic case. From the results shown in figure 6 (and its inset) it is clear that these two cases are different. The ferromagnetic system is indeed a trivial LC system, with two competing states whose energies cross each other for $\epsilon_0 = 0$. Since these two states are eigenstates of the Hamiltonian even for a finite L , the size does not play a role, so that even for $L = 29$ one can see a sharp transition between the two ground states.

On the other hand, for a small (but with an odd L) CDW system Ω shows a smooth dependence on ϵ_0 , without any non-analyticity. As a matter of fact, for any finite system the electron levels are expected to be mixed, resulting in avoided crossing of the two lowest many-body levels [14]. In other words, the CDW states are not true eigenstates of the Hamiltonian for a finite system. Indeed, indications for the sudden jump of $\frac{d\Omega}{d\epsilon_0}$ are seen only for larger system sizes (i.e. for $L \gtrsim 200$).




Figure 7. The dependence of the dot occupation on ϵ_0 for different CDW system sizes, with $V = 0.3$ and $I = 2.5$. Inset: the absolute value of the slope in the limit $\epsilon_0 \rightarrow 0$ as a function of L for short leads, together with a linear fit.

The dot-dashed line in figure 6 (inset) shows that for a short lead of an even size ($L = 30$ in this case) there is, however, a non-analyticity in Ω as a function of ϵ_0 . The comparison between even and odd lead sizes emphasizes our conclusion stated above. While for an odd lead size $\psi^{(-)}$ and $\psi^{(+)}$ have the same number of electrons, for an even size of the lead the transition from $\psi^{(-)}$ to $\psi^{(+)}$ at $\epsilon_0 = 0$ involves a decrease of the electron number by one. The Hamiltonian \hat{H} conserves the number of particles, so that in the case of L even, these two states are not coupled, and the transition between them is a simple LC, not showing a size dependence. For an odd L , however, $\psi^{(-)}$ and $\psi^{(+)}$ are coupled by \hat{H} , so that for a finite L they are actually mixtures of the CDW states, thus presenting sharper $\Omega(\epsilon_0)$ dependence for larger systems. We thus conclude, based on these two comparisons, that in the case of a CDW with an odd L this transition is a QPT, which happens for $L \rightarrow \infty$, i.e. for a semi-infinite lead¹.

As pointed out above, in the numerical results for large lead sizes, the non-analyticity of the free energy is seen clearly, so that it is obviously a first order transition. However, a similar conclusion can be drawn from the results of short leads, by scaling the results against the lead size L . Although for finite size systems the order parameter changes continuously, the slope of this change grows with the system size L . For a first order transition in d dimensions one expects a power law dependence of the slope as L^d , while for a second order transition the power law should be fixed by some universal exponents [20].

For the CDW model, the population of the dot plays the role of an order parameter, and one can check its behaviour near $\epsilon_0 = 0$ for different lead sizes. In figure 7 the occupation as a function of ϵ_0 is shown for some short lead sizes ($L = 29, 49, 69$) and for a large one ($L = 299$). It is indeed seen that while for a long enough lead there is a jump in the dot population, for short leads there is a continuous change in the occupation, with a larger slope for larger lengths. Zooming into the regime of $\epsilon_0 = 0$ we find that, although $L \gtrsim 200$ is

¹ Of course, when ϵ_0 crosses the Fermi energy, the system will remain in the previous ground state, which becomes a metastable state. In the thermodynamic limit an exponentially long time will be required for the system to switch to the true ground state.

required in order to see a clear discontinuity in Ω , $L \gtrsim 100$ is long enough to show a jump in n_{dot} . The dependence of the slope on L , for $L < 100$, is shown in the inset of figure 7. As one expects from finite size scaling predictions for first order phase transition, there is a very good linear fit of the order parameter to the lead size.

We now turn to another consequence of the particle–hole symmetry. For $\epsilon_0 \neq 0$ the symmetry results in the fact that the ground states for $+\epsilon_0$ and for $-\epsilon_0$ ($\psi^{(+)}$ and $\psi^{(-)}$ respectively) have an inverted population everywhere. It is thus clear that $\langle \psi^{(+)} | \hat{H}_{\text{lead}} | \psi^{(+)} \rangle = \langle \psi^{(-)} | \hat{H}_{\text{lead}} | \psi^{(-)} \rangle$, and similarly $\langle \psi^{(+)} | \hat{H}_{\text{dot-lead}} | \psi^{(+)} \rangle = \langle \psi^{(-)} | \hat{H}_{\text{dot-lead}} | \psi^{(-)} \rangle$. This implies $\Omega(\epsilon_0) - \Omega(-\epsilon_0) = \langle \psi^{(+)} | \hat{H}_{\text{dot}} | \psi^{(+)} \rangle - \langle \psi^{(-)} | \hat{H}_{\text{dot}} | \psi^{(-)} \rangle = \epsilon_0 n_{\text{dot}}^+ - (-\epsilon_0) n_{\text{dot}}^- = \epsilon_0$, (7) where n_{dot}^+ (n_{dot}^-) represents the dot populations of $\psi^{(+)}$ ($\psi^{(-)}$). The last equality results from the symmetric population $n_{\text{dot}}^- = 1 - n_{\text{dot}}^+$.

The relation equation (7) does not depend on the interaction, and should exist for both the LL and the CDW phases. Obviously it should be obeyed for the non-interacting case, where the energy is given by equation (6), thus leading to the following nontrivial relation:

$$-\frac{1}{\pi} \int_{-2t}^{2t} \tan^{-1} \frac{(1 - \frac{V^2}{2t^2})\epsilon - \epsilon_0}{\frac{V^2}{t} \sqrt{1 - \frac{\epsilon^2}{4t^2}}} d\epsilon = \epsilon_0, \quad (8)$$

in which the result of the integral on the LHS does not depend on the parameters V and t .

A physical insight into equation (8) can be gained by taking the derivative of both sides with respect to ϵ_0 , and using the definition of the self-energy given above. This yields

$$\frac{1}{\pi} \int_{-2t}^{2t} \frac{\text{Im } \Sigma(\epsilon)}{(\epsilon - \epsilon_0 - \text{Re } \Sigma(\epsilon))^2 + (\text{Im } \Sigma(\epsilon))^2} d\epsilon = 1, \quad (9)$$

which is evident since the LHS is the occupation of the dot [7] when $\mu > 2t$.

In figure 5 (inset) a plot of $\Omega(\epsilon_0) - \Omega(-\epsilon_0)$ as a function of ϵ_0 is shown, for values of I between 0 and 3. As can be seen, equation (7) is valid for all values of I .

In conclusion, we have shown that the occupation of a dot coupled to a one-dimensional wire can be used to identify the different phases in the wire. It is known that as a function of the interaction strength I the wire goes through three different phases (which may also be mapped onto the phases of an XXZ spin chain). For $I < -2t$ the ground state is doubly degenerate, with all sites equally full or empty, corresponding to the ferromagnetic state of a spin chain. In the intermediate range of interactions ($-2t < I < 2t$) the translational symmetry is unbroken and a Luttinger liquid phase occurs. For $I > 2t$ the ground state is doubly degenerate, corresponding to the two possible CDW states or the two different antiferromagnetic realizations for the spin chain.

For the Luttinger liquid phase the occupation of the impurity does not show a jump when the impurity level crosses the Fermi energy. In the two other phases the impurity level splits the degeneracy by favouring one of the two ground states, depending on whether the impurity level is empty or filled. Nevertheless, we have shown that the physics in these cases is different. While for the ferromagnetic phase a simple level crossing occurs with a sharp jump in occupation of the impurity for any length of the wire, in the CDW phase the position of the impurity level drives a first order QPT in the thermodynamical limit between the two CDW states, while for a finite wire the jump is smeared. This phase transition shows all the hallmarks of a first order QPT, such as a size dependence, a jump in the order parameter and a discontinuity of the derivative of the grand canonical potential.

Acknowledgment

Support from the Israel Academy of Science (grant 877/04) is gratefully acknowledged.

References

- [1] Tomonaga S-I 1950 *Prog. Theor. Phys.* **5** 544
Luttinger J M 1963 *J. Math. Phys.* **4** 1154
- [2] For a review, see Voit J 1994 *Rep. Prog. Phys.* **57** 977
- [3] Kane C L and Fisher M P A 1992 *Phys. Rev. Lett.* **68** 1220
Kane C L and Fisher M P A 1992 *Phys. Rev. B* **46** 7268
Kane C L and Fisher M P A 1992 *Phys. Rev. B* **46** 15233
- [4] Bockrath M *et al* 1999 *Nature* **397** 598
Postma H W C *et al* 2001 *Science* **293** 76
- [5] Auslander O M *et al* 2005 *Science* **308** 88
- [6] Furusaki A and Matveev K A 2002 *Phys. Rev. Lett.* **88** 226404
- [7] Sade M, Weiss Y, Goldstein M and Berkovits R 2005 *Phys. Rev. B* **71** 153301
- [8] Johnson A C, Marcus C M, Hanson M P and Gossard A C 2004 *Phys. Rev. Lett.* **93** 106803
- [9] Grishin A, Yurkevich I V and Lerner I V 2004 *Phys. Rev. B* **69** 165108
- [10] Nazarov Yu V and Glazman L I 2003 *Phys. Rev. Lett.* **91** 126804
- [11] Luther A and Peschel I 1975 *Phys. Rev. B* **12** 3908
- [12] Haldane F D M 1980 *Phys. Rev. Lett.* **45** 1358
- [13] Schulz H J 1990 *Phys. Rev. Lett.* **64** 2831
- [14] Sachdev S 1999 *Quantum Phase Transitions* (Cambridge: Cambridge University Press)
- [15] White S R 1993 *Phys. Rev. B* **48** 10345
- [16] Mikeska H J and Kolezhuk A K 2004 *Quantum Magnetism (Springer Lecture Notes in Physics vol 645)* ed U Schollwock, J Richter, D J J Farnell and R F Bishop (Berlin: Springer)
- [17] Pang H, Liang S and Annett J F 1993 *Phys. Rev. Lett.* **71** 4377
- [18] Baxter R J 1973 *J. Phys. C: Solid State Phys.* **6** L94
Baxter R J 1989 *Exactly Solved Models In Statistical Mechanics* (London: Academic Press/Harcourt Brace Jovanovich Publishers)
- [19] Mahan G D 1990 *Many Particle Physics* (New York: Plenum)
- [20] Binder K and Landau D P 1984 *Phys. Rev. B* **30** 1477