

Electromagnetic pulse-driven spin-dependent currents in semiconductor quantum rings

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2009 J. Phys.: Condens. Matter 21 145801

(<http://iopscience.iop.org/0953-8984/21/14/145801>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 129.252.86.83

The article was downloaded on 29/05/2010 at 18:58

Please note that [terms and conditions apply](#).

Electromagnetic pulse-driven spin-dependent currents in semiconductor quantum rings

Zhen-Gang Zhu and Jamal Berakdar

Institut für Physik, Martin-Luther Universität Halle-Wittenberg,
Heinrich-Damerow-Straße 4 06120 Halle, Germany

Received 2 January 2009, in final form 9 February 2009

Published 5 March 2009

Online at stacks.iop.org/JPhysCM/21/145801

Abstract

We investigate the non-equilibrium charge and spin-dependent currents in a quantum ring with a Rashba spin-orbit interaction (SOI) driven by two asymmetric picosecond electromagnetic pulses. The equilibrium persistent charge and persistent spin-dependent currents are investigated as well. It is shown that the dynamical charge and the dynamical spin-dependent currents vary smoothly with a static external magnetic flux and the SOI provides a $SU(2)$ effective flux that changes the phases of the dynamic charge and the dynamic spin-dependent currents. The period of the oscillation of the total charge current with the delay time between the pulses is larger in a quantum ring with a larger radius. The parameters of the pulse fields control to a certain extent the total charge and the total spin-dependent currents. The calculations are applicable to nanometre rings fabricated in heterojunctions of III-V and II-VI semiconductors containing several hundreds of electrons.

(Some figures in this article are in colour only in the electronic version)

1. Introduction

The study of the spin-orbit interaction (SOI) in semiconductor low-dimensional structures and its application for spintronics devices have attracted much attention recently [1]. There are two important kinds of SOI in conventional semiconductors: one is the Dresselhaus SOI induced by bulk inversion asymmetry [2], and the other is the Rashba SOI caused by structure inversion asymmetry [3]. As pointed out in [4], the Rashba SOI is dominant in a narrow gap semiconductor system and the strength of the Rashba SOI can be tuned by an external gate voltage in HgTe [5], InAs [6], $\text{In}_x\text{Ga}_{1-x}\text{As}$ [7], and GaAs [3, 8] quantum wells. Recent research has focused on the electrically induced generation of a spin-dependent current (SC) mediated by SOI-type mechanism, e.g. as in the intrinsic spin Hall effect in a three-dimensional (3D) p-doped semiconductor [9] and in a two-dimensional (2D) electron gas with Rashba SOI [10]. Here we study high quality spin-interacting quantum rings (QRs) with a radius on the nanometre scale [11, 12]. These systems show Aharonov-Bohm-type (AB) spin-interferences [13, 14]. In particular we investigate the dynamics triggered by time-dependent electric fields as provided by time-asymmetric pulses [15] or tailored

laser pulse sequences [16]. The quantity under study is the spin-resolved pulse-driven current, in analogy to the spin-independent case [17–19]. In a previous work [20], we investigated the dynamical response of the charge polarization to the pulse application. No net charge or spin-dependent current is generated because the clockwise and anti-clockwise symmetry of the carrier is not broken by one pulse or a series of pulses having the same linear polarization axis. This symmetry is lifted if two time-delayed pulses with non-collinear polarization axes are applied [17]. However, to our knowledge all previous studies on light-induced currents in quantum rings did not consider the coupling of the spin to the orbital motion (and hence to the light field), which is addressed in this work. As detailed below, having done that, it is possible to control dynamically the spin-dependent current in a 1D quantum ring with Rashba SOI by using two time-delayed linearly polarized electromagnetic pulses. For transparent interpretation of the results only the Rashba SOI is considered in this work. The presence of the Dresselhaus SOI may change qualitatively the results presented here for the spin-dependent non-equilibrium dynamics of the carriers, which can be anticipated from the findings for the equilibrium case [21].

2. Theoretical model

We study the response of charges and spins confined in a one-dimensional (1D) ballistic QR with SOI to the application of two short time-delayed linearly polarized asymmetric electromagnetic pulses [17, 22]. The effective single-particle Hamiltonian reads $\hat{H}' = \hat{H}_{\text{SOI}} + \hat{H}_1(t)$ [20], with

$$\hat{H}_{\text{SOI}} = \frac{\mathbf{p}^2}{2m^*} + V(\mathbf{r}) + \frac{\alpha_R}{\hbar}(\hat{\sigma} \times \mathbf{p})_z, \quad (1)$$

$$\hat{H}_1(t) = -e\mathbf{r} \cdot \mathbf{E}(t) + \mu_B \mathbf{B}(t) \cdot \hat{\sigma}.$$

$\mathbf{E}(t)$ and $\mathbf{B}(t)$ are the electric and the magnetic fields of the pulse. Integrating out the r dependence \hat{H}_{SOI} reads in cylindrical coordinates [23–27]

$$\hat{H}_{\text{SOI}} = \frac{\hbar\omega_0}{2} \left[\left(i\partial_\varphi + \frac{\phi}{\phi_0} - \frac{\omega_R}{2\omega_0}\sigma_r \right)^2 - \left(\frac{\omega_R}{2\omega_0} \right)^2 + \frac{\omega_B}{\omega_0}\sigma_z \right]. \quad (2)$$

$\partial_\varphi = \frac{\partial}{\partial\varphi}$, $\phi_0 = h/e$ is the flux unit, $\phi = B\pi a^2$ is the magnetic flux threading the ring, a is the radius of the ring, $\hbar\omega_0 = \hbar^2/(m^*a^2) = 2E_0$, $\hbar\omega_R = 2\alpha_R/a$, $\hbar\omega_B = 2\mu_B B$ and B are due to a possible external static magnetic field $\mathbf{B} = B\hat{\mathbf{e}}_z$. The single-particle eigenstates of \hat{H}_{SOI} are represented as $\Psi_n^S(\varphi) = e^{i(n+1/2)\varphi} v^S(\gamma, \varphi)$ where $v^S(\gamma, \varphi) = (a^S e^{-i\varphi/2}, b^S e^{i\varphi/2})^T$ are spinors in the angle-dependent local frame, and $a^\uparrow = \cos(\gamma/2)$, $b^\uparrow = \sin(\gamma/2)$, $a^\downarrow = -\sin(\gamma/2)$, $b^\downarrow = \cos(\gamma/2)$, (T means transposed) where $\tan \gamma = -Q_R = -\omega_R/\omega_0$ (if we ignore the Zeeman splitting caused by the static magnetic field [27, 28]). γ describes the direction of the spin quantization axis, as illustrated in figure 1(a). The energy spectrum of the QR with the SOI reads [20, 24–28]

$$E_n^S = \frac{\hbar\omega_0}{2} \left[\left(n - \phi/\phi_0 + \frac{1-Sw}{2} \right)^2 - \frac{Q_R^2}{4} \right], \quad (3)$$

$$w = \sqrt{1 + Q_R^2} = 1/\cos \gamma,$$

where $S = +1$ ($S = -1$) stands for spin up (spin down) in the local frame.

3. Pulse-driven single-particle dynamics

We apply two time-asymmetric pulses to the system (see figure 1(b)). The first one (at $t = 0$) propagates in the z direction and has a duration τ_d . Its E -field is along the x direction. τ_d is chosen to be much shorter than the ballistic time of the carriers in which case the QR states develop as [17, 22, 29]

$$\Psi_n^S(\varphi, t > 0) = \Psi_n^S(\varphi, t < 0) e^{i\alpha_1 \cos \varphi}, \quad (4)$$

$$\alpha_1 = eap/\hbar, \quad p = -\int_0^{\tau_d} \mathbf{E}(t) dt,$$

where $\mathbf{E}(t) = Ff(t)$, F and $f(t)$ describe the amplitude and the time dependence of the electric field of the pulse respectively. In the following, we use F_1 and F_2 to characterize




Figure 1. (a) A schematic graph of the geometry, spin configuration and the applied pulses is shown. (b) Time-delayed asymmetric pulses are schematically drawn. (c) Energy spectrum for a ring with spin-orbit interaction. Δ_S defines the distance between the spectrum symmetry axis and the smallest nearest integer.

the first and the second pulses. The pulse effect is encapsulated entirely in the *action parameter* α_1 . With the initial conditions $n(t < 0) = n_0$ and $S(t < 0) = S_0$ and using equations (4) one finds

$$\Psi_{n_0}^{S_0}(\varphi, t) = \frac{1}{\sqrt{2\pi}} \sum_{ns} C_n^S(n_0, S_0, t) e^{i(n+1/2)\varphi} e^{-iE_n^S t/\hbar} |v^S\rangle, \quad (5)$$

with

$$C_n^S = \begin{cases} \delta_{SS_0} \delta_{nm_0} & \text{for } t \leq 0, \\ \delta_{SS_0} i^{n_0-n} J_{n_0-n}(\alpha_1) & \text{for } t > 0, \end{cases} \quad (6)$$

where J_n is the n th order Bessel function. For the time-dependent energy we find

$$E_{n_0}^{S_0}(t > 0) = E_{n_0}^{S_0}(t < 0) + \frac{\hbar\omega_0}{2} \frac{\alpha_1^2}{2}, \quad (7)$$

with $E_{n_0}^{S_0}(t < 0)$ given by equation (3). Applying a second pulse at $t = \tau$ with the same duration τ_d but the electric field being along the y axis (see figure 1(b)), the wavefunctions develop as $\Psi_{n_0}^{S_0}(\varphi, t > \tau) = \Psi_{n_0}^{S_0}(\varphi, t < \tau) e^{i\alpha_2 \sin \varphi}$, where α_2 is the action parameter associated with the second pulse. $\Psi_{n_0}^{S_0}(\varphi, t = \tau^-)$ follows from equation (5). For $t > \tau$ the expansion coefficients behave as $C_n^{S'}(n_0, S_0, t > \tau) = \sum_n \delta_{S'S_0} [i^{n_0-n} J_{n_0-n}(\alpha_1) J_{n'-n}(\alpha_2)] e^{i(E_n^{S'} - E_{n_0}^{S_0})\tau/\hbar}$.

4. Non-equilibrium spin and charge currents

A single pulse does not generate in QR any net charge current because of the degeneracy of the orbital states. However,

the charge will be polarized [20, 22] and the corresponding dipole moments oscillate in the x direction with an associated optical emission. Applying a second pulse as described above leads to a non-equilibrium net current, in addition to the persistent charge current caused by the static flux and the SOI which causes a $SU(2)$ vector potential and manifests itself in an induced spin-dependent persistent charge current [27, 28, 30, 31]. Consequently, a non-equilibrium spin-dependent current is induced.

The line velocity operator is [32]

$$\hat{\mathbf{v}}_\varphi = \hat{\mathbf{e}}_\varphi \left\{ \frac{-i\hbar}{m^*a} \partial_\varphi - \frac{\hbar}{m^*a} \frac{\phi}{\phi_0} + \frac{\alpha_R}{\hbar} \sigma_r \right\}$$

which is associated with the operator of the angular velocity $\hat{\mathbf{v}}_\varphi/a$ [33]. Contributions to the persistent charge current from each QR level read [34]

$$\mathbf{I}_{n_0, S_0} = \frac{1}{2\pi} \int_0^{2\pi} d\varphi \int_{r_1}^{r_2} dr \mathbf{j}_{n_0, S_0}^\varphi(\mathbf{r}', t > \tau),$$

where

$$\mathbf{j}_{n_0, S_0}^\varphi = e \operatorname{Re}[\Psi_{n_0}^{S_0, \dagger}(\mathbf{r}', t) \hat{\mathbf{v}}_\varphi \Psi_{n_0}^{S_0}(\mathbf{r}', t)].$$

Upon algebraic manipulations we find

$$\mathbf{I}_{n_0, S_0}(t > \tau) = \mathbf{I}_{n_0, S_0}^{(0)}(t > \tau) + \mathbf{I}_{n_0, S_0}^{(1)}(t > \tau). \quad (8)$$

The index ‘(0)’ stands for the static persistent charge current (PCC) which exists in the absence of pulse field, whereas the index ‘(1)’ indicates the pulse-induced dynamic charge current (DCC). The PCC is caused by a magnetic $U(1)$ flux and has been studied extensively [35] without [34, 36, 37] or with the spin interactions [33]. It has been experimentally observed both in gold rings of radius 1.2 and 2.0 μm [38] and in a GaAs–AlGaAs ring of radius about 1 μm [39]. The SOI scattering effects were also studied [40]. The PCC carried by the states characterized by n_0 and S_0 reads (please note the current in this work is defined as flow of positive charges, which is opposite to the direction of the flow of electrons)

$$\mathbf{I}_{n_0, S_0}^{(0)}(t > \tau) = \hat{\mathbf{e}}_\varphi I_0 \left(n_0 - \frac{\phi}{\phi_0} + \frac{1 - S_0 w}{2} \right), \quad (9)$$

where $I_0 = 2E_0 a / \phi_0$ is the unit of CC, the second term on the right-hand side of equation (9) stems from the static magnetic field; the third term is a consequence of the $SU(2)$ flux of the SOI [27]. The DCC part is

$$\mathbf{I}_{n_0, S_0}^{(1)}(t > \tau) = \hat{\mathbf{e}}_\varphi I_0 \{ \alpha_2 (\cos \varphi)_{n_0}^{S_0}(\tau) \}, \quad (10)$$

where

$$\begin{aligned} (\cos \varphi)_{n_0}^{S_0}(\tau) &= \alpha_1 h(\Omega_{\tau 1}) \sin b_\tau \\ &\times \cos \left[2 \left(n_0 - \frac{\phi}{\phi_0} + \frac{1 - S_0 w}{2} \right) b_\tau \right], \end{aligned}$$

$$b_\tau = \omega_0 \tau / 2, \quad \Omega_{\tau 1} = \alpha_1 \sqrt{2(1 - \cos(2b_\tau))},$$

$$h(\Omega_{\tau 1}) = J_0(\Omega_{\tau 1}) + J_2(\Omega_{\tau 1}).$$

To obtain the total persistent charge current and the dynamic current we have to consider the spin-resolved

occupations of the single-particle states. For simplicity we operate at zero temperatures and ignore the relaxation caused by phonons or other mechanisms, i.e. we confine ourselves to times shorter than the relaxation time. The general case can be developed along the lines of [41].

At first we introduce an effective flux as

$$\phi_S = \phi - \phi_0 \frac{1 - S w}{2}. \quad (11)$$

As evident from equations (3) the spectrum is symmetric with respect to $x_S = \phi_S / \phi_0$. Further, we define the shift $\Delta_S = x_S - l(l')$, where $S = \uparrow$ or \downarrow . Here $l(l') = [x_{\uparrow(\downarrow)}]$ where $[x]$ means the nearest integer which is less than x . Δ_S is shown in figure 1. When $\Delta_S = 1$, it is equivalent to $\Delta_S = 0$. Furthermore, $\bar{\Delta}_S = |1/2 - \Delta_S|$ is the distance between the x_S and the nearest half integer.

4.1. Spinless particles

For N spinless particles we distinguish two cases: N is an even or an odd integer.

Case (1). If N is an even integer then

$$I_{\text{even}}^{(0)}(\Delta) = \operatorname{sgn}(\Delta) N \left(\frac{1}{2} - \Delta \right), \quad (12)$$

$$I_{\text{even}}^{(1)}(\Delta) = \alpha_1 \alpha_2 h(\Omega_{\tau 1}) \sin(N b_\tau) \cos(1 - 2\Delta) b_\tau,$$

where $\operatorname{sgn}(x)$ equals $+1$, for $x > 0$; 0 for $x = 0$, and -1 for $x < 0$.

Case (2). If N is an odd integer then

$$I_{\text{odd}}^{(0)}(\Delta) = -\operatorname{sgn}(\Delta) \operatorname{sgn} \left(\frac{1}{2} - \Delta \right) N \left(\frac{1}{2} - \bar{\Delta} \right), \quad (13)$$

$$I_{\text{odd}}^{(1)}(\Delta) = \alpha_1 \alpha_2 h(\Omega_{\tau 1}) \sin(N b_\tau) \cos(1 - 2\bar{\Delta}) b_\tau.$$

4.2. Particles with 1/2 spin

For spin 1/2 particles we consider four cases.

Case (0). For an even number of particle pairs, i.e. $N = 4m$, where m is an integer we find

$$I_S^{(0)}(\Delta_S) = I_{\text{even}}^{(0)}(\Delta_S), \quad I_S^{(1)}(\Delta_S) = I_{\text{even}}^{(1)}(\Delta_S). \quad (14)$$

Case (1). For an odd number of particle pairs, i.e. $N = 4m + 2$ we obtain

$$I_S^{(0)}(\Delta_S) = I_{\text{odd}}^{(0)}(\Delta_S), \quad I_S^{(1)}(\Delta_S) = I_{\text{odd}}^{(1)}(\Delta_S). \quad (15)$$

Case (2). For an even number of pairs plus one extra particle, i.e. $N = 4m + 1$ (there is one particle whose spin is unpaired as compared with case (0)) we find

$$\begin{aligned} I_{\text{ext}, S}^{(0)}(\Delta_S) &= -\operatorname{sgn}(\Delta_S) \operatorname{sgn} \left(\frac{1}{2} - \Delta_S \right) \\ &\times \left(\frac{N-1}{4} + \frac{1}{2} - \bar{\Delta}_S \right), \end{aligned} \quad (16)$$

$$\begin{aligned} I_{\text{ext}, S}^{(1)}(\Delta_S) &= \alpha_1 \alpha_2 h(\Omega_{\tau 1}) \sin(b_\tau) \\ &\times \cos \left(\frac{N-1}{2} + 1 - 2\bar{\Delta}_S \right) b_\tau. \end{aligned}$$

To determine which spin state is occupied by the extra particle one compares the distance of the symmetric axis to the nearest

half integral axis, i.e. $\bar{\Delta}_S$. The one with the larger distance will be occupied.

Case (3). For an odd number of pairs plus one extra particle, i.e. $N = 4m + 3$. Here we use case (1) and determine the contribution to the current from the extra particle

$$I_{\text{ext},S}^{(0)}(\Delta_S) = \text{sgn}(\Delta_S) \text{sgn}\left(\frac{1}{2} - \Delta_S\right) \times \left(\frac{N-3}{4} + \frac{1}{2} + \bar{\Delta}_S\right), \quad (17)$$

$$I_{\text{ext},S}^{(1)}(\Delta_S) = \alpha_1 \alpha_2 \hbar (\Omega_{\tau_1}) \sin(b_\tau) \times \cos\left(\frac{N-3}{2} + 1 + 2\bar{\Delta}_S\right) b_\tau.$$

Which spin state is occupied by the extra particle is governed by $\bar{\Delta}_S$. The level with the smaller $\bar{\Delta}_S$ is populated.

5. Spin-dependent current (SC)

In the presence of a static magnetic field and the SOI but in the absence of the pulse field, the PCC is accompanied by a persistent SC (PSC). Switching on the pulse field generates a spin-dependent charge current due to the SOI, and also a dynamic SC (DSC) that can be controlled by the parameters of the pulse field. The SC density is

$$\mathbf{j}_{n_0,S_0}^s(\mathbf{r}', t) = \text{Re}\{\Psi_{n_0}^{S_0,\dagger}(\mathbf{r}', t) \hat{\mathbf{v}}' \hat{\mathbf{s}} \Psi_{n_0}^{S_0}(\mathbf{r}', t)\},$$

where

$$\hat{\mathbf{s}} = (\hbar/2) \hat{\sigma}_z \delta(\mathbf{r}' - \mathbf{r})$$

is the local spin density. The SC associated with level n_0 , S_0 is

$$\mathbf{I}_{n_0,S_0}^s(t > \tau) = \frac{1}{2\pi} \int_0^{2\pi} d\phi \int_{r_1}^{r_2} dr' \mathbf{j}_{n_0,S_0}^s(\mathbf{r}', t) \quad (18)$$

and can be *evaluated* as

$$\mathbf{I}_{n_0,S_0}^{s_z} = \mathbf{I}_{s_0} \text{Re} \sum_n |C_n^{S_0}(n_0, S_0, t)|^2 D_{1n}^{S_0}, \quad (19)$$

where

$$\mathbf{I}_{s_0} = \hat{\mathbf{e}}_\phi E_0 a / (2\pi)$$

sets the unit SC and

$$D_{1n}^{S_0} = [(a^{S_0})^2 - (b^{S_0})^2] \left(n - \frac{\phi}{\phi_0}\right) - (b^{S_0})^2. \quad (20)$$

Here

$$(a^{S_0})^2 - (b^{S_0})^2 = S_0 \cos \gamma,$$

and $S_0 = \pm 1$. The SC after applying two pulses to the ring is a sum of two parts

$$\mathbf{I}_{n_0,S_0}^{s_z}(t > \tau) = \mathbf{I}_{n_0,S_0}^{s_z,(0)}(t > \tau) + \mathbf{I}_{n_0,S_0}^{s_z,(1)}(t > \tau), \quad (21)$$

where

$$\mathbf{I}_{n_0,S_0}^{s_z,(0)}(t > \tau) = \mathbf{I}_{s_0} [S_0 \cos \gamma] \left[\left(n - \frac{\phi}{\phi_0}\right) + \frac{1}{2} - \frac{S_0}{2 \cos \gamma} \right],$$

$$= \mathbf{I}_{s_0} [S_0 \cos \gamma] \frac{I_{n_0,S_0}^{(0)}(t > \tau)}{I_0}, \quad (22)$$

is the static PSC [27] and the DSC part is

$$\mathbf{I}_{n_0,S_0}^{s_z,(1)}(t > \tau) = \mathbf{I}_{s_0} [S_0 \cos \gamma] [\alpha_2 \langle \cos \phi \rangle_{n_0,S_0}(\tau)],$$

$$= \mathbf{I}_{s_0} [S_0 \cos \gamma] \frac{I_{n_0,S_0}^{(1)}(t > \tau)}{I_0}. \quad (23)$$

Summing over all occupied energy levels we find

$$\mathbf{I}_{S_0}^{s_z}(t > \tau) = \mathbf{I}_{S_0}^{s_z,(0)}(t > \tau) + \mathbf{I}_{S_0}^{s_z,(1)}(t > \tau), \quad (24)$$

where $(I_{S_0}^{(0),(1)}(t > \tau))$ are PCC and DCC

$$\mathbf{I}_{S_0}^{s_z,(0),(1)}(t > \tau) = \mathbf{I}_{s_0} [S_0 \cos \gamma] \frac{I_{S_0}^{(0),(1)}(t > \tau)}{I_0}. \quad (25)$$

6. Numerical results and discussions

We performed calculations for pulse-driven ballistic quantum rings fabricated by an appropriate confinement in a quantum well of $\text{In}_x\text{Ga}_{1-x}\text{As}/\text{InP}$ [42]. Our results are also valid for other III–V or II–VI semiconductor quantum rings with spin orbit, e.g. a GaAs–AlGaAs quantum well, or HgTe/HgCdTe quantum ring [43]. We shall present the total charge current (TCC) which is a sum of PCC and DCC over all the occupied states. Total spin-dependent current (TSC) is obtained in the same way [28].

Figure 2 shows how the flux and the SOI affect the PCC, DCC and TCC. Without the SOI, the jump of the PCC occurs at integer flux for even pair occupation, shown in figure 2. The jumps are different in other occupations (see [28]), here we only focus on the even pair occupation case for clarity. The periodic sawtooth dependence of the PCC on the flux has been studied before, e.g. [27, 28]. At finite SOI the jumps in PCC are shifted to $\phi/\phi_0 = l + (1 \mp w)/2$; the two jumps are the consequences of a superposition of the contributions from the two spin channels. When the SOI strength is such that $\gamma = -\arccos(1/2n)$, ($n = 1, 2, \dots$) the two jumps become at the half integer which is just the case of $4n+2$ occupation in the absence of the SOI [28]. The slope ratio between the two jumps is the same. As can be inferred from the analytic expressions, DCC (cf figure 2) depends smoothly on the flux. SOI results in a phase shift moving or even exchanging the positions of the minima and maxima, as for $\gamma = -60^\circ$. The origin of the shape of TCC is deduced from those of PCC and DCC. Here the magnitudes of the two contributions is crucial. The PCC magnitude is related to the numbers of charge carriers, while the DCC magnitude is determined primarily by the product of the α_1 and α_2 (that can be externally varied by changing the pulse intensities), the delay time τ , and the ring radius.

Figure 3 shows the TCC dependence on the ring radius (in the absence of the SOI). As expected, a larger α enhances the DCC. On the other hand, α enters the Bessel function argument whose increase suppresses the magnitude of DCC. It can be shown that the period of the oscillation with τ increases with increasing the radius. The magnitudes of the maxima and minima are larger with larger radius.

Now we discuss the spin-dependent current projected onto the z direction [27], i.e. I^{s_z} . The spin-dependent current




Figure 2. Persistent charge current (PCC), dynamic charge current (DCC) and total charge current (TCC) are shown in (a), (b) and (c), respectively. The spin orbit angles are $\gamma = 0^\circ, -20^\circ, -40^\circ$ and -50° for the solid lines, the dashed lines, the dotted lines and the dashed-dotted lines respectively in all graphs. The other parameters are $N = 100, a = 100$ nm, $\tau = 26.3$ ps, $F_1 = F_2 = 1$ kV cm $^{-1}$.




Figure 3. Contour plots of TCC on ϕ and τ are shown for different radii of the ring. $a = 100$ nm, 200 nm, 300 nm and 400 nm in (a), (b), (c) and (d), respectively. The other parameters are $N = 100, \gamma = 0^\circ, F_1 = F_2 = 500$ V cm $^{-1}$.

projected onto the γ direction (e.g. the quantization axis of the local spin frame) is $I^{S_z} = I^\gamma \cos \gamma$ [28].

PSC possesses steps at PCC jumps (figure 4(a)) as a function of ϕ . This can be understood from the ratio of PCC for different spins; the SOI only introduces a relative effective flux shift (see equation (11)) leading to a constant spin-dependent current between the jumps. For DSC versus ϕ (figure 4(b)) the effective flux leads to a shift of the DSC along the flux axis. The physics behind this shift is that SOI provides a $SU(2)$ flux, meaning that the pulse-driven (local frame) spin up electrons experience a different flux than those with down spin, leading to a substantial spin-dependent current. In contrast, a static magnetic flux does not induce a spin-dependent current in the

absence of the SOI. This shift and the jumps in the step function of the PSC explain the behaviour of TCC in figure 4(c).

The control of TCC and TSC by tuning the pulse-field parameters is demonstrated in figure 5. Because the two pulses transfer a net angular momentum setting the electrons in motion but they do not couple directly to the spins, the TCC and TSC show the same pattern with τ and F . From an experimental point of view it is essential to note that we are dealing with non-equilibrium quantities which opens the way for their detection via their emission, e.g. the TCC and TSC can be detected by measuring the current-induced magnetization of the ring and the generated electrostatic potential [44].




Figure 4. PCC and PSC, DCC and DSC, and TCC and TSC are shown in (a), (b) and (c), respectively. The solid lines are the charge current, and the dashed lines are the spin current. $\gamma = -40^\circ$, the other parameters are the same as in figure 2.




Figure 5. TCC (a) and TSC (b) versus pulse strengths $F_1 = F_2$ and delay time τ . $\gamma = -40^\circ$, $\phi = 0.2$. Other parameters are $N = 100$, $a = 400$ nm.

7. General and concluding remarks

In the presence of spin-orbit coupling, two time-delayed appropriately shaped electromagnetic pulses generate spin-dependent charge currents. As shown previously for the spin-independent case [17], the sign of the current and its magnitude are controllable via the delay time and the strengths of the pulses. From a symmetry viewpoint, similar phenomena may be expected to occur for other geometries (wires, squares, etc). However, as shown for unbiased superlattices [45] (without SOI) details of the generated currents may differ qualitatively. Application of an appropriate train of pulses opens the possibility of controlling or even stopping the current [41]. For increasing the magnitude of the current more intense pulses should be applied.

For generating currents in quantum rings one may also apply circular polarized laser pulses [18, 19]. In this context we note the following: from an electrodynamics point of view, generating currents by our pulses is a completely classical effect, i.e. currents are generated in a completely classical

system, even though in our case the subsequent excited carrier evolution is quantum mechanical. For this reason our current is robust to disorder and geometry modifications. In addition, the tunable time delay between the pulses allows an ultra-fast control of the current properties. Using circularly polarized laser pulses generates currents for quantized systems (in which case the rotating-wave approximation can be applied). For systems with level broadening of the order of level spacing, no appreciable current is generated. Our disadvantage, however, is that our pulses are much more demanding to realize experimentally, whereas laser pulses are readily available, in particular at high light intensities, thus allowing for a strong current generation.

The DSC is proportional to the DCC which can be comparable to the PCC for the small or moderate occupation number case as seen in figure 4(b). The DCC depends sensitively and dramatically on the strength of the field and the delayed time between the two pulses. We provide now an explicit calculation for the typical values of the CC and SC. For the $\text{In}_x\text{Ga}_{1-x}\text{As}/\text{InP}$ quantum well [42] we have

$m^* = 0.037m_0$. For a ring with radius of 100 nm, the line velocity is then about 5000 m s^{-1} , and the current unit is $I_0/a \sim 8 \text{ nA}$ which corresponds to the angular velocity current for one particle. If we convert this into the unit of an induced magnetization it is a radius-independent quantity $M_0 \approx 2 \text{ meV T}^{-1}$ (here we use the formula $M_0 \approx \pi a^2(I_0/a)$ valid for the rings considered here [17]).

Acknowledgment

The work is supported by the cluster of excellence ‘Nanostructured Materials’ of the state Saxony-Anhalt.

References

- [1] For a review on spintronics, see, e.g., Wolf S A *et al* 2001 *Science* **294** 1488
- [2] Dresselhaus G 1955 *Phys. Rev.* **100** 580
- [3] Rashba E I 1960 *Sov. Phys.—Solid State* **2** 1109
Bychkov Y A and Rashba E I 1984 *J. Phys. C: Solid State Phys.* **17** 6039
- [4] Lommer G *et al* 1988 *Phys. Rev. Lett.* **60** 728
- [5] Schultz M *et al* 1996 *Semicond. Sci. Technol.* **11** 1168
Zhang X C *et al* 2001 *Phys. Rev. B* **63** 245305
Gui Y S *et al* 2004 *Phys. Rev. B* **70** 115328
- [6] Luo J *et al* 1990 *Phys. Rev. B* **41** 7685
- [7] Nitta J *et al* 1997 *Phys. Rev. Lett.* **78** 1335
Hu C-M *et al* 1988 *Phys. Rev. Lett.* **60** 728
Engels G *et al* 1997 *Phys. Rev. B* **55** R1958
Schäpers Th *et al* 1998 *J. Appl. Phys.* **83** 4324
- [8] Malcher F *et al* 1986 *Superlatt. Microstruct.* **2** 267
- [9] Murakami S, Nagaosa N and Zhang S C 2003 *Science* **301** 1348
- [10] Sinova J *et al* 2004 *Phys. Rev. Lett.* **92** 126603
- [11] Fuhrer A *et al* 2001 *Nature* **413** 822
Fuhrer A *et al* 2002 *Microelectron. Eng.* **63** 47
Fuhrer A *et al* 2003 *Phys. Rev. Lett.* **91** 206802
Fuhrer A *et al* 2004 *Phys. Rev. Lett.* **93** 176803
- [12] Warburton R J *et al* 2000 *Nature* **405** 926
Lorke A *et al* 2000 *Phys. Rev. Lett.* **84** 2223
Bayer M *et al* 2003 *Phys. Rev. Lett.* **90** 186801
Keyser U F *et al* 2003 *Phys. Rev. Lett.* **90** 196601
Alén B *et al* 2007 *Phys. Rev. B* **75** 45319
- [13] Nitta J, Meijer F E and Takayanagi H 1999 *Appl. Phys. Lett.* **75** 695
- [14] Nitta J and Koga T 2003 *J. Supercond.* **16** 689
- [15] You D, Jones R R, Bucksbaum P H and Dykaar D R 1993 *Opt. Lett.* **18** 290
Bensky T J, Haeffler G and Jones R R 1997 *Phys. Rev. Lett.* **79** 2018
- [16] Bennett C H and DiVincenzo D P 2000 *Nature* **404** 247
- [17] Matos-Abiague A and Berakdar J 2005 *Phys. Rev. Lett.* **94** 166801
- [18] Pershin Y V and Piermarocchi C 2005 *Phys. Rev. B* **72** 245331
- [19] Räsänen E *et al* 2007 *Phys. Rev. Lett.* **98** 157404
- [20] Zhu Z-G and Berakdar J 2008 *Phys. Rev. B* **77** 235438
- [21] Schliemann J, Egues J C and Loss D 2003 *Phys. Rev. Lett.* **90** 146801
Wang X F and Vasilopoulou P 2005 *Phys. Rev. B* **72** 165336
- [22] Matos-Abiague A and Berakdar J 2004 *Phys. Rev. B* **70** 195338
Matos-Abiague A and Berakdar J 2005 *Europhys. Lett.* **69** 277
- [23] Meijer F E, Morpurgo A F and Klapwijk T M 2002 *Phys. Rev. B* **66** 033107
- [24] Frustaglia D and Richter K 2004 *Phys. Rev. B* **69** 235310
- [25] Molnár B *et al* 2004 *Phys. Rev. B* **69** 155335
Molnár B *et al* 2005 *Phys. Rev. B* **72** 75330
- [26] Földi P *et al* 2005 *Phys. Rev. B* **71** 33309
Földi P *et al* 2006 *Phys. Rev. B* **73** 155325
- [27] Sheng J S and Chang K 2006 *Phys. Rev. B* **74** 235315
- [28] Spletstoesser J, Governale M and Zülicke U 2003 *Phys. Rev. B* **68** 165341
- [29] Mizushima M 1967 *Prog. Theor. Phys. Suppl.* **40** 207
- [30] Oh S and Ryu C M 1995 *Phys. Rev. B* **51** 13441
- [31] Fröhlich J and Studer U M 1993 *Rev. Mod. Phys.* **65** 733
- [32] Ballentine L E 1998 *Quantum Mechanics: a Modern Development* (Singapore: World Scientific) chapter 11
- [33] Loss D and Goldbart P 1991 *Phys. Rev. B* **43** 13762
Loss D and Goldbart P 1992 *Phys. Rev. B* **45** 13544
- [34] Wendler L, Fomin V M and Krokhin A A 1994 *Phys. Rev. B* **50** 4642
- [35] For a review on quantum rings, see, e.g., Aronov A G and Sharvin Yu V 1987 *Rev. Mod. Phys.* **59** 755
Wendler L and Fomin V M 1995 *Phys. Status Solidi b* **191** 409
Eckern U and Schwab P 2002 *J. Low Temp. Phys.* **126** 1291
- [36] Büttiker M, Imry Y and Landauer R 1983 *Phys. Lett. A* **96** 365
- [37] Chakraborty T and Pietiläinen P 1994 *Phys. Rev. B* **50** 8460
- [38] Chandrasekhar V *et al* 1991 *Phys. Rev. Lett.* **67** 3578
- [39] Mailly D *et al* 1993 *Phys. Rev. Lett.* **70** 2020
- [40] Meir Y, Gefen Y and Wohlman O E 1989 *Phys. Rev. Lett.* **63** 798
Wohlman O E *et al* 1992 *Phys. Rev. B* **45** 11890
- [41] Moskalenko A S, Matos-Abiague A and Berakdar J 2006 *Phys. Rev. B* **74** 161303
Moskalenko A S, Matos-Abiague A and Berakdar J 2007 *Europhys. Lett.* **78** 57001
- [42] Schäpers Th *et al* 1998 *J. Appl. Phys.* **83** 4324
- [43] König M *et al* 2006 *Phys. Rev. Lett.* **96** 76804
- [44] Meier F and Loss D 2003 *Phys. Rev. Lett.* **90** 167204
Schütz F, Kollar M and Kopietz P 2003 *Phys. Rev. Lett.* **91** 017205
Sun Q-F *et al* 2003 arXiv:cond-mat/0301402
Nogueira F S and Bennemann K-H 2003 arXiv:cond-mat/0302528
- [45] Moskalenko A S, Matos-Abiague A and Berakdar J 2006 *Phys. Lett. A* **356** 255