

The Percolation Transition in the Zero-Temperature Domany Model

Federico Camia¹ and Charles M. Newman²

Received July 20, 2003

We analyze a deterministic cellular automaton $\sigma^{\cdot} = (\sigma^n; n \geq 0)$ corresponding to the zero-temperature case of Domany's stochastic Ising ferromagnet on the hexagonal lattice \mathbb{H} . The state space $\mathcal{S}_{\mathbb{H}} = \{-1, +1\}^{\mathbb{H}}$ consists of assignments of -1 or $+1$ to each site of \mathbb{H} and the initial state $\sigma^0 = \{\sigma_x^0\}_{x \in \mathbb{H}}$ is chosen randomly with $P(\sigma_x^0 = +1) = p \in [0, 1]$. The sites of \mathbb{H} are partitioned in two sets \mathcal{A} and \mathcal{B} so that all the neighbors of a site x in \mathcal{A} belong to \mathcal{B} and vice versa, and the discrete time dynamics is such that the σ_x^{\cdot} 's with $x \in \mathcal{A}$ (respectively, \mathcal{B}) are updated simultaneously at odd (resp., even) times, making σ_x^{\cdot} agree with the majority of its three neighbors. In ref. 1 it was proved that there is a percolation transition at $p = 1/2$ in the percolation models defined by σ^n , for all times $n \in [1, \infty]$. In this paper, we study the nature of that transition and prove that the critical exponents β , ν , and η of the dependent percolation models defined by σ^n , $n \in [1, \infty]$, have the same values as for standard two-dimensional independent site percolation (on the triangular lattice).

KEY WORDS: Dependent percolation; critical exponents; universality; cellular automaton; zero-temperature dynamics.

1. INTRODUCTION

The deterministic cellular automaton corresponding to the zero-temperature case of Domany's stochastic Ising ferromagnet on the hexagonal lattice $\mathbb{H}^{(2)}$ can be considered as a simplified version of a continuous time Markov process where an independent (rate 1) Poisson clock is assigned to

¹Forschungsinstitut für Mathematik, ETH, 8092 Zürich, Switzerland. Present address: EURANDOM, P.O. Box 513, 5600 MB Eindhoven, The Netherlands; e-mail: camia@eurandom.tue.nl

²Courant Institute of Mathematical Sciences, New York University, New York, New York 10012; e-mail: newman@courant.nyu.edu

each site $x \in \mathbb{H}$, and the spin σ_x at site x is updated when the corresponding clock rings. The rule for updating the spin is to flip it if and only if it disagrees with two or three (a majority) of its neighbors. This model has been studied both rigorously and numerically in ref. 3; the numerical results about two critical exponents obtained there strongly suggest that the dependent percolation model defined by the limiting spin configuration σ^∞ is in the same universality class as ordinary independent percolation.

In the zero-temperature Domany model studied here, the rule for updating the spins is unchanged, but the timing is different. This model has been previously used in numerical simulations,⁽⁴⁾ and its dynamical as well as percolation properties have been studied in refs. 1, 5, and 6. In ref. 1, the existence of a percolation transition for the dependent percolation models associated with the spin configurations σ^n at time n , for all $n \in [1, \infty]$, is proved; in ref. 5, it is shown that for any $n \geq 1$, the crossing probabilities converge to Cardy's formula⁽⁷⁾ when the lattice spacing δ is sent to zero (the *continuum scaling limit*), as in the case of independent percolation (at least on the triangular lattice \mathbb{T} ⁽⁸⁾); in ref. 6, the continuum scaling limit is analyzed in terms of cluster boundaries and is shown to be the same as for ordinary independent critical percolation on \mathbb{T} . This last result strongly suggests that also the critical exponents defined at the critical point should be the same as for independent percolation. Indeed, due to the above mentioned result, one can use properties of the Stochastic Loewner Evolution⁽⁹⁾ (which describes the scaling limit of ordinary critical percolation cluster interfaces) to compute certain critical exponents in the continuous model corresponding to the scaling limit. But unfortunately, the connection between critical exponents in the continuous and discrete models is not straightforward; more work and further results on the discrete model would be required to relate the discrete critical exponents with the continuous ones (see ref. 10).

Similar models on different lattices have been studied in various papers; see, for example, refs. 11–17 for models on \mathbb{Z}^d and ref. 18 for a model on the homogeneous tree of degree three. Such models are also discussed extensively in the physics literature, usually on \mathbb{Z}^d (see, for example, refs. 2 and 19). Their interest is tied to the fact that they can be obtained as zero-temperature limits of stochastic Ising models, a special class of Markov processes whose transition probabilities/rates are chosen so that the Gibbs measures (for some Hamiltonian) at temperature T are invariant for the Markov process. In systems where there are multiple (infinite-volume) Gibbs measures for T below some critical T_c , a subject of considerable interest is the $t \rightarrow \infty$ behavior of the spin configuration σ^t (with temperature $T = T_1 < T_c$) when the initial state is chosen from the (unique) Gibbs measure at $T = T_2 > T_c$. Studying the limiting case where $T_1 = 0$ and

$T_2 = \infty$ is the standard choice in much of the statistical physics literature (see, e.g., ref. 20).

In this paper, we study the percolation properties of (say) $+1$ spins in the dependent percolation models, generated by the zero-temperature Domany cellular automaton, corresponding to σ^n , with $n \in [1, \infty]$, when the initial state is chosen randomly with $P(\sigma_x^0 = +1) = p \in [0, 1]$. In ref. 1, it was shown that there is a percolation transition with $p_c = 1/2$ for all values of $n \in [1, \infty]$. We remark that p_c is *not* $1/2$ for $n = 0$, since the critical probability for independent site percolation on the hexagonal lattice is strictly larger than $1/2$, therefore the system is *driven* to criticality by the dynamics, and this after just one time step. This is in contrast to what happens in the case of the continuous time model studied in ref. 3, where it is believed that criticality is achieved again for $p = 1/2$, but only at time $t = \infty$. Nonetheless, the nature of the percolation transition, in terms of critical exponents and/or scaling limits, is presumed to be the same in the different models.

2. DEFINITION OF THE MODEL AND PRELIMINARY RESULTS

In this section, we give a more detailed description of the model and present, for completeness, results that were proved in ref. 1 which motivate (and will be used in) the next section, where the main results of this paper are presented.

Consider the homogeneous ferromagnet on the hexagonal lattice \mathbb{H} (embedded in \mathbb{R}^2 so that the elementary cells are regular hexagons with side length 1—see, for example, Fig. 1) with states denoted by $\sigma = \{\sigma_x\}_{x \in \mathbb{H}}$, $\sigma_x = \pm 1$, and with (formal) Hamiltonian

$$\mathcal{H}(\sigma) = - \sum_{\langle x, y \rangle} \sigma_x \sigma_y, \quad (1)$$

where $\sum_{\langle x, y \rangle}$ denotes the sum over all pairs of neighbor sites, each pair counted once. We write $\mathcal{N}_{\mathbb{H}}(x)$ for the set of three neighbors of x , and indicate with

$$\Delta_x \mathcal{H}(\sigma) = 2 \sum_{y \in \mathcal{N}_{\mathbb{H}}(x)} \sigma_x \sigma_y \quad (2)$$

the change in the Hamiltonian when the *spin* σ_x at site x is flipped (i.e., changes sign).

Partition the sites of the hexagonal lattice \mathbb{H} into two subsets, \mathcal{A} and \mathcal{B} , in such a way that all three neighbors of a site x in \mathcal{A} (resp., \mathcal{B}) are in \mathcal{B} (resp., \mathcal{A}). By joining two sites of \mathcal{A} whenever they are next-nearest




Fig. 1. A star-triangle transformation relating triangular and hexagonal lattice.

neighbors in the hexagonal lattice (two steps away from each other), we get a triangular lattice (the same with \mathcal{B} —see Fig. 1). The synchronous dynamics that we consider here is such that all the sites in \mathcal{A} (resp., \mathcal{B}) are updated simultaneously.

We now define the cellular automaton σ^n , $n \in \mathbb{N}$, with state space $\mathcal{S}_{\mathbb{H}} = \{-1, +1\}^{\mathbb{H}}$, which is the zero temperature limit of a model of Domany,⁽²⁾ as follows:

- The initial state σ^0 is chosen from a Bernoulli product measure P_p , with $P_p(\sigma_0^0 = +1) = p$.
- At odd times $n = 1, 3, \dots$, the spins in \mathcal{A} are updated according to the following rule: σ_x , $x \in \mathcal{A}$, is flipped if and only if $\Delta_x \mathcal{H}(\sigma) < 0$.
- At even times $n = 2, 4, \dots$, the spins in \mathcal{B} are updated according to the same rule as for those in \mathcal{A} .

We denote by σ^∞ the limiting state of the cellular automaton σ^n . $\sigma^\infty = \lim_{n \rightarrow \infty} \sigma^n$ exists with probability one⁽¹⁴⁾ and, like σ^n for $1 \leq n < \infty$, defines a dependent percolation model on \mathbb{H} .

The following observations are useful in understanding the behavior of the model and will help in the proof of our main result, Theorem 3, which is presented in the next section of the paper.

- The values of the spins in \mathcal{A} at time 0 are irrelevant since after the first update those values are uniquely determined by the values of the spins in \mathcal{B} .
- Once the initial spin configuration in \mathcal{B} is chosen, the dynamics is completely deterministic.

- A spin can no longer flip once it belongs to either a loop or a “barbell” of constant sign, where a barbell consists of two disjoint loops connected by a path.

There is an alternative but equivalent way of describing the discrete time dynamics as a deterministic cellular automaton (with random initial state) on the triangular lattice \mathbb{T} (corresponding to the set \mathcal{B} of sites). Given some site $\bar{x} \in \mathbb{T}$, group its six \mathbb{T} -neighbors $y_i^{\bar{x}}$ in three disjoint pairs $\{y_1^{\bar{x}}, y_2^{\bar{x}}\}$, $\{y_3^{\bar{x}}, y_4^{\bar{x}}\}$, $\{y_5^{\bar{x}}, y_6^{\bar{x}}\}$, so that $y_1^{\bar{x}}$ and $y_2^{\bar{x}}$ are \mathbb{T} -neighbors, and so on for the other two pairs. Translate this construction to all sites $x \in \mathbb{T}$, thus producing three pairs of sites $\{y_1^x, y_2^x\}$, $\{y_3^x, y_4^x\}$, $\{y_5^x, y_6^x\}$ associated to each site $x \in \mathbb{T}$. (Note that this construction does not need to specify how \mathbb{T} is embedded in \mathbb{R}^2 .) Site x is updated at times $m = 1, 2, \dots$ according to the following rule: the spin at site x is changed from σ_x to $-\sigma_x$ if and only if at least two of its pairs of neighbors have all four sites with the same sign $-\sigma_x$.

The models on the hexagonal and on the triangular lattice are related through a *star-triangle transformation* (see Fig. 1 and, for example, p. 335 of ref. 22). More precisely, the dynamics on the triangular lattice \mathbb{T} is equivalent to the zero-temperature Domany dynamics on the hexagonal lattice \mathbb{H} when restricted to the sublattice \mathcal{B} for even times $n = 2m$.

To see this, start with \mathbb{T} and construct an hexagonal lattice \mathbb{H}' by means of a star-triangle transformation such that a site is added at the center of each of the triangles (x, y_1^x, y_2^x) , (x, y_3^x, y_4^x) , and (x, y_5^x, y_6^x) . \mathbb{H}' may be partitioned into two triangular sublattices \mathcal{A}' and \mathcal{B}' with $\mathcal{B}' = \mathbb{T}$. One can now see that the dynamics on \mathbb{T} for $m = 1, 2, \dots$ and the zero-temperature Domany dynamics on \mathbb{H}' restricted to \mathcal{B}' for even times $n = 2m$ are the same.

An immediate consequence of this equivalence between the two cellular automata is that the two families of percolation models that they produce are also equivalent in an obvious way through a star-triangle transformation. To be more precise, the percolation models defined on \mathbb{T} for times $m = 1, 2, \dots$ are the same as those defined on \mathcal{B} by the zero-temperature Domany model for even times $n = 2m$.

We now present (without proof) the results of ref. 1 on the zero-temperature Domany model. Theorem 1 says that the convergence to the limiting state is exponentially fast, while Theorem 2 identifies the critical point of the percolation transition, $p_c = 1/2$.

Theorem 1. Let $P_{\mathcal{A}}(n)$ denote the probability that a deterministic site in \mathcal{A} flips after time n and similarly for $P_{\mathcal{B}}(n)$. Then, for any p , there is a constant $c \in (0, \infty)$ such that

$$P_{\mathcal{A}}(n), P_{\mathcal{B}}(n) \leq e^{-cn}. \quad (3)$$

Theorem 2. If $p > 1/2$ (resp., $< 1/2$), there is percolation of $+1$ (resp., -1) spins in σ^n for any $n \in [1, \infty]$ for almost every σ^0 .

If $p = 1/2$:

1. For $n \in [0, \infty]$, there is no percolation in σ^n of either $+1$ or -1 spins, for almost every σ^0 .
2. The mean cluster size is infinite in σ^n for all $n \in [1, \infty]$ (but finite for $n = 0$).

3. CRITICAL EXPONENTS

We will consider three percolation critical exponents, namely the exponents β (related to the percolation probability), ν (related to the correlation length) and η (related to the connectivity function). The existence of these exponents has been proved, and their predicted values confirmed rigorously, in recent papers,^(10,21) for the case of independent site percolation on the triangular lattice. Such exponents are believed to be universal for independent percolation in the sense that their value should depend only on the number of dimensions and not on the structure of the lattice or on the nature of the percolation model (e.g., whether it is site or bond percolation); that type of universality has not yet been proved.

Consider an independent percolation model with distribution P_p on a two-dimensional lattice \mathbb{L} such that $0 < p_c < 1$. Let C_x be the open cluster containing site x and $|C_x|$ be its cardinality, then $\theta(p) = \theta_x(p) = P_p(|C_x| = \infty)$ is the *percolation probability*. Arguments from theoretical physics suggest that $\theta(p)$ behaves roughly like $(p - p_c)^\beta$ as p approaches p_c from above.

It is also believed that the *connectivity function*

$$\tau_p(x, y) = P_p(x \text{ and } y \text{ belong to the same cluster}) \quad (4)$$

behaves, for the Euclidean length $\|x - y\|$ large, like $\|x - y\|^{-\eta}$ if $p = p_c$, and like $\exp(-\|x - y\|/\xi(p))$ if $0 < p < p_c$, for some $\xi(p)$ satisfying $\xi(p) \rightarrow \infty$ as $p \uparrow p_c$. The *correlation length* $\xi(p)$ is defined by

$$\xi(p)^{-1} = \lim_{\|x-y\| \rightarrow \infty} \left\{ -\frac{1}{\|x-y\|} \log \tau_p(x, y) \right\}. \quad (5)$$

$\xi(p)$ is expected to behave like $(p_c - p)^{-\nu}$ as $p \uparrow p_c$.

It is not clear how strong one may expect such asymptotic relations to be (for more details about critical exponents and *scaling theory* in percolation, see ref. 22 and references therein). For this reason, the logarithmic

relation is usually employed. That means that the previous conjectures are usually stated in the following form:

$$\lim_{p \downarrow p_c} \frac{\log \theta(p)}{\log(p - p_c)} = \beta, \tag{6}$$

$$\lim_{\|x-y\| \rightarrow \infty} \frac{\log \tau_{p_c}(x, y)}{\log \|x - y\|} = -\eta, \tag{7}$$

$$\lim_{p \uparrow p_c} \frac{\log \zeta(p)}{\log(p_c - p)} = -\nu. \tag{8}$$

In the rest of this section and in the next one, $\theta_x(p)$, $\tau_p(x, y)$, and $\zeta(p)$ will indicate the percolation probability, connectivity function and correlation length for independent site percolation on the triangular lattice \mathbb{T} (identified with \mathcal{B}). We will denote by $\theta_x(p, n)$, $\tau_{p,n}(x, y)$, and $\zeta(p, n)$ the corresponding quantities for the percolation models on the hexagonal lattice \mathbb{H} at time $n = 0, 1, 2, \dots$. The main theorem of this paper is the following.

Theorem 3. There exist constants $0 < c_1, c_2, c_3, c_4 < \infty$ such that, $\forall n \in [1, \infty]$, and $x, y \in \mathbb{H}$ and suitably chosen $x', y' \in \mathcal{B}$ with $\|x - x'\|, \|y - y'\| \leq 1$,

$$c_1 \theta_{x'}(p) \leq \theta_x(p, n) \leq c_2 \theta_{x'}(p), \quad \text{for } p \in (1/2, 1], \tag{9}$$

$$p^{c_3} \tau_p(x', y') \leq \tau_{p,n}(x, y) \leq p^{-c_4} \tau_p(x', y'), \quad \text{for } p \in (0, 1/2], \tag{10}$$

$$\zeta(p, n) = \zeta(p), \quad \text{for } p \in (0, 1/2]. \tag{11}$$

The next corollary is an immediate consequence of Theorem 3 and its main application; it says that the dependent percolation models defined by σ^n , with $n \in [1, \infty]$, are in the universality class of ordinary independent percolation.

Corollary 3.1. The critical exponents β , η , and ν exist for the dependent percolation models on \mathbb{H} defined by σ^n , with $n \in [1, \infty]$, and have the same numerical values as for independent site percolation on \mathbb{T} .

Remark 3.1. As already mentioned, the existence of the exponents β , ν , and η for independent site percolation on \mathbb{T} has been recently proved, and their predicted values confirmed rigorously.^(10, 21)

4. PROOFS

Before we can prove the main results of this paper, we need some notation. We will denote by $P_{p,n}$ the distribution of σ^n with initial density p of plus spins (i.e., with initial distribution $P_{p,0} = P_p$).

For a site $x \in \mathbb{H}$, we will denote by $\mathcal{N}_{\mathbb{H}}(x)$ the set of its three neighbors in \mathbb{H} . For a site $x \in \mathcal{B}$, we will denote by $\mathcal{N}_{\mathcal{B}}(x)$ the set of its six neighbors in \mathcal{B} endowed with the graph structure of a triangular lattice (as explained in Section 2—see also Fig. 1).

We will call an \mathbb{H} -path a path on the hexagonal lattice \mathbb{H} and a \mathcal{B} -path a path on the triangular lattice \mathcal{B} . A path whose spins are all plus (resp., minus) will be called a plus (resp., minus) path. Similarly, we will call an \mathbb{H} -loop a (simple) loop on the hexagonal lattice \mathbb{H} and \mathcal{B} -loop a (simple) loop on the triangular lattice \mathcal{B} . A loop whose spins are all plus (resp., minus) will be called a plus (resp., minus) loop. Notice that constant-sign \mathbb{H} -loops, doubly-infinite \mathbb{H} -paths and “barbells” (two loops connected by a path) are stable for the dynamics, in the sense that, once formed, their spins will never flip again.

For two subsets C and D of \mathbb{H} , we indicate with $\{C \overset{\mathbb{H}}{\longleftrightarrow} D\}$ the event that some site in C is connected to some site in D by a *plus* \mathbb{H} -path, with $\{C \overset{\mathbb{H}}{\longleftrightarrow} \infty\}$ the event that some site in C belongs to an infinite *plus* \mathbb{H} -path. For two subsets C and D of \mathcal{B} , endowed with the graph structure of a triangular lattice, we indicate with $\{C \overset{\mathcal{B}}{\longleftrightarrow} D\}$ the event that some site in C is connected to some site in D by a *plus* \mathcal{B} -path, with $\{C \overset{\mathcal{B}}{\longleftrightarrow} \infty\}$ the event that some site in C belongs to an infinite *plus* \mathcal{B} -path.

Proof of Theorem 3. Let us first assume that x and y belong to \mathcal{B} ; we then take $x' = x, y' = y$ in (9)–(10). The lower bound for $\theta_x(p, n)$ in Eq. (9) comes from the following bound

$$\theta_x(p, n) = P_{p,n}(x \overset{\mathbb{H}}{\longleftrightarrow} \infty) \quad (12)$$

$$\geq P_{p,0}(\{x \overset{\mathcal{B}}{\longleftrightarrow} \infty\} \cap \{x \text{ belongs to a plus } \mathbb{H}\text{-loop}\}) \quad (13)$$

$$\geq P_{p,0}(x \overset{\mathcal{B}}{\longleftrightarrow} \infty) P_{p,0}(x \text{ belongs to a plus } \mathbb{H}\text{-loop}) \quad (14)$$

$$\geq p^6 \theta_x(p) \quad (15)$$

$$> \frac{1}{2^6} \theta_x(p), \quad (16)$$

where we have used the fact that at time 1 (when the sites in \mathcal{A} are updated for the first time) the dynamics transforms any constant sign \mathcal{B} -path into a constant sign \mathbb{H} -path, the FKG inequality and the fact that the events

$\{0 \overset{\mathcal{B}}{\leftrightarrow} \infty\}$ and $\{0 \text{ belongs to a plus } \mathbb{H}\text{-loop}\}$ are increasing (see, for example, ref. 22), and the fact that the smallest \mathbb{H} -loop contains 6 sites.

The lower bound for $\tau_{p,n}(x, y)$ in Eq. (10) is obtained in a similar way. Mimicking the proof of the lower bound in Eq. (9), we have immediately

$$\tau_{p,n}(x, y) = P_{p,n}(x \overset{\mathbb{H}}{\leftrightarrow} y) \geq p^{12} \tau_p(x, y). \tag{17}$$

For the upper bound of Eq. (9) we rely on the following observation. If no site in $\mathcal{N}_{\mathcal{B}}(x)$ belongs to an infinite plus \mathcal{B} -path at time 0, then, by the self-matching property of the triangular lattice, site x must be surrounded by a minus \mathcal{B} -loop, which will produce a stable \mathbb{H} -loop at time 1. Therefore, site x will not belong to an infinite plus \mathbb{H} -path at any later time. Thus,

$$\theta_x(p, n) \leq P_{p,0}(\mathcal{N}_{\mathcal{B}}(x) \overset{\mathcal{B}}{\leftrightarrow} \infty). \tag{18}$$

Since the event $\{x \overset{\mathcal{B}}{\leftrightarrow} \infty\}$ can be written as $\{\sigma_x^0 = +1\} \cap \{\mathcal{N}_{\mathcal{B}}(x) \overset{\mathcal{B}}{\leftrightarrow} \infty\}$, using the FKG inequality we have

$$P_{p,0}(x \overset{\mathcal{B}}{\leftrightarrow} \infty) \geq p P_{p,0}(\mathcal{N}_{\mathcal{B}}(x) \overset{\mathcal{B}}{\leftrightarrow} \infty). \tag{19}$$

From this and Eq. (18), we get

$$\theta_x(p, n) \leq p^{-1} \theta_x(p) \leq 2\theta_x(p), \tag{20}$$

as required.

For the upper bound of Eq. (10), we first note that for bounded $\|x - y\|$, the inequality is trivial by choosing c_4 big enough so that the right-hand side of (10) exceeds 1. Next, for $\|x - y\|$ large enough, we notice that unless $\{\mathcal{N}_{\mathcal{B}}(x) \overset{\mathcal{B}}{\leftrightarrow} \mathcal{N}_{\mathcal{B}}(y)\}$ at time 0, x and y must be separated by a minus \mathcal{B} -loop surrounding one of them or by a doubly-infinite minus \mathcal{B} -path, and therefore it cannot be the case that $\{x \overset{\mathbb{H}}{\leftrightarrow} y\}$ at any later time because at time 1 a stable minus \mathbb{H} -loop or doubly-infinite \mathbb{H} -path will be formed. This yields, for $\|x - y\|$ large enough,

$$\tau_{p,n}(x, y) \leq P_{p,0}(\mathcal{N}_{\mathcal{B}}(x) \overset{\mathcal{B}}{\leftrightarrow} \mathcal{N}_{\mathcal{B}}(y)). \tag{21}$$

Since the event $\{x \overset{\mathcal{B}}{\leftrightarrow} y\}$ can be written as $\{\sigma_x^0 = \sigma_y^0 = +1\} \cap \{\mathcal{N}_{\mathcal{B}}(x) \overset{\mathcal{B}}{\leftrightarrow} \mathcal{N}_{\mathcal{B}}(y)\}$, using the FKG inequality we have

$$P_{p,0}(x \overset{\mathcal{B}}{\leftrightarrow} y) \geq p^2 P_{p,0}(\mathcal{N}_{\mathcal{B}}(x) \overset{\mathcal{B}}{\leftrightarrow} \mathcal{N}_{\mathcal{B}}(y)). \tag{22}$$

From this and Eq. (21), we get

$$\tau_{p,n}(x, y) \leq p^{-2} \tau_p(x, y), \tag{23}$$

as required.

If x and y belong to \mathcal{A} , the proof is analogous, but one has to consider slightly different events. In this case we take $x' \in \mathcal{N}_{\mathbb{H}}(x)$ and then we have, for the lower bound in Eq. (9),

$$\theta_x(p, n) = P_{p, n}(x \overset{\mathbb{H}}{\leftrightarrow} \infty) \quad (24)$$

$$\geq P_{p, 0}(\{x' \overset{\mathcal{B}}{\leftrightarrow} \infty\} \cap \{x \text{ belongs to a plus } \mathbb{H}\text{-loop}\}) \quad (25)$$

$$\geq P_{p, 0}(x' \overset{\mathcal{B}}{\leftrightarrow} \infty) P_{p, 0}(x \text{ belongs to a plus } \mathbb{H}\text{-loop}) \quad (26)$$

$$\geq p^6 \theta_{x'}(p) \quad (27)$$

$$> \frac{1}{2^6} \theta_{x'}(p), \quad (28)$$

where $\|x - x'\| = 1$ because x' is an \mathbb{H} -neighbor of x .

Analogously, for the lower bound in Eq. (10), we get

$$\tau_{p, n}(x, y) = P_{p, n}(x \overset{\mathbb{H}}{\leftrightarrow} y) \geq p^{12} \tau_p(x', y'), \quad (29)$$

with $x' \in \mathcal{N}_{\mathbb{H}}(x)$ and $y' \in \mathcal{N}_{\mathbb{H}}(y)$.

For the upper bound of Eq. (9), we notice that, if no site in $\bigcup_{z \in \mathcal{N}_{\mathbb{H}}(x)} \mathcal{N}_{\mathcal{B}}(z)$ belongs to an infinite plus \mathcal{B} -path at time 0, then, by the self-matching property of the triangular lattice, site x must be surrounded by a minus \mathcal{B} -loop that will produce a stable \mathbb{H} -loop at time 1. Therefore, site x will not belong to an infinite plus \mathbb{H} -path at any later time. Thus,

$$\theta_x(p, n) \leq P_{p, 0} \left(\bigcup_{z \in \mathcal{N}_{\mathbb{H}}(x)} \mathcal{N}_{\mathcal{B}}(z) \overset{\mathcal{B}}{\leftrightarrow} \infty \right). \quad (30)$$

Since for $x' \in \mathcal{N}_{\mathbb{H}}(x)$, $\{x' \overset{\mathcal{B}}{\leftrightarrow} \infty\} \supset \{\sigma_z^0 = +1, \forall z \in \mathcal{N}_{\mathbb{H}}(x)\} \cap \{\bigcup_{z \in \mathcal{N}_{\mathbb{H}}(x)} \mathcal{N}_{\mathcal{B}}(z) \overset{\mathcal{B}}{\leftrightarrow} \infty\}$, using the FKG inequality we have

$$P_{p, 0}(x' \overset{\mathcal{B}}{\leftrightarrow} \infty) \geq p^3 P_{p, 0} \left(\bigcup_{z \in \mathcal{N}_{\mathbb{H}}(x)} \mathcal{N}_{\mathcal{B}}(z) \overset{\mathcal{B}}{\leftrightarrow} \infty \right), \quad (31)$$

which yields

$$\theta_x(p, n) \leq p^{-3} \theta_{x'}(p) \leq 8 \theta_{x'}(p), \quad (32)$$

with $\|x - x'\| = 1$.

The upper bound of Eq. (10), still trivial for bounded $\|x - y\|$, follows again from a similar observation for $\|x - y\|$ large enough: unless $\{\bigcup_{z \in \mathcal{N}_{\mathbb{H}}(x)} \mathcal{N}_{\mathcal{B}}(z) \overset{\mathcal{B}}{\leftrightarrow} \bigcup_{z' \in \mathcal{N}_{\mathbb{H}}(y)} \mathcal{N}_{\mathcal{B}}(z')\}$ at time 0, x and y must be separated

by a minus \mathcal{B} -loop surrounding one of them or by a doubly-infinite minus \mathcal{B} -path. At time 1, a stable minus \mathbb{H} -loop or doubly-infinite \mathbb{H} -path will be formed, making the event $\{x \overset{\mathbb{H}}{\leftrightarrow} y\}$ impossible at any later time. This yields, for $\|x - y\|$ large enough,

$$\tau_{p,n}(x, y) \leq P_{p,0} \left(\bigcup_{z \in \mathcal{N}_{\mathbb{H}}(x)} \mathcal{N}_{\mathcal{B}}(z) \overset{\mathcal{B}}{\leftrightarrow} \bigcup_{z' \in \mathcal{N}_{\mathbb{H}}(y)} \mathcal{N}_{\mathcal{B}}(z') \right). \tag{33}$$

Since for $x' \in \mathcal{N}_{\mathbb{H}}(x)$ and $y' \in \mathcal{N}_{\mathbb{H}}(y)$, $\{x' \overset{\mathcal{B}}{\leftrightarrow} y'\} \supset \{\sigma_z^0 = +1, \forall z \in \mathcal{N}_{\mathbb{H}}(x)\} \cap \{\sigma_{z'}^0 = +1, \forall z' \in \mathcal{N}_{\mathbb{H}}(y)\} \cap \{\bigcup_{z \in \mathcal{N}_{\mathbb{H}}(x)} \mathcal{N}_{\mathcal{B}}(z) \overset{\mathcal{B}}{\leftrightarrow} \bigcup_{z' \in \mathcal{N}_{\mathbb{H}}(y)} \mathcal{N}_{\mathcal{B}}(z')\}$, using the FKG inequality we have

$$P_{p,0}(x' \overset{\mathcal{B}}{\leftrightarrow} y') \geq p^6 P_{p,0} \left(\bigcup_{z \in \mathcal{N}_{\mathbb{H}}(x)} \mathcal{N}_{\mathcal{B}}(z) \overset{\mathcal{B}}{\leftrightarrow} \bigcup_{z' \in \mathcal{N}_{\mathbb{H}}(y)} \mathcal{N}_{\mathcal{B}}(z') \right), \tag{34}$$

which yields

$$\tau_{p,n}(x, y) \leq p^{-6} \tau_p(x', y'), \tag{35}$$

with $x' \in \mathcal{N}_{\mathbb{H}}(x)$ and $y' \in \mathcal{N}_{\mathbb{H}}(y)$.

The proof of Eq. (10) in the remaining case (namely, $x \in \mathcal{A}$, $y \in \mathcal{B}$ or vice versa) should now be clear.

Equation (11) is an immediate consequence of Eq. (10) and the definition of $\xi(p)$; it is enough to observe that

$$\lim_{\|x-y\| \rightarrow \infty} \left\{ -\frac{1}{\|x'-y'\|} [\log \tau_p(x', y') + c_3 \log p] \right\} = \xi(p)^{-1} \tag{36}$$

and

$$\lim_{\|x-y\| \rightarrow \infty} \left\{ -\frac{1}{\|x'-y'\|} [\log \tau_p(x', y') - c_4 \log p] \right\} = \xi(p)^{-1}. \blacksquare \tag{37}$$

Proof of Corollary 3.1. It follows from Eqs. (9) and (10) that, for $p \in (1/2, 1]$ and $\|x - y\|$ large enough,

$$\frac{\log c_1 + \log \theta_{x'}(p)}{\log(p-1/2)} \leq \frac{\log \theta_x(p, n)}{\log(p-1/2)} \leq \frac{\log c_2 + \log \theta_{x'}(p)}{\log(p-1/2)}, \tag{38}$$

$$\frac{\log \tau_{1/2}(x', y') + c_3 \log \frac{1}{2}}{\log \|x - y\|} \leq \frac{\log \tau_{1/2,n}(x, y)}{\log \|x - y\|} \leq \frac{\log \tau_{1/2}(x', y') - c_4 \log \frac{1}{2}}{\log \|x - y\|}. \tag{39}$$

Using these two equations, together with Eq. (11) and the definitions of the critical exponents, and taking the appropriate limits gives the desired results. \blacksquare

ACKNOWLEDGMENTS

Research partially supported by the U.S. NSF under Grant DMS-01-04278.

REFERENCES

1. F. Camia, C. M. Newman, and V. Sidoravicius, Approach to fixation for zero-temperature stochastic Ising models on the hexagonal lattice, in *In and Out of Equilibrium: Probability with a Physics Flavor*, V. Sidoravicius, ed., Progress in Probability, Vol. 51 (Birkhäuser, 2002), pp. 163–183.
2. E. Domany, Exact results for two- and three-dimensional Ising and Potts models, *Phys. Rev. Lett.* **52**:871–874 (1984).
3. C. D. Howard and C. M. Newman, The percolation transition for the zero-temperature stochastic Ising model on the hexagonal lattice, *J. Stat. Phys.* **111**:57–72 (2003).
4. B. Nienhuis, private communication (2001).
5. F. Camia, C. M. Newman, and V. Sidoravicius, Cardy's formula for some dependent percolation models, *Bull. Brazilian Math. Soc.* **33**:147–156 (2002).
6. F. Camia, C. M. Newman, and V. Sidoravicius, A particular bit of universality: Scaling limits of some dependent percolation models, *Commun. Math. Phys.*, to appear.
7. J. L. Cardy, Critical percolation in finite geometries, *J. Phys. A* **25**:L201–L206 (1992).
8. S. Smirnov, Critical percolation in the plane: Conformal invariance, Cardy's formula, scaling limits, *C. R. Acad. Sci. Paris* **333**:239–244 (2001).
9. O. Schramm, Scaling limits of loop-erased random walks and uniform spanning trees, *Israel J. Math.* **118**:221–288 (2000).
10. S. Smirnov and W. Werner, Critical exponents for two-dimensional percolation, *Math. Rev. Lett.* **8**:279–744 (2001).
11. F. Camia, E. De Santis, and C. M. Newman, Clusters and recurrence in the two-dimensional zero-temperature stochastic Ising model, *Ann. Appl. Probab.* **12**:565–580 (2002).
12. L. R. Fontes, R. H. Schonmann, and V. Sidoravicius, Stretched exponential fixation in stochastic Ising models at zero temperature, *Commun. Math. Phys.* **228**:495–518 (2002).
13. A. Gandolfi, C. M. Newman, and D. L. Stein, Zero-temperature dynamics of $\pm J$ spin glasses and related models, *Commun. Math. Phys.* **214**:373–387 (2000).
14. S. Nanda, C. M. Newman, and D. L. Stein, Dynamics of Ising spin systems at zero temperature, in *On Dobrushin's Way (from Probability Theory to Statistical Mechanics)*, R. Minlos, S. Shlosman, and Y. Suhov, eds. (AMS, Providence, 2000).
15. C. M. Newman and D. L. Stein, Blocking and persistence in zero-temperature dynamics of homogeneous and disordered Ising models, *Phys. Rev. Lett.* **82**:3944–3947 (1999).
16. C. M. Newman and D. L. Stein, Equilibrium pure states and nonequilibrium chaos, *J. Stat. Phys.* **94**:709–722 (1999).
17. C. M. Newman and D. L. Stein, Zero-temperature dynamics of Ising spin systems following a deep quench: Results and open problems, *Physica A* **279**:156–168 (2000).
18. C. D. Howard, Zero-temperature Ising spin dynamics on the homogeneous tree of degree three, *J. Appl. Probab.* **37**:736–747 (2000).
19. J. L. Lebowitz, C. Maes, and E. R. Speer, Statistical mechanics of probabilistic cellular automata, *J. Stat. Phys.* **59**:117–170 (1990).
20. A. J. Bray, Theory of phase ordering kinetics, *Adv. Phys.* **43**:565–580 (1994).
21. G. R. Grimmett, *Percolation*, second edition (Springer, Berlin, 1999).
22. G. F. Lawler, O. Schramm, and W. Werner, One-arm exponent for critical 2D percolation, *Electr. J. Probab.* **7**:000–000 (2002).