

Large Scale Dynamics of the Persistent Turning Walker Model of Fish Behavior

Pierre Degond · Sébastien Motsch

Received: 4 October 2007 / Accepted: 14 March 2008 / Published online: 1 April 2008
© Springer Science+Business Media, LLC 2008

Abstract This paper considers a new model of individual displacement, based on fish motion, the so-called Persistent Turning Walker (PTW) model, which involves an Ornstein-Uhlenbeck process on the curvature of the particle trajectory. The goal is to show that its large time and space scale dynamics is of diffusive type, and to provide an analytic expression of the diffusion coefficient. Two methods are investigated. In the first one, we compute the large time asymptotics of the variance of the individual stochastic trajectories. The second method is based on a diffusion approximation of the kinetic formulation of these stochastic trajectories. The kinetic model is a Fokker-Planck type equation posed in an extended phase-space involving the curvature among the kinetic variables. We show that both methods lead to the same value of the diffusion constant. We present some numerical simulations to illustrate the theoretical results.

Keywords Individual based model · Fish behavior · Persistent Turning Walker model · Ornstein-Uhlenbeck process · Kinetic Fokker-Planck equation · Asymptotic analysis · Diffusion approximation

1 Introduction

This paper considers a new model of individual displacement, the so-called ‘Persistent Turning Walker’ (PTW) model, which has recently been introduced to describe fish behavior [28]. The fish evolves with a velocity of constant magnitude and its trajectory is subject to random turns (i.e. random changes of curvature) on the one hand and to curvature relaxation to zero on the other hand. The random changes of curvature can be interpreted as a way for the fish to explore its surroundings while relaxation to zero curvature just expresses that the

P. Degond · S. Motsch (✉)

Institute of Mathematics of Toulouse UMR 5219 (CNRS-UPS-INSA-UT1-UT2), Université Paul Sabatier, 118, route de Narbonne, 31062 Toulouse cedex, France
e-mail: motsch@mip.ups-tlse.fr

P. Degond
e-mail: degond@mip.ups-tlse.fr

fish cannot sustain too strongly curved trajectories and that, when the curvature becomes too large, the fish tries to return to a straight line trajectory. The combination of these two antagonistic behaviors gives rise to an Ornstein-Uhlenbeck process on the curvature. The curvature is the time derivative of the director of the velocity, while the velocity itself is the time derivative of position. The PTW process collects all these considerations into a system of stochastic differential equations.

This model is, to the knowledge of the authors, original, and has appeared for the first time in the works by Gautrais, Theraulaz, and coworkers [28]. It has been deduced from a statistical analysis of data issued from experiments run in La Réunion islands during years 2001 and 2002. The studied species is a pelagic fish named *Kuhlia Mugil*. Its typical size ranges between 20 and 25 cm. The first experiments have been made with a single fish in a basin of 4 meters diameter during two minutes. A video records the positions of the fish every 12-th of a second (see Fig. 1 below). Then, after a filtering of the oscillatory swimming motion of the fish, the trajectories have been fitted to model trajectories.

The more classical process consisting of a fish moving along straight lines and undergoing random changes of its velocity direction according to a Poisson process (like a photon in a diffusive medium) has been discarded. Indeed, statistical tests have shown [28] that the befitting of the actual trajectory of a fish to a broken line interpolating the trajectory at random points did not lead to a satisfactory agreement. On the other hand, a better fitting was obtained when the trajectory was interpolated by a sequence of circular arcs connected to each other with a continuous tangent. It was then observed that the curvatures of these circles followed a process which can be viewed as a realization of an Ornstein-Uhlenbeck process on the curvature. This was the starting point of the PTW process. The experiments and the data analysis are reported in full detail in [28].

Then, the experiments were reproduced with 2, 5, 15 and 30 fish in the basin. The analyses of the interactions among the fish population are still in progress. In the present work, we will concentrate on the modeling of a single fish behaviour based on the PTW model and we will postpone the modeling of interacting individuals to future time, when the analysis of the data will be available. However, we will have in mind that the models issued from the analysis of a single fish behaviour must be easily expanded to model large groups of interacting fish.

The present paper considers the large time and space scale dynamics of a two-dimensional particle subject to this PTW process. It rigorously shows (in the mathematical sense) that, at large scales, the dynamics of the particle can be described by a diffusion process and it provides a formula for the diffusion coefficient. To prove this result, two methods are considered.

In the first method, the stochastic differential system itself is considered and the variance of the position is shown to behave, at large times, like a linear function of time. The diffusion coefficient is identified as the slope of this linear function. Because the curvature and the velocity angle can be explicitly computed, an explicit formula for the diffusion coefficient, involving some special functions, can be obtained.

The second method considers the forward Kolmogorov equation of the stochastic process. This equation gives the evolution of the probability distribution function of the particle in the extended phase space (position, velocity angle, curvature) as a function of time. It is a Fokker-Planck type equation. The passage from the microscopic to the macroscopic scales relies on a rescaling of the Kolmogorov equation. This rescaling depends on a small parameter $\varepsilon \ll 1$, which describes the ratio of the typical microscopic to macroscopic space units. After this rescaling, the problem has the typical form of the diffusion approximation of a kinetic problem (see references below). The goal is then to study the behaviour

of the solution as $\varepsilon \rightarrow 0$. It is shown that the solution converges to some ‘thermodynamical equilibrium’ which is a Gaussian distribution of the curvature and a uniform distribution of the velocity angle. The equilibrium depends parametrically on the density which satisfies a spatial diffusion equation.

Finally, the connection between the two methods is made by showing that the diffusion tensor in the second approach can be represented by a formula involving the solution of the stochastic differential equation of the first approach. Additionally, this representation leads to explicit computations which show that the two formulas for the diffusion coefficient actually coincide. This seemingly innocuous result is actually quite powerful. Indeed, the diffusion approximation method leads to a non-explicit expression of the diffusion coefficient, involving the moments of a particular solution of a stationary equation involving the leading order operator of the Fokker-Planck equation. That this non-explicit formula is equivalent to the explicit formula given by the stochastic trajectory method is by far not obvious. In this respect, the stochastic trajectory method is more powerful than the diffusion approximation approach, because it directly leads to the most simple expression of the diffusion constant.

A third route could have been taken and has been dismissed. This third method would actually use the stochastic differential equation itself to perform the diffusion approximation in the forward Kolmogorov equation. We have preferred to use partial differential equation techniques. One reason for this choice is that these techniques can be more easily extended to more complex situations. One typical example of these more complex situations are the nonlinear systems which are obtained when interactions between individual are included. The inclusion of interactions between individuals within the PTW model is actually work in progress.

From the biological viewpoint, one should not restrict the content of the paper to the sole expression of the diffusion coefficient. Indeed, once interactions between individuals will be included in the PTW model, it is not clear at all that the explicit computations which led to this expression will still be tractable. In the absence of an explicit solution of the stochastic differential system, there is little grasp to get information about the large scale behaviour of the system. By contrast, the diffusion approximation approach gives a systematic tool to study the large scale behavior of such systems, in all kinds of situations, be they linear or nonlinear. By its flexibility and its versatility, the diffusion approximation approach is the method of choice to study these problems.

The ultimate goal of this work programme is the establishment of a macroscopic model for large groups of fish. However, such a model must be based on a reliable model for the individual displacements, since interactions lead to an alteration of the individual behaviour in the absence of interaction. Therefore, the present work, which has a solid experimental basis, can be viewed as the first step towards the establishment of such an experimentally based model.

One previously proposed models to describe fish behavior is the discrete Vicsek model (VM) [1, 14, 31, 51] (see also [2, 10, 25, 27, 39, 42, 43] for related models). For a large scale modeling of fish behavior, it is efficient to look at continuum models, which use macroscopic variables such as mean density, mean velocity and so on. Several such models based on phenomenological observations, exist (see e.g. [26, 38, 49, 50]). Several attempts to derive continuum models from the VM model are also reported in the literature [36, 46, 47]. In [19, 20], a derivation of a continuum model from a kinetic version of the VM model is proposed. However, few Individual Based Models for fish have been validated against experimental data with a comparable care as in [28] for the PTW process. As such, the continuum model derived in this paper has a firm experimental basis, although further work needs certainly to be done to fully validate its biological foundations. Additional references on swarm aggregation and fish schooling can be found in [11]. Among other types of animal societies,

insects, and in particular ants [35, 48] or cockroaches [34] have been the subject of a vast literature (see references therein).

The derivation of macroscopic models from particle behavior has been initiated by the seminal works of Boltzmann, and later Hilbert, Chapman and Enskog. We refer to [13] for a mathematical perspective and to [16] for an introduction to the subject from a modeling perspective. More recently, the derivation of macroscopic models from microscopic behavior has been very productive in other context like traffic [4, 33] or supply-chains [3]. Diffusion approximation problems for kinetic equations have been widely studied in the literature, in the context of neutron transport (see e.g. [6, 8]), semiconductors [7, 21, 29, 45], plasmas [17, 18, 22] or polymeric fluids [23].

The outline of the paper is as follows: in Sect. 2, the PTW process is introduced and the main results are stated. In Sect. 3 the diffusion coefficient is obtained by direct inspection of the trajectories of the stochastic differential system. In Sect. 4, the diffusion approximation of the forward Kolmogorov equation of the stochastic process is performed. Section 5 is devoted to proving that the trajectory method and the diffusion approximation method give rise to the same value of the diffusion coefficient. In Sect. 6, the theoretical results are illustrated by and complemented with some numerical simulations. A conclusion is drawn in Sect. 7. Several proofs of auxiliary results, which are inessential for the main discussion are collected in three appendix (A, B and C).

2 The Persistent Turning Walker Model: Presentation and Main Results

The starting point of the present work is a new model of fish motion based on experimental data taken from experiments run in La Réunion islands [28] (see a schematic description of the experiments in the introduction). Figure 1 displays a typical trajectory of a fish as recorded during the experiment.

The statistical analysis of the data recorded during these experiments have led to the observation that the trajectories are well described by the following systems of stochastic differential equations:

$$\frac{d\mathbf{x}}{dt} = c\boldsymbol{\tau}(\theta), \quad (2.1)$$

$$\frac{d\theta}{dt} = c\kappa, \quad (2.2)$$

$$d\kappa = -a\kappa dt + b dB_t, \quad (2.3)$$

where $\mathbf{x} = (x_1, x_2) \in \mathbb{R}^2$ is the (two-dimensional) position vector of the (centroid of the) fish, $\boldsymbol{\tau}(\theta) = (\cos \theta, \sin \theta)$ is the director of the velocity vector with the angle $\theta \in \Pi = \mathbb{R}/2\pi\mathbb{Z}$ measured from the x_1 direction, $\kappa \in \mathbb{R}$ is the curvature of the trajectory and dB_t is the standard Brownian motion. The magnitude of the velocity is constant and denoted by $c > 0$. The constant a is a relaxation frequency and b quantifies the intensity of the random curvature jumps. b has the dimension of $1/(L\sqrt{T})$ where L and T stand for the dimensions of length and time.

The κ -dynamics is a standard Ornstein-Uhlenbeck process. The term “ $b dB_t$ ” models a diffusion process in curvature space while the term “ $-a \kappa dt$ ” expresses the tendency of the individual to return to a straight line trajectory. The curvature cannot increase endlessly as a consequence of the diffusion process, but rather, must relax to zero and the relaxation is stronger as the curvature gets larger. This model has been called the *Persistent Turning*




Fig. 1 One trajectory of a *Kuhlia Mugil* fish

Walker model (PTW) because it allows large excursions of the curvature towards positive or negative values, during which the spinning of the trajectory persists for a certain time.

We stress the difference with more standard diffusion processes (such as those suffered by photons in a diffusive medium), in which the Brownian motion acts on the velocity itself (or, in the case of a velocity of constant magnitude, on the angle θ). In this case, the diffusion process acts on the second derivative of the particle positions, and the associated kinetic equation is of Fokker-Planck type. This model of photon diffusion is also relevant for a certain number of animal species [41].

In the PTW model, the diffusion process acts on the curvature, i.e. on the third derivative of the position vector. An intuitive justification of the relevance of this model for animal behaviour is by considering the non-differentiability of the Brownian motion. Because of this feature, the photon diffusion process involves infinite second derivatives of the position, i.e. infinite forces. However, an animal body can only exert finite forces and the muscles act only in such a way that the velocity angle undergoes smooth variations. The PTW model precisely presents this feature of having smooth second order derivatives, i.e. smooth forces.

Our goal in the present work is to study the large-scale dynamics of the stochastic differential system (2.1)–(2.3). This is best done in scaled variables, where the dimensionless parameters of the model are highlighted. We use $t_0 = a^{-1}$ as time unit, $x_0 = ca^{-1}$ as space unit, and $\kappa_0 = x_0^{-1}$ as curvature unit, and we introduce the dimensionless time, space and curvature as $t' = t/t_0$, $x' = x/x_0$ and $\kappa' = \kappa/\kappa_0$. For simplicity, we omit the primes. In

scaled variables, the PTW model is written:

$$\frac{d\mathbf{x}}{dt} = \boldsymbol{\tau}(\theta), \tag{2.4}$$

$$\frac{d\theta}{dt} = \kappa, \tag{2.5}$$

$$d\kappa = -\kappa dt + \sqrt{2\alpha} dB_t, \tag{2.6}$$

where the only dimensionless parameter left is α such that

$$\alpha^2 = \frac{b^2 c^2}{2a^3}. \tag{2.7}$$

The meaning of α^2 is the following: b/\sqrt{a} is the amplitude of a curvature change during a relaxation time a^{-1} , while c/a is obviously the distance travelled by the particle during this time. The product of these two quantities is dimensionless and is equal to $\sqrt{2\alpha}$. It quantifies the strength of the curvature jumps relative to the other phenomena.

The individual dynamics can be translated in terms of a probability distribution $f(t, \mathbf{x}, \theta, \kappa) d\mathbf{x} d\theta d\kappa$ of finding particles at times t with position in small neighborhoods $d\mathbf{x} d\theta d\kappa$ of position \mathbf{x} , velocity angle θ and curvature κ . The link between the individual dynamics and the evolution of the probability distribution f is given by the forward Kolmogorov equation:

$$\partial_t f + \boldsymbol{\tau} \cdot \nabla_{\mathbf{x}} f + \kappa \partial_{\theta} f - \partial_{\kappa}(\kappa f) - \alpha^2 \partial_{\kappa^2} f = 0. \tag{2.8}$$

This equation is an exact transcription of the individual dynamics, where the initial value f_0 at time $t = 0$ is given by the probability distribution of the initial conditions of the stochastic differential system (2.4)–(2.6). For more detailed considerations about the forward Kolmogorov equation and its link with stochastic differential systems, we refer the reader to [5, 40].

In order to capture the macroscopic dynamics, two possible routes can be taken, using either the stochastic differential system (2.4)–(2.6) or the partial differential equation (2.8). In this work, we follow both routes and verify that they lead to the same large-scale behaviour. The advantage of working directly on the stochastic system is that it is simpler and it leads to explicit formulas. However, as soon as the system gets more complicated, and in particular nonlinear, explicit solutions can no longer be found and this methodology can hardly be pursued. On the other hand, the PDE approach, which, in the present case is more complicated, is also more systematic and more general. In particular, it is generally usable in the more complex nonlinear cases (see e.g. [19, 20]). A particular important complex situation is the case of many interacting fish. In future work, we plan to extend the PTW model to populations of interacting fish and to use the PDE approach to extract the large-scale dynamics of the system.

From the analysis of the individual trajectories, explicit exact expressions for κ and θ in terms of stochastic integrals can be found. Unfortunately, there is no such explicit result for the position $\mathbf{x}(t)$, but we can calculate the first two moments of the probability distribution of $\mathbf{x}(t)$ explicitly, using the expressions of κ and θ . We show that the mean of the position vector stays at the origin: $\mathbb{E}\{\mathbf{x}(t)\} = (0, 0)$ (where \mathbb{E} denotes the expectation over all sources of randomness, in the initial data and in the stochastic process) and that the variance grows asymptotically linearly in time. More exactly, we prove:

Theorem 2.1 *Under assumptions on the initial conditions that will be specified later on (see (3.1)–(3.4)), the solution of system (2.4)–(2.6) satisfies:*

$$\text{Var}\{\mathbf{x}(t)\} \stackrel{t \rightarrow +\infty}{\sim} 2\mathcal{D}t, \quad \text{with } \mathcal{D} = \int_0^\infty \exp(-\alpha^2(-1 + s + e^{-s})) ds. \tag{2.9}$$

The notation Var is for the variance over all sources of randomness. The asymptotic linear growth of the variance (2.9) suggests that the dynamics of the system is of diffusive type at large times with diffusion coefficient \mathcal{D} . We can find an expression of \mathcal{D} in terms of special functions. Indeed, we have

Proposition 2.1 *The following expression holds true:*

$$\mathcal{D} = \left(\frac{e}{\alpha^2}\right)^{\alpha^2} \gamma(\alpha^2, \alpha^2), \tag{2.10}$$

where $\gamma(z, u)$ is the incomplete gamma function:

$$\gamma(z, u) = \int_0^u e^{-t} t^{z-1} dt. \tag{2.11}$$

\mathcal{D} has the following series representation:

$$\mathcal{D} = e^{\alpha^2} \sum_{n=0}^\infty \frac{(-1)^n \alpha^{2n}}{n!(n + \alpha^2)}. \tag{2.12}$$

It is a decreasing function of α which has the following asymptotic behavior:

$$\mathcal{D} \sim \frac{1}{\alpha^2} \quad \text{as } \alpha \rightarrow 0, \quad \mathcal{D} \sim \sqrt{\frac{\pi}{2}} \frac{1}{\alpha} \quad \text{as } \alpha \rightarrow \infty. \tag{2.13}$$

To investigate the large scale dynamics of the solution of the kinetic equation (2.8) (the existence of which can be easily proved, see Proposition 4.2), we need to rescale the variables to the macroscopic scale. Indeed, in (2.8), all the coefficients are supposed to be of order unity. This means that the time and space scales of the experiment are of the same order as the typical time and length scales involved in the dynamics, such as, the relaxation time or the inverse of the typical random curvature excursions. Of course, in most experiments, this is not true, since the duration of the experiment and the size of the experimental region are large compared with the time and length scales involved in the dynamics.

To translate this observation, we change the space unit x_0 to a new space unit $x'_0 = x_0/\varepsilon$, where $\varepsilon \ll 1$ is a small parameter. This induces a change of variables $x' = \varepsilon x$. We make a similar operation on the time unit $t'_0 = t_0/\eta$, $t' = \eta t$ with $\eta \ll 1$. Now, the question of linking η to ε is a subtle one and is largely determined by the nature of the asymptotic regime which is achieved by the system. In the present case, we expect that the asymptotic regime will be of diffusive nature, in view of Theorem 2.1 and so, we will investigate the so-called ‘diffusion approximation’ which involves a quadratic relationship between η and ε : $\eta = \varepsilon^2$.

For this reason, we introduce the diffusive rescaling:

$$t' = \varepsilon^2 t; \quad \mathbf{x}' = \varepsilon \mathbf{x}, \tag{2.14}$$

and we make the following change of variable in the distribution f :

$$f^\varepsilon(t', \mathbf{x}', \theta, \kappa) = \frac{1}{\varepsilon^2} f\left(\frac{t'}{\varepsilon^2}, \frac{\mathbf{x}'}{\varepsilon}, \theta, \kappa\right).$$

The scaling of the magnitude of the distribution function is unnecessary, since the problem is linear. However, it is chosen in order to preserve the total number of particles. Introducing (2.14) into (2.8) leads to the following problem for f^ε :

$$\varepsilon \partial_t f^\varepsilon + \boldsymbol{\tau} \cdot \nabla_{\mathbf{x}} f^\varepsilon + \frac{1}{\varepsilon} [\kappa \partial_\theta f^\varepsilon - \partial_\kappa(\kappa f^\varepsilon) - \alpha^2 \partial_{\kappa^2} f^\varepsilon] = 0. \tag{2.15}$$

In order to analyze the large-scale dynamics of (2.15), we need to investigate the limit $\varepsilon \rightarrow 0$. We show that f^ε converges to an equilibrium distribution function (i.e. a function which cancels the $O(\varepsilon^{-1})$ term of (2.15)) f^0 which depends parametrically on the particle density $n^0(x, t)$ and n^0 evolves according to a diffusion equation. More precisely, we prove:

Theorem 2.2 *Under Hypothesis 4.1 on the initial data to be precised below, the solution f^ε of (2.15) converge weakly in a Banach space also to be specified below, (see (4.21)) X :*

$$f^\varepsilon \xrightarrow{\varepsilon \rightarrow 0} n^0 \frac{M(\kappa)}{2\pi} \text{ in } X \text{ weak star}, \tag{2.16}$$

where M is a Gaussian distribution of the curvature with zero mean and variance α^2 (see (4.4)) and $n^0 = n^0(x, t)$ is the solution of the system:

$$\partial_t n^0 + \nabla_{\mathbf{x}} \cdot \mathbf{J}^0 = 0, \tag{2.17}$$

$$\mathbf{J}^0 = -D \nabla_{\mathbf{x}} n^0, \tag{2.18}$$

where the initial datum n_0^0 and the diffusion tensor D will be defined later on (see (4.31) and (4.28) respectively).

The following theorem connects the two methods by showing that the tensor D is related to \mathcal{D} given by (2.9):

Theorem 2.3 *The tensor D defined by (4.28) satisfies:*

$$D = \frac{\mathcal{D}}{2} Id, \tag{2.19}$$

where \mathcal{D} is given by (2.9) and Id denotes the 2×2 identity tensor.

This theorem confirms that the trajectory method and the asymptotic PDE method are equivalent. The factor 2 between the two coefficients comes from the dimension of the problem. Indeed, \mathcal{D} is the average of $|x|^2 = |x_1|^2 + |x_2|^2$ while D is the diffusion coefficient in a given direction.

The graphical representation of \mathcal{D} is given in Fig. 2. The expression \mathcal{D}_0 of the diffusion coefficient in ‘physical’ variables is obtained by multiplying the dimensionless expression \mathcal{D} by the appropriate scaling units. The scaling unit for a diffusion coefficient is the square

of the space scale divided by the time scale. Therefore, in the present case, its value is c^2/a . Therefore, we find:

$$\mathcal{D}_0 = \frac{c^2}{a} \mathcal{D} \left(\frac{b^2 c^2}{2a^3} \right). \tag{2.20}$$

Expression (2.20) and the fact that $\mathcal{D}(\alpha)$ is a decreasing function of α shows that the diffusion coefficient is decreasing with respect to b for fixed a and c . This is explained by the fact that, with an increasing noise intensity, the trajectory has a larger probability to reach strong curvatures (in magnitude). Therefore, the trajectory spins more, and the distance travelled in straight line is shorter.

On the other hand, the monotonicity of \mathcal{D}_0 with respect to a and c is unclear. Still, we can investigate the asymptotic limits and find that for $c \rightarrow \infty$ or $b \rightarrow \infty$ or $a \rightarrow 0$, (each limit being taken with the other two parameters kept fixed), we have $\mathcal{D}_0 \sim \sqrt{\pi} \sqrt{ac}/b$. In particular, for large c or small a , the diffusion increases with respect to both c and a . Conversely, if $c \rightarrow 0$ or $b \rightarrow 0$ or $a \rightarrow \infty$, (again, each limit being taken with the other two parameters kept fixed), we have $\mathcal{D}_0 \sim 2a^2/b^2$. Here, the diffusion is increasing with a but is independent of c .

The increase of \mathcal{D}_0 with a is easily explained: with a stronger relaxation parameter a , the curvature is more likely to be small, and the trajectory resembles more a straight line.

On the other hand, the diffusion \mathcal{D}_0 is independent of the velocity c when c is small. This is somehow paradoxical since one would expect that, as the particle moves faster, it travels larger distances. However, as c increases, the spinning of the trajectory increases, because the particle moves along a circle before undergoing a random change of curvature or a relaxation. Therefore, the average linear distance from the origin does not increase so much when the velocity is increased, at least for small velocities. For large velocities, the intuitive feeling that the diffusion should increase with the velocity is actually true: the diffusion is asymptotically proportional c for large values of c .

3 Large-Scale Dynamics of the PTW Model by the Trajectory Method

In this section, we want to show Theorem 2.1 and Proposition 2.1. We first specify the initial conditions. First, we fix the starting point of the particle at the origin:

$$\mathbf{x}(t = 0) = (0, 0). \tag{3.1}$$

We suppose that the initial velocity angle is uniformly distributed on the one-dimensional sphere, i.e.:

$$dP\{\theta|_{t=0} = \theta\} = \frac{d\theta}{2\pi}. \tag{3.2}$$

For the curvature, we make the following observation: (2.6) predicts that the process $\kappa(t)$ converges exponentially fast to its stationary state, which is a Gaussian distribution with zero mean and variance equal to α^2 [40]. We denote such a Gaussian distribution by $\mathcal{N}(0, \alpha^2)$. For this reason, we suppose:

$$dP\{\kappa|_{t=0} = \kappa\} = \mathcal{N}(0, \alpha^2)(\kappa). \tag{3.3}$$




Fig. 2 Scaled diffusion coefficient \mathcal{D} as a function of α^2 in log-log scale (solid line). Asymptotic behavior of \mathcal{D} as $\alpha \rightarrow 0$ or $\alpha \rightarrow \infty$ (dashed lines): $\mathcal{D} \sim \alpha^{-2}$ as $\alpha \rightarrow 0$ and $\mathcal{D} \sim \sqrt{\frac{\pi}{2}}\alpha^{-1}$ as $\alpha \rightarrow \infty$

The last hypothesis on the initial conditions is the following:

$$\text{The processes } \theta(t = 0), \kappa(t = 0) \text{ and } B_t \text{ are independents.} \tag{3.4}$$

We stress that this choice of initial conditions is for simplicity only. Completely arbitrary initial conditions would lead to the same large time behaviour, but the computation would be slightly more complicated. Since we are mainly interested in the explicit expression of \mathcal{D} , a choice of initial conditions which simplifies the calculations is legitimate.

We begin by proving the following proposition:

Proposition 3.1 *The solution of the stochastic differential equation (2.4)–(2.6) with initial condition given by (3.1)–(3.4) satisfies:*

$$\mathbb{E}\{\mathbf{x}(t)\} = (0, 0), \quad \forall t \geq 0, \tag{3.5}$$

$$\text{Var}\{\mathbf{x}(t)\} = 2 \int_{s=0}^t (t - s) \exp(-\alpha^2 (-1 + s + e^{-s})) ds. \tag{3.6}$$

To prove this proposition, we first establish explicit formulae for the solutions of (2.5) and (2.6). The proof is deferred to Appendix A.

Lemma 3.1 *The solution of the stochastic differential system (2.5), (2.6) with initial conditions (3.2)–(3.4) is given by:*

$$\theta(t) = \theta_0 + \kappa_0 - \kappa(t) + \sqrt{2}\alpha B_t, \tag{3.7}$$

$$\kappa(t) = e^{-t}\kappa_0 + \sqrt{2}\alpha e^{-t} \int_0^t e^s dB_s. \tag{3.8}$$

Additionally,

$$\theta(t) = \theta_0 + K_0^t, \tag{3.9}$$

where K_0^t is a Gaussian random variable independent of θ_0 with zero mean and variance β_t^2 given by:

$$\beta_t^2 = \text{Var}\{K_0^t\} = 2\alpha^2(-1 + t + e^{-t}). \tag{3.10}$$

Proof of Proposition 3.1 Using Lemma 3.1, we can compute the first two moments of $\mathbf{x}(t)$. Let us start with the computation of the mean. If we write $\mathbf{x}(t) = (x_1(t), x_2(t))$, we have:

$$x_1(t) = \int_0^t \cos \theta(s) ds, \quad x_2(t) = \int_0^t \sin \theta(s) ds,$$

and, computing the mean:

$$\mathbb{E}\{x_1(t)\} = \mathbb{E}\left\{ \int_0^t \cos \theta(s) ds \right\} = \int_0^t \mathbb{E}\{\cos \theta(s)\} ds.$$

Now, we can develop $\theta(s)$ using (3.9):

$$\mathbb{E}\{\cos \theta(s)\} = \mathbb{E}\{\cos(\theta_0 + K_0^s)\} = \mathbb{E}\{\cos \theta_0 \cos K_0^s - \sin \theta_0 \sin K_0^s\}.$$

By the independence of θ_0 and K_0^s we finally have:

$$\mathbb{E}\{\cos \theta(s)\} = \mathbb{E}\{\cos \theta_0\}\mathbb{E}\{\cos K_0^s\} - \mathbb{E}\{\sin \theta_0\}\mathbb{E}\{\sin K_0^s\} = 0,$$

since the expectations of $\cos \theta_0$ and $\sin \theta_0$ over the uniform probability distribution on θ_0 are zero. Finally, we have $\mathbb{E}\{x_1(t)\} = 0$, and similarly for x_2 . This proves (3.5).

Now for the variance of $\mathbf{x}(t)$, we write:

$$\text{Var}\{\mathbf{x}(t)\} = \mathbb{E}\{x_1^2(t) + x_2^2(t)\} = 2\mathbb{E}\{x_1^2(t)\} \tag{3.11}$$

by the isotropy of the problem. Then,

$$\begin{aligned} \mathbb{E}\{x_1^2(t)\} &= \mathbb{E}\left\{ \left(\int_0^t \cos \theta(s) ds \right)^2 \right\} \\ &= \int_0^t \int_0^t \mathbb{E}\{\cos \theta(s) \cos \theta(u)\} ds du \\ &= 2 \int_0^t du \int_0^u ds \mathbb{E}\{\cos \theta(s) \cos \theta(u)\}. \end{aligned}$$

Since $u \geq s$, we can write $\theta(u)$ as follows:

$$\theta(u) = \theta_0 + \int_0^s \kappa(z) dz + \int_s^u \kappa(z) dz = \theta_0 + K_0^s + K_s^u,$$

where K_0^s and K_s^u are Gaussian random variables independent of θ_0 with zero mean and variances β_s^2 and β_{u-s}^2 respectively, thanks to (3.10). Then, using standard identities for trigonometric functions, we get

$$\begin{aligned} \mathbb{E}\{\cos \theta(s) \cos \theta(u)\} &= \mathbb{E}\{\cos(\theta_0 + K_0^s) \cos(\theta_0 + K_0^s + K_s^u)\} \\ &= \frac{1}{2}(\cos(2\theta_0 + 2K_0^s + K_s^u) + \cos(-K_s^u)). \end{aligned}$$

But since θ_0 is independent of K_0^s and K_s^u we have $\mathbb{E}\{\cos(2\theta_0 + 2K_0^s + K_s^u)\} = 0$ since the mean of a $\cos(\theta_0 + C)$ over the uniform distribution of θ_0 is zero whatever the value of C . Then:

$$\begin{aligned} \mathbb{E}\{\cos \theta(s) \cos \theta(u)\} &= \frac{1}{2} \mathbb{E}\{\cos(-K_s^u)\} \\ &= \frac{1}{2} \int_{\mathbb{R}} \cos(y) \frac{1}{\sqrt{2\pi} \beta_{u-s}} e^{-\frac{y^2}{2\beta_{u-s}^2}} dy \\ &= \frac{1}{2} e^{-\frac{1}{2}\beta_{u-s}^2}. \end{aligned}$$

Indeed, an elementary computation shows that for any Gaussian random variable Z with zero mean and variance σ^2 , one has

$$\mathbb{E}\{\cos(Z)\} = \exp(-\sigma^2/2). \tag{3.12}$$

Thus,

$$\mathbb{E}\{x_1(t)^2\} = \int_{u=0}^t \int_{s=0}^u \exp(-\alpha^2(-1 + |u - s| + e^{-|u-s|})) ds du.$$

Using the change of unknowns $w = u - s$ and $y = u$ and inverting the order of integration we find:

$$\mathbb{E}\{x_1(t)^2\} = \int_{w=0}^t (t - w) \exp(-\alpha^2(-1 + w + e^{-w})) dw.$$

Using (3.11), we finally find (3.6), which ends the proof of the proposition. □

In order to prove 2.1, we investigate the behavior of the variance $\text{Var}\{x(t)\}$ (given by (3.6)) when $t \rightarrow +\infty$.

End of proof of Theorem 2.1 We write, thanks to (3.6):

$$\text{Var}\{x(t)\} - 2Dt = -2 \int_{s=0}^t s e^{-\alpha^2(-1+s+e^{-s})} ds - 2 \int_{s=t}^{\infty} t e^{-\alpha^2(-1+s+e^s)} ds.$$

We have to show that the difference is bounded independently of t . For the first term, we have:

$$\left| \int_{s=0}^t s e^{-\alpha^2(-1+s+e^{-s})} ds \right| \leq \int_0^t e^{\alpha^2} s e^{-\alpha^2 s} ds,$$

and integrating by parts, we find:

$$\left| \int_{s=0}^t s e^{-\alpha^2(-1+s+e^s)} ds \right| \leq \frac{e^{\alpha^2}}{\alpha^2} \left[-t e^{-\alpha^2 t} - \frac{e^{-\alpha^2 t}}{\alpha^2} + \frac{1}{\alpha^2} \right] \leq C_1.$$

For the second term, we have:

$$\left| \int_{s=t}^\infty t e^{-\alpha^2(-1+s+e^s)} ds \right| \leq t \int_t^\infty e^{\alpha^2} e^{-\alpha^2 s} ds \leq t e^{\alpha^2} \frac{e^{-t\alpha^2}}{\alpha^2} \leq C_2.$$

This proves that the difference is $\text{Var}\{x(t)\} - 2\mathcal{D}t$ is bounded independently of t and completes the proof. □

We now prove Proposition 2.1 which gives an explicit approximation of the diffusion coefficient. This approximation is useful for practical simulations.

Proof of Proposition 2.1 The change of variables $t = \alpha^2 e^{-s}$ in the integral (2.9) leads to (2.10). The series representation (2.12) follows from a similar series representation of the incomplete gamma function (see e.g. formula (8.354) of [30]). The series representation can also be found by expanding the exponential in the integral (2.11) in power series (this point is left to the reader). That \mathcal{D} is a decreasing function of α follows from (2.9) and the fact that the function $g(s) = -1 + s + \exp(-s)$ is non-negative for $s \geq 0$. The behavior of \mathcal{D} for $\alpha \rightarrow 0$ (first formula (2.13)) is obtained by keeping only the zero-th order term in the series expansion (2.12). From (2.9), the behaviour of \mathcal{D} for $\alpha \rightarrow \infty$ is controlled by the behaviour of $g(s)$ near $s = 0$. Since $g(s) \sim s^2/2$, we find $\mathcal{D} \sim \int_0^\infty \exp(-\alpha^2 s^2/2) ds$, which leads to the second formula (2.13), and ends the proof. □

4 Large-Scale Dynamics of the PTW Model Through the Diffusion Approximation of the Associated Kinetic Equation

4.1 Formal Asymptotics

In this section, for the reader's convenience, we give a formal proof of Theorem 2.2. We write (2.15) as follows:

$$\varepsilon \partial_t f^\varepsilon + \boldsymbol{\tau} \cdot \nabla_x f^\varepsilon + \frac{1}{\varepsilon} A f^\varepsilon = 0, \tag{4.1}$$

where we define the operator A acting on functions $u(\theta, \kappa)$ as follows:

$$Au = \kappa \partial_\theta u - \partial_\kappa (\kappa u) - \alpha^2 \partial_{\kappa^2} u. \tag{4.2}$$

The formal investigation of the limit $\varepsilon \rightarrow 0$ usually starts by considering the Hilbert expansion (see e.g. [16] for the general theory or [18] for an application in the context of Fokker-Planck equations):

$$f^\varepsilon = f^0 + \varepsilon f^1 + O(\varepsilon^2), \tag{4.3}$$

with f^k being independent of ε and inserting it into (4.1). Then, collecting all the terms of comparable orders with respect to ε , we are led to a sequence of equations. The first one, corresponding to the leading $O(\varepsilon^{-1})$ term is $Af^0 = 0$, which means that f^0 lies in the kernel of A . In Sect. 4.3, we show that the kernel of A is composed of functions of the form $f^0(t, \mathbf{x}, \theta, \kappa) = n^0(t, \mathbf{x})M(\kappa)/(2\pi)$ where $M(\kappa)$ is a normalized Gaussian with zero mean and variance α^2 :

$$M(\kappa) = \frac{1}{\sqrt{2\pi}\alpha^2} e^{-\frac{\kappa^2}{2\alpha^2}}, \tag{4.4}$$

and $n^0(t, \mathbf{x})$ is a function still to be determined.

In order to determine n^0 , we first integrate (4.1) with respect to $(\theta, \kappa) \in \Pi \times \mathbb{R}$ and use that $\int Au \, d\theta \, d\kappa = 0$. Defining the density $n^\varepsilon(t, \mathbf{x})$ and the flux $J^\varepsilon(t, \mathbf{x})$ by

$$n^\varepsilon(t, \mathbf{x}) = \int_{\theta, \kappa} f^\varepsilon \, d\kappa \, d\theta, \quad J^\varepsilon(t, \mathbf{x}) = \int_{\theta, \kappa} \frac{f^\varepsilon}{\varepsilon} \boldsymbol{\tau}(\theta) \, d\kappa \, d\theta, \tag{4.5}$$

we find:

$$\partial_t n^\varepsilon + \nabla_x \cdot J^\varepsilon = 0. \tag{4.6}$$

We note that this continuity equation is valid for all values of ε . Then, letting $\varepsilon \rightarrow 0$, we formally have $n^\varepsilon \rightarrow n^0$. If we prove that J^0 given by (2.18) is the limit of J^ε , as $\varepsilon \rightarrow 0$, then, we can pass to the limit in (4.6) and find (2.17).

System (2.17) and (2.18) is a diffusion system, which completely determines $n^0(t, \mathbf{x})$, given its initial datum $n_0^0(\mathbf{x})$. Here, for simplicity, we assume that the initial datum for (4.1) is of the form $f^\varepsilon(0, \mathbf{x}, \theta, \kappa) = n_0(\mathbf{x})M(\kappa)/(2\pi)$ and the resulting initial condition for n^0 is therefore $n_0^0 = n_0$ (in this formal convergence proof, we admit the functions and the convergences are as smooth as required).

So, the only points left in the proof are the existence of a limit for J^ε and the validity of (2.18) for J^0 . Note that the existence of a limit is not obvious because of the factor ε at the denominator of the integral (4.5) defining J^ε . To prove that the limit exists, we use the Hilbert expansion (4.3) again and compute f^1 . Since A is linear, collecting the terms of order $O(\varepsilon^0)$ leads to:

$$-Af^1 = \frac{M(\kappa)}{2\pi} \boldsymbol{\tau} \cdot \nabla_x n^0. \tag{4.7}$$

Again using the linearity of A and the fact that it operates only with respect to the (θ, κ) variables, we can write the solution of (4.7) as $f^1 = -\boldsymbol{\chi} \cdot \nabla_x n^0$, where $\boldsymbol{\chi} = (\chi_1, \chi_2)$ is a solution of the problem

$$A\boldsymbol{\chi} = \frac{M(\kappa)}{2\pi} \boldsymbol{\tau}. \tag{4.8}$$

This equation must be understood componentwise (i.e χ_1 is associated with $\tau_1 = \cos \theta$ and χ_2 with $\tau_2 = \sin \theta$). Since the right-hand side of (4.8) has zero average with respect to (θ, κ) , Proposition 4.3 below shows that it has a unique solution, up to an element of the kernel of A . We can single out a unique solution by requesting that $\boldsymbol{\chi}$ has zero average with respect to (θ, κ) as well. Then, all solutions f^1 to (4.7) can be written as

$$f^1 = -\boldsymbol{\chi} \cdot \nabla_x n^0 + n^1(t, \mathbf{x}) \frac{M(\kappa)}{2\pi}, \tag{4.9}$$

where the second term of (4.9) is an arbitrary element of the kernel of A . We shall see that the determination of n^1 is unnecessary.

Now, inserting the Hilbert expansion (4.3) into the integral (4.5) defining J^ε , we find:

$$\begin{aligned} J^\varepsilon(t, \mathbf{x}) &= \frac{1}{\varepsilon} \int_{\theta, \kappa} f^0 \boldsymbol{\tau}(\theta) d\kappa d\theta + \int_{\theta, \kappa} f^1 \boldsymbol{\tau}(\theta) d\kappa d\theta + O(\varepsilon) \\ &= 0 + \int_{\theta, \kappa} f^1 \boldsymbol{\tau}(\theta) d\kappa d\theta + O(\varepsilon), \end{aligned} \tag{4.10}$$

because f^0 is independent of θ and $\int \boldsymbol{\tau}(\theta) d\theta = 0$. Therefore, J^ε has a limit when $\varepsilon \rightarrow 0$ and this limit is given by

$$J^0(t, \mathbf{x}) = \int_{\theta, \kappa} f^1 \boldsymbol{\tau}(\theta) d\kappa d\theta. \tag{4.11}$$

To compute J^0 we insert expression (4.9) into (4.11) and find

$$J^0(t, \mathbf{x}) = \int_{\theta, \kappa} \left(-\boldsymbol{\chi} \cdot \nabla_x n^0 + n^1 \frac{M}{2\pi} \right) \boldsymbol{\tau}(\theta) d\kappa d\theta. \tag{4.12}$$

The second term vanishes and the first one can be written

$$J^0(t, \mathbf{x}) = - \left(\int_{\theta, \kappa} \boldsymbol{\tau} \otimes \boldsymbol{\chi} d\theta d\kappa \right) \nabla_x n^0, \tag{4.13}$$

which is nothing but formula (2.18) with the diffusivity tensor D given by (4.28).

This shows the formal convergence of the solution of the Fokker-Planck equation (2.15) to that of the diffusion system (2.17), (2.18).

Now, to make this proof rigorous, we need to justify all the formal convergences. In the framework of the Hilbert expansion, this requires to work out the regularity of the various terms of the expansion. This is doable and actually leads to stronger convergences than the one we are going to prove, but this is a bit technical (see e.g. [18]).

What we are going to do instead is proving a convergence result in a weaker topology without using the Hilbert expansion technique. The method is close to the so-called moment method, which consists in integrating the equation against suitable test functions. This convergence proof is developed in Sect. 4.3, but before that, we state an existence result for the original Fokker-Planck equation (4.1).

4.2 Functional Setting and Existence Result

We define the differential operator D acting on smooth functions $f(\kappa)$ by:

$$Df = \partial_\kappa(\kappa f) + \alpha^2 \partial_{\kappa^2} f. \tag{4.14}$$

We state some properties of D , the proofs of which are easy and left to the reader. We recall that $M(\kappa)$ denotes the normalized Gaussian with zero mean and variance α^2 (4.4).

Proposition 4.1 *Let f and g be smooth functions decreasing at infinity. The following identities hold true:*

$$Df = \alpha^2 \frac{\partial}{\partial \kappa} \left(M \frac{\partial f}{\partial \kappa} M \right), \tag{4.15}$$

$$\int_{\mathbb{R}} Dfg \frac{d\kappa}{M} = -\alpha^2 \int_{\mathbb{R}} M \partial_{\kappa} \left(\frac{f}{M} \right) \partial_{\kappa} \left(\frac{g}{M} \right) d\kappa = \int_{\mathbb{R}} f Dg \frac{d\kappa}{M}, \tag{4.16}$$

$$\int_{\mathbb{R}} Dff \frac{d\kappa}{M} = -\alpha^2 \int_{\mathbb{R}} M \left| \partial_{\kappa} \left(\frac{f}{M} \right) \right|^2 d\kappa \leq 0, \tag{4.17}$$

$$Df = 0 \Leftrightarrow \exists c \in \mathbb{R}, \quad f = cM. \tag{4.18}$$

The first identity translates the fact that M is the stationary measure of the Ornstein-Uhlenbeck process. The second one that D is formally self-adjoint with respect to the measure $d\kappa/M$. The third one shows that D is dissipative. The same inequality holds with any non-decreasing function $\eta(f)$, indeed, $\int D(\eta(f))\eta(f)M^{-1}d\kappa \leq 0$. If η is the logarithm function, the corresponding quantity would be the relative entropy dissipation of f with respect to M . Entropy plays an important role in kinetic theory (see [13] for a review). Finally, the last quantity states that the kernel of D is one-dimensional and spanned by M .

Proposition 4.1 shows that the natural L^2 norm associated with this operator has a weight M^{-1} and that the natural H^1 semi-norm is given by the right-hand side of (4.17). This motivates the introduction of the following functional spaces, endowed with their naturally associated Hilbert structures and norms:

$$H = \left\{ u : \Pi_{\theta} \times \mathbb{R}_{\kappa} \rightarrow \mathbb{R} / \int_{\theta, \kappa} |u(\theta, \kappa)|^2 \frac{d\theta d\kappa}{M} < +\infty \right\}, \tag{4.19}$$

$$V = \left\{ u \in H / \int_{\Pi, \mathbb{R}} M \left| \partial_{\kappa} \left(\frac{u}{M} \right) \right|^2 d\kappa d\theta < +\infty \right\}, \tag{4.20}$$

$$L_M^2 = L^2(\mathbb{R}_x^2, H), \quad X = L^2([0, T] \times \mathbb{R}_x^2, V). \tag{4.21}$$

Identifying H with its dual, we have a Hilbertian triple $V \subset H \subset V'$, where V' is the dual of V and all injections are continuous. They are not compact because V does not bring any regularity with respect to θ .

The existence proof follows closely the existence proof of [15] (see Appendix A of this reference) and for this reason, is omitted (see also [18]). The proof relies on an existence theorem due to J.L. Lions [37].

Proposition 4.2 *Let $\varepsilon > 0$. We assume that f_0 belongs to L_M^2 defined by (4.21). Then there exists a unique solution f^ε to (2.15) with initial datum f^0 in the class of functions Y defined by:*

$$Y = \{ f \in X / \partial_t f + \varepsilon^{-1} \tau \cdot \nabla_x f + \varepsilon^{-2} \kappa \partial_{\theta} f \in X' \}.$$

Moreover, we have the inequality for any $T > 0$:

$$\|f^\varepsilon(T)\|_{L_M^2}^2 + \frac{\alpha^2}{\varepsilon^2} \int_0^T \int_{x, \theta, \kappa} M \left| \partial_{\kappa} \left(\frac{f^\varepsilon}{M} \right) \right|^2 d\kappa d\theta dx dt = \|f^\varepsilon(0)\|_{L_M^2}^2. \tag{4.22}$$

Estimate (4.22) is obtained via a Green and a trace formula for functions belonging to Y which can be deduced from the one proved in [15].

4.3 Rigorous Asymptotics

We first study operator A given by (4.2), i.e. $Af = \kappa \partial_{\theta} f - Df$ and state some properties which will be proved in Appendix B. We view A as an unbounded operator on the Hilbert

space H with DOMAIN $D(A)$ given by:

$$D(A) = \{u(\theta, \kappa) \in V / Au \in H\}.$$

Lemma 4.1 *Operator A is maximal monotone. Moreover its kernel (or Null-space) is given by:*

$$\text{Ker}(A) = \{cM, c \in \mathbb{R}\}, \tag{4.23}$$

with M defined by (4.4).

Lemma 4.2 *The adjoint A^* of A in H is given by $A^*f = -\kappa\partial_\theta f - Df$. It is a Maximal monotone operator with domain $D(A^*) = D(A)$ and $\text{Ker}(A^*) = \text{Ker}(A)$.*

Proposition 4.3 *Let $g \in H$. Then, there exists $u \in D(A)$ such that*

$$Au = g, \tag{4.24}$$

if and only if g satisfies the following solvability condition:

$$\int_{\theta,\kappa} g(\theta, \kappa) d\theta d\kappa = 0. \tag{4.25}$$

Moreover, the solution u is unique up to a constant times M . A unique solution can be singled out by prescribing the condition

$$\int_{\theta,\kappa} u(\theta, \kappa) d\theta d\kappa = 0. \tag{4.26}$$

The same lemma applies to the equation $A^*u = g$.

As an application of this lemma, let χ be the solution of:

$$A\chi = \tau(\theta)\frac{M}{2\pi}, \tag{4.27}$$

with $\tau(\theta) = (\cos\theta, \sin\theta)$. Since τ has zero average over θ and κ , χ is well-defined and unique thanks to Proposition 4.3. Then, we define the tensor D by:

$$D = \int_{\theta,\kappa} \tau(\theta) \otimes \chi d\theta d\kappa. \tag{4.28}$$

Note that, since $\int_{\theta,\kappa} \tau(\theta)M(\kappa) d\theta d\kappa = 0$, it would not change the value of D to add any element of $\text{Ker}(A)$ to χ .

Lemma 4.3 *Let R denote the reflection operator $u(\theta, \kappa) \rightarrow Ru(\theta, \kappa) = u(\theta, -\kappa)$. Then, $\chi^* = R\chi$ is the unique solution (satisfying (4.26)) of*

$$A^*\chi^* = \tau(\theta)\frac{M}{2\pi}, \tag{4.29}$$

and we have

$$D = \int_{\theta, \kappa} \boldsymbol{\tau}(\theta) \otimes \boldsymbol{\chi}^* d\theta d\kappa. \tag{4.30}$$

Proof Obviously, D commutes with R : $DR = RD$ while $\kappa \partial_\theta$ anticommutes with R : $\kappa \partial_\theta (Ru) = -R(\kappa \partial_\theta u)$. Therefore, $RA = A^*R$. Since the right-hand side of (4.27) is invariant by R , applying R to both sides of (4.27) leads to (4.29). Then, the change of variables $\kappa' = -\kappa$ in the integral at the right-hand side of (4.30) shows that it is equal to D . \square

To study the limit $\varepsilon \rightarrow 0$, we make the following hypothesis on the initial conditions.

Hypothesis 4.1 We suppose that the initial condition f_0^ε is uniformly bounded in L^2_M and converges weakly in L^2_M to f^0 as $\varepsilon \rightarrow 0$.

We can now prove Theorem 2.2. The initial datum for the diffusion system (2.17), (2.18) will be shown to be:

$$n^0_{t=0} = n^0_0 = \int_{\theta, \kappa} f^0_0(\mathbf{x}, \theta, \kappa) d\kappa d\theta. \tag{4.31}$$

Proof of Theorem 2.2 By Hypothesis 4.1 inequality (4.22) implies:

$$\|f^\varepsilon(T)\|_{L^2_M}^2 + \frac{\alpha^2}{\varepsilon^2} \int_0^T \int_{\mathbf{x}, \theta, \kappa} M \left| \partial_\kappa \left(\frac{f^\varepsilon}{M} \right) \right|^2 d\kappa d\theta d\mathbf{x} dt \leq C, \tag{4.32}$$

with C independent of ε . So $(f^\varepsilon)_\varepsilon$ is a bounded sequence in $L^\infty(0, T, L^2_M)$ and satisfies

$$\int_0^T \int_{\mathbf{x}, \theta, \kappa} M \left| \partial_\kappa \left(\frac{f^\varepsilon}{M} \right) \right|^2 d\kappa d\theta d\mathbf{x} dt \leq C\varepsilon^2, \tag{4.33}$$

for any time interval T (by the diagonal process, we will eventually be able to take an increasing sequence of times T tending to infinity, so that the result will be valid on the whole interval $t \in (0, \infty)$). Therefore, there exists $f^0 \in L^\infty(0, T, L^2_M)$ and a subsequence, still denoted by f^ε , satisfying:

$$f^\varepsilon \xrightarrow{\varepsilon \rightarrow 0} f^0 \quad \text{in } L^\infty(0, T, L^2_M) \text{ weak star.}$$

Furthermore, with (4.33), we deduce that $f^0 = C(\mathbf{x}, \theta, t)M(\kappa)$. Then, letting $\varepsilon \rightarrow 0$ in (2.15), we get that $Af^0 = 0$ in the distributional sense. This implies that $C(\mathbf{x}, \theta, t)$ is independent of θ and we can write

$$f^0(t, \mathbf{x}, \theta, \kappa) = n^0(t, \mathbf{x}) \frac{M(\kappa)}{2\pi}, \tag{4.34}$$

the quantity $n^0(t, \mathbf{x}) = \int f^0(t, \mathbf{x}, \theta, \kappa) d\theta d\kappa$ being the density associated with f^0 .

Our next task is to show that n^0 satisfies the diffusion model (2.17), (2.18) with initial condition (4.31). We first note that f^ε is a weak solution of (2.15) with initial condition f^ε_0 in the following sense: f^ε satisfies:

$$\int_0^T \int_{\mathbf{x}, \theta, \kappa} f^\varepsilon \left(-\varepsilon \partial_t \varphi - \boldsymbol{\tau} \cdot \nabla_{\mathbf{x}} \varphi + \frac{1}{\varepsilon} A^*(\varphi) \right) \frac{d\kappa d\theta d\mathbf{x}}{M} dt = \varepsilon \int_{\mathbf{x}, \theta, \kappa} f^\varepsilon_0 \varphi_{t=0} \frac{d\kappa d\theta d\mathbf{x}}{M}, \tag{4.35}$$

for all test functions φ in the space $C_c^2([0, T] \times \mathbb{R}_x^2 \times \Pi_\theta \times \mathbb{R}_\kappa)$ of twice continuously differentiable functions with compact support in $[0, T] \times \mathbb{R}_x^2 \times \Pi_\theta \times \mathbb{R}_\kappa$. Again, the trace at $t = 0$ has a meaning, thanks to a trace formula for functions in Y which is proven in [15].

We recall the definition of the flux (4.5). We prove that J^ε has a weak limit as $\varepsilon \rightarrow 0$. To this aim, in the weak formulation (4.35), we take as a test function $\varphi = \phi(t, \mathbf{x}) \cdot \chi^*(\theta, \kappa)$ with χ^* the auxiliary function defined as the solution to (4.29) and ϕ is a smooth compactly supported vector test function of (\mathbf{x}, t) . Although φ does not have a compact support, a standard truncation argument (which is omitted here) can be used to bypass this restriction. This allows us to write:

$$\begin{aligned} & \int_0^T \int_{\mathbf{x}, \theta, \kappa} \left[f^\varepsilon(-\varepsilon \partial_t - (\boldsymbol{\tau} \cdot \nabla_{\mathbf{x}}))(\phi \cdot \chi^*) + \frac{1}{\varepsilon} f^\varepsilon \boldsymbol{\tau} \frac{M}{2\pi} \cdot \phi \right] \frac{d\kappa d\theta d\mathbf{x}}{M} dt \\ &= \varepsilon \int_{\mathbf{x}, \theta, \kappa} f_0^\varepsilon \phi_{t=0} \cdot \chi \frac{d\kappa d\theta d\mathbf{x}}{M}. \end{aligned}$$

Taking the limit $\varepsilon \rightarrow 0$, we find:

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_0^T \int_{\mathbf{x}} J^\varepsilon \cdot \phi d\mathbf{x} dt &= 2\pi \int_0^T \int_{\mathbf{x}, \theta, \kappa} f^0(\boldsymbol{\tau} \cdot \nabla_{\mathbf{x}})(\phi \cdot \chi^*) \frac{d\kappa d\theta d\mathbf{x}}{M} dt \\ &= \int_0^T \int_{\mathbf{x}} n^0 \nabla_{\mathbf{x}} \cdot \left(\left(\int_{\theta, \kappa} \chi^* \otimes \boldsymbol{\tau} d\kappa d\theta \right)^T \phi \right) d\mathbf{x} dt, \end{aligned}$$

where the exponent T denotes the transpose of a matrix. Using (4.30) and taking the limit $\varepsilon \rightarrow 0$ shows that J^ε converges weakly (in the distributional sense) towards J^0 satisfying

$$\int_0^T \int_{\mathbf{x}} J^0 \cdot \phi d\mathbf{x} dt = \int_0^T \int_{\mathbf{x}} n^0 \nabla_{\mathbf{x}} \cdot ((D)^T \phi) d\mathbf{x} dt.$$

This last equation is the weak form of (2.18).

Finally, to prove (2.17), we apply the weak formulation (4.35) to a test function of the form $\varphi = \phi(t, \mathbf{x})M(\kappa)$, where again, $\phi(x, t)$ is a scalar, smooth and compactly supported test function of (\mathbf{x}, t) in $\mathbb{R}^2 \times [0, T]$. This gives:

$$- \int_0^T \int_{\mathbf{x}, \theta, \kappa} f^\varepsilon((\varepsilon \partial_t + (\boldsymbol{\tau} \cdot \nabla_{\mathbf{x}}))\varphi) d\kappa d\theta d\mathbf{x} dt = \varepsilon \int_{\mathbf{x}, \theta, \kappa} f_0^\varepsilon \phi_{t=0} d\kappa d\theta d\mathbf{x}.$$

Dividing by ε and taking the limit $\varepsilon \rightarrow 0$, we get:

$$- \int_0^T \int_{\mathbf{x}} (n^0 \partial_t \varphi + J^0 \cdot \nabla_{\mathbf{x}} \phi) d\mathbf{x} dt = \int_{\mathbf{x}} n_0^0 \phi_{t=0} d\mathbf{x},$$

where n_0^0 is defined by (4.31). This last equation is exactly the weak formulation of (2.17), with initial datum n_0^0 . This concludes the proof. \square

5 Equivalence of the Two Methods

In this section, we show that both methods lead to the same value of the diffusion coefficient (Theorem 2.3).

The first step is to show that we can approximate the solution of (4.24) by the solution of the associated evolution equation. More precisely, in Appendix C, we prove the following lemma:

Lemma 5.1 *Let g in H satisfying (4.25) and u_∞ in $D(A)$ be the solution of (4.24) satisfying (4.26). Let $u_0 \in D(A)$ satisfying (4.26). Then, the solution $u(t)$ of the evolution problem:*

$$\partial_t u = -Au + g, \quad u_{t=0} = u_0, \tag{5.1}$$

weakly converges to u_∞ in H as t tends to ∞ .

With this lemma we can explicitly calculate the tensor D and prove the Theorem 2.3:

Proof of Theorem 2.3 Let $\chi(t)$ be the solution of

$$\partial_t \chi = -A\chi + \tau(\theta) \frac{M(\kappa)}{2\pi}, \quad \chi(t=0) = 0. \tag{5.2}$$

Thanks to Lemma 5.1, $\chi(t)$ weakly converges to χ in H when $t \rightarrow \infty$. It follows that:

$$\int_{\kappa, \theta} \chi(t) \otimes \tau \, d\kappa d\theta \xrightarrow{t \rightarrow +\infty} \int_{\kappa, \theta} \chi \otimes \tau \, d\kappa d\theta. \tag{5.3}$$

Let us consider the first component of $\chi(t)$, which we denote by $u(t)$ and the integrals $\int_{\kappa, \theta} u(t) \cos \theta \, d\kappa d\theta$ and $\int_{\kappa, \theta} u(t) \sin \theta \, d\kappa d\theta$. Because u satisfies (5.2), it admits the following representation (see the proof of Lemma 5.1):

$$u(t) = \int_0^t T_s \left(\cos \theta \frac{M(\kappa)}{2\pi} \right) ds,$$

where T_t is the semi-group generated by $-A$ (see [44]). With this expression, we evaluate the integral of $u(t)$ against $\cos \theta$:

$$\begin{aligned} \int_{\kappa, \theta} u(t) \cos \theta \, d\kappa d\theta &= \int_{\kappa, \theta} \int_0^t T_s \left(\cos \theta \frac{M(\kappa)}{2\pi} \right) ds \cos \theta \, d\kappa d\theta \\ &= \int_0^t \int_{\kappa, \theta} \cos \theta \frac{M(\kappa)}{2\pi} T_s^*(\cos \theta) \, d\kappa d\theta ds, \end{aligned}$$

where T^* is the adjoint operator of T in $L^2(\theta, \kappa)$ generated by $-A^*$, where

$$A^*(f) = -\kappa \partial_\theta f + \kappa \partial_\kappa f - \alpha^2 \partial_{\kappa^2} f.$$

Note that we are referring here to the adjoint in the standard L^2 sense and not in the weighted space H . This is why A^* does not coincide with A^* defined in Lemma 4.2. The semi-group T_t^* admits a probabilistic representation: for all regular functions $f(\theta, \kappa)$

$$T_t^*(f)(\theta, \kappa) = \mathbb{E}\{f(\theta(t), \kappa(t)) | \theta_0 = \theta, \kappa_0 = \kappa\},$$

where $(\kappa(t), \theta(t))$ is the solution of the stochastic differential equation (2.5), (2.6). Using this representation, we have:

$$\begin{aligned} \int_{\kappa} M(\kappa) T_s^*(\cos \theta) d\kappa &= \int_{\kappa} M(\kappa) \mathbb{E}\{\cos \theta_s | \theta_0 = \theta, \kappa_0 = \kappa\} d\kappa \\ &= \mathbb{E}\{\cos \theta_s | \theta_0 = \theta, \kappa_0 = Z\}, \end{aligned}$$

where Z is a random variable independent of B_t with density M . Using Lemma 3.1, we have:

$$\mathbb{E}\{\cos \theta_s | \theta_0 = \theta, \kappa_0 = Z\} = \mathbb{E}\{\cos(\theta + Y_s)\},$$

with Y_s a Gaussian random variable with zero mean and variance β_s^2 given by (3.10). Then:

$$\mathbb{E}\{\cos(\theta + Y_s)\} = \mathbb{E}\{\cos \theta \cos Y_s - \sin \theta \sin Y_s\} \cos \theta \mathbb{E}\{\cos Y_s\},$$

because the density of Y_s is even and implies that $\mathbb{E}\{\sin Y_s\} = 0$. Finally using (3.12), we have:

$$\int_{\kappa} M(\kappa) T_s^*(\cos \theta) d\kappa = \cos \theta e^{-\frac{\beta_s^2}{2}}.$$

Then, the first integral is given by:

$$\int_{\kappa, \theta} u(t) \cos \theta d\kappa d\theta = \int_0^t \int_{\theta} \frac{\cos \theta}{2\pi} \cos \theta e^{-\frac{\beta_s^2}{2}} d\theta ds \int_0^t \frac{1}{2} e^{-\frac{\beta_s^2}{2}} ds.$$

We can proceed similarly to evaluate the integral of $u(t)$ against $\sin(t)$. This gives:

$$\int_{\kappa, \theta} u(t) \sin \theta d\kappa d\theta = \int_0^t \int_{\theta} \frac{\cos \theta \sin \theta}{2\pi} \mathbb{E}\{\cos Y_s\} d\theta ds = 0.$$

It remains to evaluate the integrals involving the second component of vector $\chi(t)$ which we denote by $v(t)$. By the same method as for $u(t)$, we get:

$$\int_{\kappa, \theta} v(t) \cos \theta d\kappa d\theta = 0 \quad \text{and} \quad \int_{\kappa, \theta} v(t) \sin \theta d\kappa d\theta = \int_0^t \frac{1}{2} e^{-\frac{\beta_s^2}{2}} dt.$$

Collecting these formulae, we can write:

$$\int_{\kappa, \theta} \chi(t) \otimes \tau d\kappa d\theta = \frac{\mathcal{D}(t)}{2} Id,$$

with $\mathcal{D}(t) = \int_0^t e^{-\alpha^2(-1+u+e^{-u})} du$. Taking the limit $t \rightarrow +\infty$ and using (5.3), shows that (2.19) holds true and completes the proof of the theorem. □

6 Numerical Simulation

We simulate individual trajectories satisfying (2.4)–(2.6) with initial conditions given by (3.1)–(3.4). If we fix a time step Δt , using (3.7), (3.8), we have:

$$\begin{cases} \kappa_{(n+1)\Delta t} = \gamma \kappa_{n\Delta t} + G_{(n+1)}, \\ \theta_{(n+1)\Delta t} = \theta_0 + \kappa_0 - \kappa_{(n+1)\Delta t} + \sqrt{2\alpha} B_{(n+1)\Delta t} \end{cases}$$

with $\gamma = e^{-\Delta t}$ and $G_{(n+1)}$ a Gaussian random variable with zero mean and variance $2\alpha^2(1 - e^{-2\Delta t})$ independent of $\kappa_{n\Delta t}$. With this formula, we can simulate recursively the process $(\kappa_{n\Delta t}, \theta_{n\Delta t})_n$ exactly (in the sense that it has the same law as the exact solution). To generate the Brownian motion, we just compute the increments $B_{(n+1)\Delta t} - B_{n\Delta t}$ since they are Gaussian and independent of $B_{n\Delta t}$. On the other hand, these increments are not independent of $G_{(n+1)}$. Fortunately, we can compute the covariance matrix of the Gaussian vector $(G_{(n+1)}, B_{(n+1)\Delta t} - B_{n\Delta t})$:

$$\begin{pmatrix} G_{(n+1)} \\ B_{(n+1)\Delta t} - B_{n\Delta t} \end{pmatrix} \sim \mathcal{N}(0, C),$$

where $\mathcal{N}(0, C)$ is a two-dimensional Gaussian vector with zero mean and covariance matrix C given by:

$$C = \begin{bmatrix} 2\alpha^2(1 - e^{-2\Delta t}) & \sqrt{2}\alpha(1 - e^{-\Delta t}) \\ \sqrt{2}\alpha(1 - e^{-\Delta t}) & \Delta t \end{bmatrix}.$$

Knowing this covariance matrix, we can simulate the Gaussian vector $(G_{(n+1)}, B_{(n+1)\Delta t} - B_{n\Delta t})$ using the Cholesky method: we generate (X_1, X_2) a vector of two independent normal law, and take $\sqrt{C}(X_1, X_2)^T$ as realization of the Gaussian vector.

Now for the position \mathbf{x} , since we do not have any explicit expression, we use a discrete approximation scheme of order $O((\Delta t)^2)$. For example, the first component x_1 of \mathbf{x} is approximated by:

$$\begin{aligned} x_1((n + 1)\Delta t) &= x_1(n\Delta t) + \int_{n\Delta t}^{(n+1)\Delta t} \cos \theta(s) ds \\ &\approx x_1(n\Delta t) + \frac{\Delta t}{2} (\cos \theta(n\Delta t) + \cos \theta((n + 1)\Delta t)). \end{aligned}$$

We present four trajectories obtained with different values of the parameter α in Fig. 3. As the parameter α increases, the excursions towards large positive or negative curvatures become larger. As a consequence, the spinning of the trajectory around itself increases and, from almost a straight line when $\alpha = 0.1$, the trajectory shrinks and looks closer and closer to a wool ball. In this way, we can visualize the decay of \mathcal{D} with respect to α .

To illustrate Theorem 2.1, we use a Monte-Carlo method to simulate the variance of the process \mathbf{x} . We simulate N independent trajectories and we compute the variance of the sample at each time step. In Fig. 4, we compare the result obtained with $N = 2000$ and the theoretical prediction given by the (3.6). The figure shows an excellent agreement between the computation and the theoretical prediction. Additionally, after an initial transient, the growth of the variance is linear, in accordance with the theoretical result (2.9). We can use the slope of the asymptotically linear part of the curve to give a numerical estimate of the diffusion coefficient \mathcal{D} . For this purpose, we fit a straight line (in the mean-square sense) between times $T/2$ and T . We remove the data between 0 and $T/2$ because the initial transient is not linear and including them would deteriorate the accuracy of the measurement. We compare the slope of the fitted line with the theoretical value (2.9). We report the result of this comparison for two values of α ($\alpha = 0.1$ and $\alpha = 2.$, with $T = 1200$ time units in Table 1). The approximation is quite good, with an error comprised between 2 and 3%, which can be attributed to numerical noise and to an insufficient approximation of the asymptotic state. To illustrate the influence of the initial transient, we take $T = 120$ time units in the




Fig. 3 Four trajectories simulated with different value of α : $\alpha = 0.1$ (top left), $\alpha = 0.5$ (top right), $\alpha = 1$. (bottom left) and $\alpha = 2$. (bottom right). The simulation is run during 120 time units with a time step $dt = 0.05$ time unit




Fig. 4 Variance of the process $x(t)$: comparison between the numerical simulation (points) and the theoretical prediction (solid line) for two values of α : $\alpha = 0.1$ (left) and $\alpha = 2$ (right)

case $\alpha = 0.1$ and report the result in Table 2. There, the approximation is quite poor, because the asymptotic state has not yet been reached.

Table 1 Diffusion coefficient: comparison of the numerical estimate obtained by fitting the numerical values with a straight line over the time interval $[T/2, T]$ with the theoretical prediction (2.9). $T = 1200$ units of time

	D simulation	D theoretical	Relative error
$\alpha = 0.1$	98.5	101	2.5%
$\alpha = 2$	0.708	0.725	2.4%

Table 2 Same as Table 1 but with $T = 120$ units of time. The agreement is poor because the asymptotic state is not reached

	D simulation	D theoretical	Relative error
$\alpha = 0.1$	58.8	101	72 %





Fig. 5 The spatial distribution n for $\alpha = 2$ at time $T = 30$ time units

In order to illustrate Theorem 2.2, we plot the spatial density $n(t, \mathbf{x})$ of the distribution $f(t, \mathbf{x}, \theta, \kappa)$ using a Monte-Carlo algorithm for $\alpha = 2$ and $T = 30$ time units on Fig. 5. We see that the density has the Gaussian shape of the solution of a diffusion equation, in accordance with the prediction of the theorem. To make a more quantitative comparison, we compare it with the asymptotic prediction, i.e. the solution of the diffusion equation (2.17), (2.18), by computing the difference in L^1 norm. The results are reported in Fig. 6. We plot the L^1 norm of the difference for $\alpha = 1$. and for four values of the parameter ε : $1, \frac{1}{2}, \frac{1}{5}$ and $\frac{1}{10}$. As expected, the agreement is better as ε is smaller. However, at large times, all

Fig. 6 L^1 norm of the difference between $n^\varepsilon(t, \cdot) = \int f^\varepsilon(t, \cdot, \theta, \kappa) d\theta d\kappa$ and its asymptotic limit $n^0(t)$ as a function of time t , with $\alpha = 1$ and $N = 10^4$ simulation particles



solutions are eventually close to the solution of the diffusion equation. Roughly speaking, the time at which the solution of the diffusion equation starts to be a good approximation of the solution of the kinetic equation scales like ε . This means that, after an initial transient the duration of which may depend on ε , the solution is close to that of the diffusion equation, no matter the value of ε .

7 Conclusion

In this paper, the large-scale dynamics of the ‘Persistent Turning Walker’ (PTW) model of fish behavior has been analyzed. It has been shown, by two different methods, that the large scale limit of this model is of diffusion type, and an explicit formula for the diffusion coefficient has been provided. While the direct analysis of the stochastic trajectories provides a direct route to the value of the diffusion constant, the diffusion approximation of the associated forward Kolmogorov equation, which is of Fokker-Planck type, gives a more systematic way to extend the theory to more complex nonlinear cases. Such a nonlinear situation will be encountered when, in the near future, the nonlinear interactions between the individuals will be introduced within the PTW model (when the experimental data concerning the interactions among groups of fish will be analyzed). We expect that, in this context, the diffusion approximation methodology will have to be exploited thoroughly to allow access to the large scale behaviour of the system.

Acknowledgements The authors wish to thank Guy Théraulaz and Jacques Gautrais of the ‘Centre de Recherches sur la Cognition Animale’ in Toulouse, for introducing them to the model and for stimulating discussions.

Appendix A: Proofs of Sect. 3

Proof of Lemma 3.1 Formula (3.8) is standard in the theory of Ornstein-Uhlenbeck processes [40]. To obtain (3.7), we integrate $\kappa(t)$ with respect to time:

$$\int_0^t \kappa(s) ds = (1 - e^{-t})\kappa_0 + \sqrt{2}\alpha \int_0^t \int_0^s e^{-s} e^u dB_u ds.$$

Interchanging the order of integrations and integrating with respect to s , we deduce:

$$\int_0^t \kappa(s) ds = (1 - e^{-t})\kappa_0 + \sqrt{2}\alpha \int_0^t (1 - e^{-(t-u)}) dB_u.$$

Then we develop the integral:

$$\int_0^t \kappa(s) ds = (1 - e^{-t})\kappa_0 + \sqrt{2}\alpha B_t - (\kappa(t) - e^{-t}\kappa_0).$$

This formula can be rewritten:

$$\int_0^t \kappa(s) ds = \kappa_0 - \kappa(t) + \sqrt{2}\alpha B_t,$$

which easily leads to (3.7).

We now calculate the mean and the variance of $K_0^t = \int_0^t \kappa(s) ds$. Since $\kappa(s)$ is of zero mean, its integral K_0^t is also of zero mean: $\mathbb{E}\{K_0^t\} = 0$. Now for the variance of K_0^t , we can write:

$$\text{Var}\{K_0^t\} = \mathbb{E} \left\{ \left((1 - e^{-t})\kappa_0 - \sqrt{2}\alpha e^{-t} \int_0^t e^s dB_s + \sqrt{2}\alpha B_t \right)^2 \right\}.$$

Using that κ_0 and B_s are independent, we can develop the square and get:

$$\text{Var}\{K_0^t\} = (1 - e^{-t})^2 \mathbb{E}\{\kappa_0^2\} + 2\alpha^2 \mathbb{E} \left\{ \left(-e^{-t} \int_0^t e^s dB_s + B_t \right)^2 \right\}.$$

Let us consider the second term. By Ito's formula, we have

$$\begin{aligned} \mathbb{E} \left\{ \left(-e^{-t} \int_0^t e^s dB_s + B_t \right)^2 \right\} &= e^{-2t} \mathbb{E} \left\{ \left(\int_0^t e^s dB_s \right)^2 \right\} - 2e^{-t} \mathbb{E} \left\{ B_t \int_0^t e^s dB_s \right\} + \mathbb{E} \{ B_t^2 \} \\ &= e^{-2t} \int_0^t e^{2s} ds - 2e^{-t} \mathbb{E} \left\{ B_t \left(e^t B_t - \int_0^t e^s B_s ds \right) \right\} + t, \end{aligned}$$

where the Ito correction term is zero due to the fact that $\exp s$ is a deterministic process. We can simplify this expression again since $\mathbb{E}\{B_t B_s\} = \min(t, s)$ and get:

$$\begin{aligned} \mathbb{E} \left\{ \left(-e^{-t} \int_0^t e^s dB_s + B_t \right)^2 \right\} &= \frac{1 - e^{-2t}}{2} - 2e^{-t} \left(e^t t - \int_0^t e^s s ds \right) + t \\ &= \frac{1 - e^{-2t}}{2} - 2(1 - e^{-t}) + t. \end{aligned}$$

Using also that $\mathbb{E}\{\kappa_0^2\} = \alpha^2$, the variance of K_0^t is written:

$$\text{Var}\{K_0^t\} = (1 - e^{-t})^2 \alpha^2 + 2\alpha^2 \left(\frac{1 - e^{-2t}}{2} - 2(1 - e^{-t}) + t \right).$$

Developing and simplifying the expression, we find (3.10), which ends the proof. □

Appendix B: Proofs of Sect. 4

Proof of Lemma 4.1 Let $u \in D(A)$. Then, $\kappa \partial_\theta u \in V'$ and Lemma A1 of [15] shows that the Green formula for functions $u \in V$ such that $\kappa \partial_\theta u \in V'$ is legitimate. Therefore, taking the inner product of $A(u)$ against u , we find:

$$\langle A(u), u \rangle_H = \int_{\theta, \kappa} \alpha^2 M \left| \partial_\kappa \left(\frac{u}{M} \right) \right|^2 d\theta d\kappa \geq 0. \tag{B.1}$$

So, A is a monotone operator in H . To show that A is maximal monotone, we prove that for any $g \in H$, there exists $u \in D(A)$ such that:

$$u + Au = g. \tag{B.2}$$

Taking the inner product of (B.2) against a test function φ in the space $\mathcal{D}(\Pi_\theta \times \mathbb{R}_\kappa)$ of infinitely differentiable and compactly supported functions on $\Pi_\theta \times \mathbb{R}_\kappa$ leads to the variational problem:

$$\int_{\kappa, \theta} \left[u(\varphi - \kappa \partial_\theta \varphi) \frac{1}{M} + M \partial_\kappa \left(\frac{u}{M} \right) \partial_\kappa \left(\frac{\varphi}{M} \right) \right] d\theta d\kappa = \int_{\theta, \kappa} g \varphi \frac{d\kappa d\theta}{M}. \tag{B.3}$$

Again, the same theory as in the Appendix A of [15] (based the result by J.L. Lions in [37]) applies to prove the existence of a solution to (B.3) with u in V such that $\kappa \partial_\theta u \in V'$. From there, it immediately follows that $u \in D(A)$.

It is immediate to see that any function of the form $u(\theta, \kappa) = CM(\kappa)$ for any constant C belongs to the kernel of A . Conversely, suppose that $u \in \text{Ker} A$. Then, by (B.1), there exists a function $C(\theta) \in L^2(\Pi)$ such that $u(\theta, \kappa) = C(\theta)M(\kappa)$. But again, $A(u) = 0$ implies that $\kappa \partial_\theta C(\theta)M = 0$. So $C(\theta)$ is a constant, which proves (4.23). \square

Proof of Proposition 4.3 The ‘only if’ part of the theorem is obvious since, using Green’s formula (again, obtained by adapting that of Appendix B of [15], we have $\int Aud\theta d\kappa = 0$).

To prove the ‘if’ part, we borrow a method from (for instance) [12]. To find a solution to (4.24), we look at a perturbed equation:

$$\lambda u + Au = g, \tag{B.4}$$

with $\lambda > 0$. Since A is maximal monotone in H (Lemma 4.1), (B.4) admits a solution u_λ for all positive λ [9]. To prove the existence of a solution to (4.24), we want to extract a subsequence, still denoted by (u_λ) which converges weakly in H . For this purpose, it is enough to show that there exists a bounded subsequence.

We proceed by contradiction, supposing that the (full) sequence $N_\lambda = \|u_\lambda\|_H \xrightarrow{\lambda \rightarrow 0} +\infty$. We define $U_\lambda = \frac{u_\lambda}{N_\lambda}$. U_λ satisfies $\|U_\lambda\|_H = 1$ for all λ and

$$\lambda U_\lambda + AU_\lambda = \frac{g}{N_\lambda}. \tag{B.5}$$

Since $(U_\lambda)_\lambda$ is a bounded sequence in H , we can extract a subsequence (still denoted by U_λ) such that $U_\lambda \rightharpoonup U$ in H weak as $\lambda \rightarrow 0$. Taking the limit $\lambda \rightarrow 0$ in (B.5), gives $A(U) = 0$. If we take the inner product of (B.5) with U_λ and then pass to the limit $\lambda \rightarrow 0$, we also find

that U belongs to V . So Lemma 4.1 applies and gives $U = cM$ with a constant $c \in \mathbb{R}$. Using (4.25), we also have:

$$\langle \lambda U_\lambda + AU_\lambda, M \rangle_H = \left\langle \frac{g}{N_\lambda}, M \right\rangle_H = 0.$$

So $\int_{\kappa, \theta} U_\lambda d\theta d\kappa = 0$ for all λ . Taking the limit $\lambda \rightarrow 0$ leads to $\int_{\kappa, \theta} U d\theta d\kappa = \int_{\kappa, \theta} CM(\kappa) d\theta d\kappa = 2\pi C = 0$, which implies $U = 0$. This proves:

$$U_\lambda \rightharpoonup 0 \quad \text{in } H \text{ weak.} \tag{B.6}$$

To get a contradiction, we now prove that the convergence is strong.

To this aim, we introduce a decomposition of the space H into two orthogonal subspaces. Let L be the closed subspace of H defined by:

$$L = \{c(\theta)M/c(\theta) \in L^2(\Pi_\theta)\},$$

with M defined by (4.4). So $H = L \oplus L^\perp$. We also define the orthogonal projector P of H onto L such that $Pf = (\int_\kappa f(\kappa, \theta) d\kappa)M$. Using this projection, we decompose the sequence $(U_\lambda)_\lambda$ as follows:

$$U_\lambda = c_\lambda(\theta)M + v_\lambda, \tag{B.7}$$

with $v_\lambda \in L^\perp$, i.e. $\int_\kappa v_\lambda d\kappa = 0$. To demonstrate that $U_\lambda \xrightarrow{\lambda \rightarrow 0} 0$ in H strongly, we first demonstrate that $v_\lambda \xrightarrow{\lambda \rightarrow 0} 0$ in H strongly.

Taking the inner product of the equation satisfied by U_λ (B.5) with U_λ gives:

$$\lambda \|U_\lambda\|_H^2 + \int_{\theta, \kappa} M \left| \partial_\kappa \frac{U_\lambda}{M} \right|^2 d\theta d\kappa = \frac{1}{N_\lambda} \langle g, U_\lambda \rangle_H.$$

Since $\partial_\kappa \frac{U_\lambda}{M} = \partial_\kappa \frac{v_\lambda}{M}$ and $\|U_\lambda\|_H = 1$, we get by taking the limit $\lambda \rightarrow 0$:

$$\int_{\theta, \kappa} M \left| \partial_\kappa \frac{v_\lambda}{M} \right|^2 d\theta d\kappa \xrightarrow{\lambda \rightarrow 0} 0. \tag{B.8}$$

Now Gross inequality [32] gives, for any $v \in V$:

$$\alpha^2 \int_{\mathbb{R}} \left| \partial_\kappa \left(\frac{f}{M} \right) \right|^2 M d\kappa \geq \int_{\mathbb{R}} |f|^2 \frac{d\kappa}{M} - \left(\int_{\mathbb{R}} f d\kappa \right)^2. \tag{B.9}$$

Then, since $\int_\kappa v_\lambda d\kappa = 0$, we deduce:

$$\alpha^2 \int_{\mathbb{R}} \left| \partial_\kappa \frac{v_\lambda}{M} \right|^2 M d\kappa \geq \int_{\mathbb{R}} \frac{|v_\lambda|^2}{M} d\kappa.$$

Integrating this inequality with respect to θ and using (B.8), we find:

$$\|v_\lambda\|_H \xrightarrow{\lambda \rightarrow 0} 0, \quad \text{in } H \text{ strong.} \tag{B.10}$$

To prove the convergence of c_λ , we define the bounded operator $T: H \rightarrow L^2(\Pi_\theta)$ such that $Tf = \int_\kappa \kappa f d\kappa$. Having T acting on (B.5) and taking the limit $\lambda \rightarrow 0$, leads to:

$$TAU_\lambda \xrightarrow{\lambda \rightarrow 0} 0 \quad \text{in } L^2(\Pi_\theta) \text{ strong.} \tag{B.11}$$

If we develop the left-hand side, we find:

$$\begin{aligned} TAU_\lambda &= \int_\kappa \kappa^2 \partial_\theta U_\lambda d\kappa - \int_\kappa [\kappa \partial_\kappa (\kappa U_\lambda) - \alpha^2 \kappa \partial_{\kappa^2} U_\lambda] d\kappa \\ &= \int_\kappa \kappa^2 \partial_\theta U_\lambda d\kappa + \int_\kappa \kappa U_\lambda d\kappa. \end{aligned}$$

But using the decomposition $U_\lambda = c_\lambda M + v_\lambda$ (B.7), we have:

$$\left\| \int_\kappa \kappa U_\lambda d\kappa \right\|_{L^2(\theta)} = \left\| \int_\kappa \kappa v_\lambda d\kappa \right\|_{L^2(\theta)} \xrightarrow{\lambda \rightarrow 0} 0.$$

So, (B.11) leads to:

$$\int_\kappa \kappa^2 \partial_\theta U_\lambda d\kappa \xrightarrow{\lambda \rightarrow 0} 0 \quad \text{in } L^2(\Pi_\theta) \text{ strong.} \tag{B.12}$$

If we define $h_\lambda(\theta) = \int_\kappa \kappa^2 U_\lambda d\kappa$, (B.12) is equivalent to saying that $\|\partial_\theta h_\lambda\|_{L^2(\theta)} \xrightarrow{\lambda \rightarrow 0} 0$. Using the Poincare-Wirtinger inequality [9], there exists a constant C_0 such that:

$$\|h_\lambda - \bar{h}_\lambda\|_{L^2(\theta)} \leq C_0 \|\partial_\theta h_\lambda\|_{L^2(\theta)}, \tag{B.13}$$

with $\bar{h}_\lambda = \frac{1}{2\pi} \int_0^{2\pi} h_\lambda(\theta) d\theta$. Then, we develop \bar{h}_λ . We get:

$$\bar{h}_\lambda = \frac{1}{2\pi} \int_0^{2\pi} \int_\kappa \kappa^2 U_\lambda d\kappa d\theta = \langle U_\lambda, M \kappa^2 \rangle_H \xrightarrow{\lambda \rightarrow 0} 0 \quad \text{in } \mathbb{R},$$

since U_λ converges weakly to zero (see (B.6)). So, (B.13) leads to $h_\lambda \xrightarrow{\lambda \rightarrow 0} 0$ in $L^2(\Pi_\theta)$ strong. If we develop h_λ we find:

$$h_\lambda(\theta) = \int_\kappa \kappa^2 (c_\lambda(\theta)M + v_\lambda) d\kappa = \alpha^2 c_\lambda(\theta) + \int_\kappa \kappa^2 v_\lambda d\kappa.$$

Now, $\int_\kappa \kappa^2 v_\lambda d\kappa$ converges to zero in $L^2(\theta)$ strong because of (B.10) and we finally have:

$$c_\lambda(\theta) \xrightarrow{\lambda \rightarrow 0} 0 \quad \text{in } L^2(\theta) \text{ strong.}$$

Using the convergence of c_λ and v_λ , we can now prove the strong convergence of U_λ to 0 in H :

$$\|U_\lambda\|_H^2 = \|c_\lambda M\|_H^2 + \|v_\lambda\|_H^2 = \|c_\lambda\|_{L^2(\theta)}^2 + \|v_\lambda\|_H^2 \xrightarrow{\lambda \rightarrow 0} 0,$$

which contradicts the fact that U_λ has unit norm in H . This shows that there exists a bounded subsequence in the sequence u_λ . In fact, since the same proof can be applied to any subsequence, this shows that the whole sequence u_λ is bounded, but this is useless for our purpose.

We conclude the proof of Proposition 4.3 as follows: there exists a subsequence u_λ and a function u in H such that $u_\lambda \rightharpoonup u$ in H weak. Taking the limit of (B.4) as $\lambda \rightarrow 0$, we deduce that $Au = g$ in the sense of distributions. However, since $g \in H$, eq. $Au = g$ also holds in H . Moreover if we take the inner product of (B.4) with u_λ and pass to the limit $\lambda \rightarrow 0$, we find that u belongs to V . So u belongs to $D(A)$, which ends the proof of the ‘if’ part of the statement.

Finally, to prove uniqueness, we just remark that, two solutions of (4.24) differ from an element of the kernel of A and we apply (4.23). This ends the proof. \square

Appendix C: Proofs of Sect. 5

Proof of Lemma 5.1 The proof borrows some ideas from [24], but is simpler, due to the linear character of the problem. The difficulty is getting some compactness in time. Here, instead of considering time translates of the solution as in [24], we will consider time integrals over a fixed interval length Δt .

Since operator A is maximal monotone on H (see Lemma 4.1), operator $-A$ generates a semi-group of contractions T_t on H . Moreover the solution of (5.1) is given by:

$$u(t) = T_t(u_0) + \int_0^t T_s(g) ds.$$

We define $f(t) = u(t) - u_\infty$ which satisfies:

$$\partial_t f = -Af, \quad f_{t=0} = f_0, \tag{C.1}$$

with $f_0 = u_0 - u_\infty$ and $\int_\kappa f_0(\kappa) d\kappa = 0$. To prove the weak convergence of $u(t)$ to u_∞ , we have to prove that $f(t)$ converges to zero weakly in H .

To this aim, we make an orthogonal decomposition of $f(t)$ as in the proof of Proposition 4.3: $f(t) = c(t)M + v(t)$, with $c(t) \in L^2(\Pi_\theta)$, $v(t) \in H$ and $\int_\kappa v(t) d\kappa = 0$. Taking the inner product of (C.1) with f , we get:

$$\frac{1}{2} \partial_t \|f\|_H^2 = - \int_{\kappa, \theta} \alpha^2 M \left[\partial_\kappa \left(\frac{f}{M} \right) \right]^2 d\kappa d\theta.$$

Using the decomposition of $f(t)$ and noticing that $\partial_\kappa \left(\frac{f}{M} \right) = \partial_\kappa \left(\frac{v}{M} \right)$, this equality becomes:

$$\frac{1}{2} \partial_t (\|c(t)\|_{L^2}^2 + \|v(t)\|_H^2) = - \int_{\kappa, \theta} \alpha^2 M \left[\partial_\kappa \left(\frac{v(t)}{M} \right) \right]^2 d\kappa d\theta. \tag{C.2}$$

If we apply the Gross inequality (B.9), we get:

$$\frac{1}{2} \partial_t (\|c(t)\|_{L^2}^2 + \|v(t)\|_H^2) \leq -\|v(t)\|_H^2.$$

Since $c(t)$ is bounded by $\|f_0\|_H^2$, by integrating with respect to time, we have:

$$\frac{1}{2} \|v(t)\|_H^2 \leq - \int_0^t \|v(s)\|_H^2 ds + C.$$

Using the Gronwall lemma, we deduce that $v(t)$ decays exponentially fast to zero strongly in H :

$$v(t) \xrightarrow{t \rightarrow +\infty} 0 \quad \text{in } H \text{ strong.}$$

It remains to prove the convergence of $c(t)$ to zero. We integrate (C.1) with respect to κ . This gives, using that $\int_\kappa M(\kappa) d\kappa = 1$ and $\int_\kappa v(t) d\kappa = 0$:

$$\partial_t c(t) = \partial_\theta \int_\kappa \kappa v(t) d\kappa. \tag{C.3}$$

Now if we pre-multiply by κ before integrating with respect to κ , we obtain:

$$\partial_t \int_{\kappa} \kappa v(t) d\kappa = \alpha^2 \partial_{\theta} c(t) + \partial_{\theta} \int_{\kappa} \kappa^2 v(t) d\kappa - \int_{\kappa} \kappa v(t) d\kappa. \tag{C.4}$$

We fix a time interval Δt and integrate (C.4) over this time interval. This leads to:

$$\begin{aligned} \int_{\kappa} \kappa (v(t + \Delta t) - v(t)) d\kappa &= \alpha^2 \partial_{\theta} \int_t^{t+\Delta t} c(s) ds + \partial_{\theta} \int_{\kappa} \kappa^2 \int_t^{t+\Delta t} v(s) ds d\kappa \\ &\quad - \int_{\kappa} \kappa \int_t^{t+\Delta t} v(s) ds d\kappa. \end{aligned}$$

Since $v(t)$ converges to zero in H , we have, in the sense of distributions:

$$\alpha^2 \partial_{\theta} \int_t^{t+\Delta t} c(s) ds \xrightarrow{t \rightarrow +\infty} 0. \tag{C.5}$$

Since c belongs to $L^{\infty}((0, \infty)_t, L^2(\Pi_{\theta}))$ (see (C.2)), we have $\int_t^{t+\Delta t} c(s) ds$ which belongs to $L^{\infty}((0, \infty)_t, L^2(\Pi_{\theta}))$. So there exists a subsequence such that $\int_t^{t+\Delta t} c(s) ds$ is weakly convergent in $L^2(\Pi_{\theta})$. Actually, (C.5) implies that there exists a constant function with respect to θ , depending on Δt and denoted by $L(\Delta t)$ such that

$$\int_t^{t+\Delta t} c(s) ds \xrightarrow{t \rightarrow +\infty} L(\Delta t).$$

To deduce the convergence of $c(t)$, we have to control the derivative of $c(t)$ in time. For this purpose, we rewrite:

$$\begin{aligned} \int_t^{t+\Delta t} c(s) ds &= \int_0^{\Delta t} \left(c(t) + \int_0^s \partial_t c(t+z) dz \right) ds \\ &= \Delta t c(t) + \int_0^{\Delta t} \int_0^s \partial_{\theta} \int_{\kappa} \kappa v(t+z) d\kappa dz ds. \end{aligned}$$

Using again the convergence of $v(t)$ to zero, we find:

$$\Delta t c(t) \xrightarrow{t \rightarrow +\infty} L(\Delta t),$$

or defining the constant $C = \frac{L(\Delta t)}{\Delta t}$, we have $c(t) \xrightarrow{t \rightarrow +\infty} C$ in $L^2(\Pi_{\theta})$ weak.

To complete the proof, it remains to prove that C is equal to zero. Now, since (C.1) is mass preserving i.e.:

$$\partial_t \int_{\kappa, \theta} f(t) d\kappa d\theta = - \int_{\kappa, \theta} A f(t) d\kappa d\theta = 0,$$

we have $\int_{\kappa, \theta} f(t) d\kappa d\theta = \int_{\kappa, \theta} f(0) d\kappa d\theta = 0$. Also:

$$\int_{\kappa, \theta} f(t) d\kappa d\theta = \int_{\kappa, \theta} (c(t)M + v(t)) d\kappa d\theta \xrightarrow{t \rightarrow +\infty} \int_{\theta} C d\theta 2\pi C.$$

So $C = 0$. This proves $f(t) \xrightarrow{t \rightarrow +\infty} 0$ in H weak and completes the proof. □

References

1. Aldana, M., Huepe, C.: Phase transitions in self-driven many-particle systems and related non-equilibrium models: a network approach. *J. Stat. Phys.* **112**(1/2), 135–153 (2003)
2. Aoki, I.: A simulation study on the schooling mechanism in fish. *Bull. Jpn. Soc. Sci. Fish.* **48**, 1081–1088 (1982)
3. Armbruster, D., Degond, P., Ringhofer, C.: A model for the dynamics of large queuing networks and supply chains. *SIAM J. Appl. Math.* **66**, 896–920 (2006)
4. Aw, A., Klar, A., Rasche, M., Materne, T.: Derivation of continuum traffic flow models from microscopic follow-the-leader models. *SIAM J. Appl. Math.* **63**, 259–278 (2002)
5. Bass, R.: *Diffusions and Elliptic Operators*. Springer, New York (1997)
6. Bardos, C., Santos, R., Sentis, R.: Diffusion approximation and computation of the critical size. *Trans. Am. Math. Soc.* **284**, 617–649 (1984)
7. Ben Abdallah, N., Degond, P., Mellet, A., Poupaud, F.: Electron transport in semiconductor superlattices. *Q. Appl. Math.* **61**, 161–192 (2003)
8. Bensoussan, A., Lions, J.L., Papanicolaou, G.C.: Boundary layers and homogenization of transport processes. *J. Publ. RIMS Kyoto Univ.* **15**, 53–157 (1979)
9. Brézis, H.: *Analyse Fonctionnelle*. Dunod, Paris (1983)
10. Brillinger, D.R., Preisler, H.K., Ager, A.A., Kie, J.G., Stewart, B.S.: Employing stochastic differential equations to model wildlife motion. *Bull. Braz. Math. Soc.* **33**, 385–408 (2002)
11. Camazine, S., Deneubourg, J.-L., Franks, N.R., Sneyd, J., Theraulaz, G., Bonabeau, E.: *Self-Organization in Biological Systems*. Princeton University Press, Princeton (2002)
12. Castella, F., Degond, P., Goudon, T.: Diffusion dynamics of classical systems driven by an oscillatory force. *J. Stat. Phys.* **124**, 913–950 (2006)
13. Cercignani, C., Illner, R., Pulvirenti, M.: *The Mathematical Theory of Dilute Gases*. Springer, New York (1991)
14. Couzin, I.D., Krause, J., James, R., Ruxton, G.D., Franks, N.R.: Collective memory and spatial sorting in animal groups. *J. Theor. Biol.* **218**, 1–11 (2002)
15. Degond, P.: Global existence of solutions for the Vlasov-Fokker-Planck equation in 1 and 2 space dimensions. *An. Sci. Ec. Norm. Sup.* **19**, 519–542 (1986)
16. Degond, P.: Macroscopic limits of the Boltzmann equation: a review. In: Degond, P., Pareschi, L., Russo, G. (eds.) *Modeling and Computational Methods for Kinetic Equations. Modeling and Simulation in Science, Engineering and Technology Series*, pp. 3–57. Birkhäuser, Boston (2003)
17. Degond, P., Mancini, S.: Diffusion driven by collisions with the boundary. *Asymptot. Anal.* **27**, 47–73 (2001)
18. Degond, P., Mas-Gallic, S.: Existence of solutions and diffusion approximation for a model Fokker-Planck equation. *Transp. Theory Stat. Phys.* **16**, 589–636 (1987)
19. Degond, P., Motsch, S.: Continuum limit of self-driven particles with orientation interaction. *Math. Models Methods Appl. Sci. (M3AS)* (to appear)
20. Degond, P., Motsch, S.: Macroscopic limit of self-driven particles with orientation interaction. *C. R. Acad. Sci. Paris, Ser. I* **345**, 555–560 (2007)
21. Degond, P., Zhang, K.: Diffusion approximation of a scattering matrix model of a semiconductor superlattice. *SIAM J. Appl. Math.* **63**, 279–298 (2002)
22. Degond, P., Latocha, V., Mancini, S., Mellet, A.: Diffusion dynamics of an electron gas confined between two plates. *Methods Appl. Anal.* **9**, 127–150 (2002)
23. Degond, P., Lemou, M., Picasso, M.: Viscoelastic fluid models derived from kinetic equations for polymers. *SIAM J. Appl. Math.* **62**, 1501–1519 (2002)
24. Desvillettes, L., Dolbeault, J.: On long time asymptotics of the Vlasov-Poisson-Boltzmann equation. *Commun. Partial Differ. Equ.* **16**, 451–489 (1991)
25. D’Orsogna, M.R., Chuang, Y.L., Bertozzi, A.L., Chayes, L.: Self-propelled particles with soft-core interactions: patterns, stability and collapse, *Phys. Rev. Lett.* (2006)
26. Edelstein-Keshet, L.: Mathematical models of swarming and social aggregation, invited lecture. In: *The 2001 International Symposium on Nonlinear Theory and its Applications (NOLTA 2001)*, Miyagi, Japan (Oct. 28–Nov. 1, 2001)
27. Friedrich, B.M., Julicher, F.: Chemotaxis of sperm cells. *Proc. Natl. Acad. Sci. USA* **104**, 13256–13261 (2007)
28. Gautrais, J., Motsch, S., Jost, C., Soria, M., Campo, A., Fournier, R., Bianco, S., Theraulaz, G.: Analyzing fish movement as a persistent turning walker (in preparation)
29. Golse, F., Poupaud, F.: Limite fluide des équations de Boltzmann des semiconducteurs pour une statistique de Fermi-Dirac. *Asymptot. Anal.* **6**, 135–160 (1992)

30. Gradshteyn, I.S., Ryzhik, I.M.: Tables of Integrals, Series and Products, 6th edn. Academic, New York (2000)
31. Grégoire, G., Chaté, H.: Onset of collective and cohesive motion. *Phys. Rev. Lett.* **92**, 025702 (2004)
32. Gross, L.: Logarithmic Sobolev Inequalities and Contractivity Properties of Semigroups. *Lectures Notes in Mathematics*, vol. 1563. Springer, Berlin (1992), pp. 54–88
33. Helbing, D.: Traffic and related self-driven many-particle systems. *Rev. Mod. Phys.* **73**, 1067–1141 (2001)
34. Jeanson, R., Blanco, S., Fournier, R., Deneubourg, J.L., Fourcassié, V., Theraulaz, G.: A model of animal movements in a bounded space. *J. Theor. Biol.* **225**, 443–451 (2003)
35. Jost, C., et al.: From individual to collective ant displacements in heterogenous environments. *J. Theor. Biol.* **250**, 424–434 (2008)
36. Kulinskii, V.L., Ratushnaya, V.I., Zvelindovsky, A.V., Bedeaux, D.: Hydrodynamic model for a system of self-propelling particles with conservative kinematic constraints. *Europhys. Lett.* **71**, 207–213 (2005)
37. Lions, J.L.: *Equations différentielles opérationnelles et problèmes aux limites*. Springer, New York (1961)
38. Mogilner, A., Edelstein-Keshet, L.: A non-local model for a swarm. *J. Math. Biol.* **38**, 534–570 (1999)
39. Mogilner, A., Edelstein-Keshet, L., Bent, L., Spiros, A.: Mutual interactions, potentials, and individual distance in a social aggregation. *J. Math. Biol.* **47**, 353–389 (2003)
40. Oksendal, B.: *Stochastic Differential Equations*. Springer, New York (1992)
41. Othmer, H.G., Hillen, T.: The diffusion limit of transport equations II: Chemotaxis equations. *SIAM J. Appl. Math.* **62**, 1222–1250 (2002)
42. Parrish, J.K., Viscido, S.V.: Traffic rules of fish schools: a review of agent-based approaches. In: Hemelrijk, C.K. (ed.) *Self-Organization and Complexity*. Cambridge University Press, Cambridge (2003)
43. Parrish, J.K., Viscido, S.V., Grünbaum, D.: Self-organized fish schools: an examination of emergent properties. *Biol. Bull.* **202**, 296–305 (2002)
44. Pazy, A.: *Semigroups of Linear Operators and Applications to Partial Differential Equations*. Springer, New York (1983)
45. Poupaud, F.: Diffusion approximation of the linear semiconductor equation: analysis of boundary layers. *Asymptot. Anal.* **4**, 293–317 (1991)
46. Ratushnaya, V.I., Bedeaux, D., Kulinskii, V.L., Zvelindovsky, A.V.: Collective behaviour of self-propelling particles with kinematic constraints; the relations between the discrete and the continuous description. *Physica A* **381**, 39–46 (2007)
47. Ratushnaya, V.I., Kulinskii, V.L., Zvelindovsky, A.V., Bedeaux, D.: Hydrodynamic model for the system of self-propelling particles with conservative kinematic constraints; two dimensional stationary solutions. *Physica A* **366**, 107–114 (2006)
48. Theraulaz, G., et al.: Spatial patterns in ant colonies. *Proc. Natl. Acad. Sci.* **99**, 9645–9649 (2002)
49. Topaz, C.M., Bertozzi, A.L.: Swarming patterns in a two-dimensional kinematic model for biological groups. *SIAM J. Appl. Math.* **65**, 152–174 (2004)
50. Topaz, C.M., Bertozzi, A.L., Lewis, M.A.: A nonlocal continuum model for biological aggregation. *Bull. Math. Biol.* **68**, 1601–1623 (2006)
51. Vicsek, T., Czirók, A., Ben-Jacob, E., Cohen, I., Shochet, O.: Novel type of phase transition in a system of self-driven particles. *Phys. Rev. Lett.* **75**, 1226–1229 (1995)