

The role of extensional tectonics in the Caledonides of south Norway: Reply

HAAKON FOSSEN

Statoil, GF/PETEK, N-5020 Bergen, Norway

(Received 19 April 1993; accepted 4 May 1993)

I WELCOME Andersen's interest in my paper, and appreciate the opportunity to discuss the extensional deformation in the Caledonides of southern Norway, further.

Andersen and co-workers have been studying a relatively small area of SW Norway near the Nordfjord-Sogn detachment (NSD), and have suggested that the model involving gravity-driven extensional collapse due to a detaching and descending cold orogenic root of the lithospheric mantle may apply to this area. In my paper I presented data from a much larger part of the Caledonides in south Norway, and particularly from the décollement zone which extends from the hinterland to the foreland. This regional approach is essential when large-scale orogenic models are to be discussed and evaluated. It appears that Andersen accepts the main thesis of my paper; i.e. that the contractional fabrics were consistently reworked by W- to NW-directed nappe translations and W-dipping shear zones related to horizontal extension of the Caledonian crust. It is not an *assumption*, however, that the contractional fabrics everywhere pre-date the extensional deformation, but a conclusion that was based on detailed and systematic field work and data analysis. The age relationship between the SE-directed (contractional) and WNW-directed (extensional) kinematic structures is surprisingly consistent. This conclusion is supported by much recent work in various parts of the Caledonides (Kvale 1960, Naterstad *et al.* 1973, Andresen, 1974, Brewer & Smythe 1984, Milnes & Koestler 1985, McClay *et al.* 1986, Olesen 1986, Milnes *et al.* 1988, Fossen & Rykkelid 1988, Holdsworth 1989, Powell & Glendinning 1990, Fossen 1991, 1992a,b, in press, Fossen & Rykkelid 1992a,b, Rykkelid 1992, Rykkelid & Andresen in press) and one would have to disregard the usefulness of almost any kind of kinematic indicators to argue against this conclusion.

Some of my statements are somewhat incorrectly quoted by Andersen, so the interested reader is urged to read the original paper carefully. In particular, I did not claim that the extension "chiefly was driven by divergent plate motions between Baltica and Laurentia rather than internal body forces generated within the thickened lithosphere". I wrote that the extension "was closely related to post-collisional, Lower to Middle Devonian plate divergence", and that "the exact reason

for the plate divergence is unknown, but may possibly have been triggered or enhanced by the push exerted on the plates by an earlier (pre- D_2) extensional collapse in the central parts of the orogen. The back movement would in this case take place after, rather than during an extensional collapse of the central parts". Hence, there is room in my model for the general extensional collapse model envisaged by Dewey (1988), but kinematic indicators and overprinting relationships show that this must have occurred prior to the top-to-the-west movement recorded in the décollement zone east of the area where Andersen recently has been working.

In other words, extensional deformation may have been going on in the central part of the orogen as gravity-driven top-to-the-ESE thrusting was still being recorded in the décollement zone under the Jotun and other nappes (see fig. 14b in my original paper and Fig. 1a in this paper). Perhaps the coaxial fabrics described from the basement by Andersen and co-workers formed during this stage (?), together with normal faults and shear zones in the overlying orogenic edifice (now mostly removed by erosion) and thrust motion along the décollement zone towards the foreland. But the reversal of the shear sense along the décollement zone clearly *post-dates* this deformation, as does the extension along the NSD. As pointed out before (Milnes *et al.* 1988, Fossen 1992a, in press), although neglected by Andersen, the rotation of the décollement zone by the HFSZ and the NSD must have caused the back movement of Caledonian nappes to have ceased, while the NSD continued to be active, transecting and cutting out the entire overlying orogenic wedge (a displacement of 30–45 km if the NSD were dipping 20–30° to the west).

I am surprised that Andersen does not realize the necessity of plate divergence in the light of the kinematic data presented in my paper. It is simply not possible for the orogenic wedge to move back into the central and thickened part of the collision zone unless 'space' is simultaneously being provided by some sort of divergent motion. Extensional collapse of the central part of an orogenic belt can perhaps thin the crust in this area almost to normal thickness, provided that rock volumes may flow freely toward the foreland (by thrust motion along the basal décollement zone, Fig. 1a). However, since this deformation is gravity driven, the central part




Fig. 1. Illustration of particle displacement pattern during (a) extensional collapse without plate divergence (as suggested by Andersen) and (b) the extensional deformation actually observed in the field. (a) corresponds to fig. 14(b) in my original paper (Fossen 1992a), and (b) is a modified version of fig. 14(c). NSD = Nordfjord-Sogn Detachment, HSZ = Hardangerfjord shear zone, RB = Røragen Basin, RSS = Ringerike sandstone.

can never be thinned more than to the thickness of the surrounding crust, and a gravitational collapse process alone therefore cannot cause the well documented backsliding of the orogenic wedge towards the hinterland. The only realistic explanation for this is that Baltoscandia and Laurentia experienced divergent motions at the time, possibly during, and certainly after, a possible gravitational collapse. Hence an extensional collapse model that does not take into account divergent plate motions, fails to explain the unequivocal kinematic data from the décollement zone.

It is interesting to note that Andersen himself insists on horizontal extension in the order of 180 km (Andersen *et al.* 1991). If this undocumented estimate should turn out to be correct, and if one disregards plate divergence, then this implies 180 km of gravity driven nappe translations (thrusting) towards the foreland while the central parts of the orogen experienced extension. In addition to being geologically and mechanically unsound, this would be incompatible with clear-cut kinematic data from the décollement zone, as emphasized above. Furthermore, the kinematics in Andersen's 'collapse-without-divergence' model is also incompatible with available data from the area where Andersen has been working. Somewhat simplified, one may say that Andersen's model implies a flow of particles from the elevated central zone towards the foreland (Fig. 1a). However, most of the extension is reported to have been taken up along a several kilometers wide low-angle, W-dipping shear zone (Norton 1986, Séranne & Séguret 1987, Chauvet & Séranne 1988), resulting in a displacement path in a significant part of the orogen which is opposite to that predicted by Andersen's 'collapse-without-divergence' model (Fig. 1b).

Andersen avoids discussing the discrepancies between his model and the kinematic data presented in my

original paper, and instead argues that stratigraphic data support contemporaneous thrusting in the Oslo region (foreland) and extension in western Norway. The stratigraphic control in the Devonian basins is, however, not very well constrained, nor is the calibration of the Devonian time scale (cf. Gale *et al.* 1980). In fact, given the uncertainties involved, stratigraphic and geochronological data do *not* exclude the possibility that thrusting and extension were separated in time (Fig. 2). This requires a very rapid switch from contraction to extension, as well as rapid extensional deformation, as supported by new geochronological data (Chauvet & Dallmeyer 1992). It is, however, possible that extension actually started in the collapsing internal part of the collision zone as thrusting was still going on near Oslo, as suggested by Andersen and indicated in my original paper (cf. fig. 14b). A third possibility is that the extensional deformation in the décollement zone cut up-section somewhere southeast of the Jotun Nappe, leaving an unstable orogenic ridge that subsequently collapsed to form the large, open folds and the small-scale thrust structures observed in the late Silurian Ringerike sandstone (Fig. 1b). The latter explanation would account for the small difference in age between the gentle deformation of the Ringerike sandstone and the deposition of Lowermost Devonian sediments along the MTFZ north of the study area.

A very important fact which has been omitted by Andersen is that the Lower Devonian sediments of the Røragen basin also contain plant fossils that denote an early Devonian age (Høeg 1936); i.e. a quite similar age to the basins associated with the MTFZ and older than the Devonian sediments of SW Norway. The extensional Røragen basin is also of the same age as the age of the thrusting of the Ringerike sandstone suggested by Andersen. This casts additional doubt on Andersen's idea that most or all of the extension in the hinterland




Fig. 2. Black boxes represent minimum age of youngest sediments involved in D_1 thrusting, and Nd/Sm ages dating the D_1 continent-continent collision. White boxes represent radiometric ages or lower age of sediments related to D_2 . The height of the boxes represents uncertainties from calibration of the time scale (cf. Gale *et al.* 1980), uncertainties related to fossil evidence (see Steel *et al.* 1985 and references therein for the ORS), or error limits/spread in radiometric ages. MTFZ = Møre-Trøndelag fault zone. See text for discussion.

was accompanied by thrusting in the foreland, since the extensional Røragen basin is situated close to the foreland.

One should also keep in mind that the MTFZ acted as a strike-slip zone both during the late stages of contraction and during the Devonian extension. The geotectonic implication of the related Devonian basins is therefore less obvious, since extensional basins tend to develop in conjunction with strike slip faults whether they are situated in an extensional or contractional regional setting. Perhaps a key point in this discussion is to be aware of the fact that extensional deformation must have been an active internal process within the upper plate (orogenic wedge) during the Scandian contraction as well as during the later extension, and extensional basins formed on the upper part of an orogenic wedge can therefore not be taken as evidence of large-scale extensional tectonics (e.g. Platt 1986). It is therefore possible that some of the Devonian sediments are older than the extensional deformation recorded in the deeper part of the Caledonian crust, and presently preserved in the hangingwall of a later extensional detachment that is not directly related to the basin formation. It is very important to be aware of the difference in significance between kinematic data from the upper plate and those from the basal décollement between the basement and the orogenic wedge. The basal décollement recorded the bulk direction of displacement of the wedge, and therefore contains the critical information about the general kinematic and tectonic history of the orogen.

This discussion emphasizes the need and usefulness, but also the complexities involved in integrating kinematic, geochronologic, tectonic and stratigraphic data on a regional as well as local scale to recognize and constrain the evolution of orogenic belts. In the Caledonides, the switch from contractional to extensional tectonics happened so quickly that both stratigraphy and geochronology are on the edge of resolution, and the kinematic-structural evidence seems most convincing. We are lucky to have a significant part of the décollement zone with its kinematic indicators preserved more or less continuously from the foreland to the hinterland, and should acknowledge the significance of data obtainable from this zone. I feel that the best way to better constrain some of the issues discussed here may be to try and date the D_1 and D_2 deformation fabrics in this zone using various modern high-precision analytical methods.

REFERENCES

- Andersen, T. B., Jamtveit, B., Dewey, J. F. & Swenson, E. 1991. Subduction and exhumation of continental crust: major mechanisms during continent-continent collision and orogenic extensional collapse, a model based on the south Norwegian Caledonides. *Terra Nova* **3**, 303–310.
- Andresen, A. 1974. Petrographic and structural history of the Caledonian rocks north of Haukelisæter, Hardangervidda. *Norges geol. unders. Bull.* **314**, 1–52.
- Brewer, M. S. & Smythe, D. K. 1984. MOIST and the continuity of crustal reflector geometry along the Caledonian–Appalachian orogen. *J. geol. Soc. Lond.* **141**, 105–120.
- Chauvet, A. & Dallmeyer, R. D. 1992. $^{40}\text{Ar}/^{39}\text{Ar}$ mineral dates related to Devonian extension in the southwestern Scandinavian Caledonides. *Tectonophysics* **210**, 155–177.
- Chauvet, A. & Séranne, M. 1988. Microtectonic evidence of Devonian extensional westward shearing in southwest Norway. In: *The Caledonian and Related Geology of Scandinavia* (edited by Gayer, R.). Graham and Trotman, London.
- Dewey, J. F. 1988. Extensional collapse of orogens. *Tectonics* **7**, 1123–1139.
- Fossen, H. 1991. Evidence for post-contractional crustal instabilities in the Caledonian orogenic wedge of South Norway. (Abs.) *Terra Nova* **3**, 15.
- Fossen, H. 1992a. The role of extensional tectonics in the Caledonides of South Norway. *J. Struct. Geol.* **14**, 1033–1046.
- Fossen, H. 1992b. Devonian extensional deformation in the Caledonian orogen, southern Norway. Unpublished Ph.D. thesis, University of Minnesota.
- Fossen, H. In press. Structural evolution of the Bergsdalen Nappes, Southwest Norway. *Norges geol. unders. Bull.*
- Fossen, H. & Rykkelid, E. 1988. Skjøernesbevegelser i Bergensområdet—foreløpige resultater. *Geolognytt* **22**, 25.
- Fossen, H. & Rykkelid, E. 1992a. The interaction between oblique and layer-parallel shear in high-strain zones: observations and experiments. *Tectonophysics* **207**, 331–343.
- Fossen, H. & Rykkelid, E. 1992b. Post-collisional extension of the Caledonide orogen in Scandinavia: structural expressions and tectonic significance. *Geology* **20**, 737–740.
- Gale, N. H., Beckinsale, R. D. & Wedge, A. J. 1980. Discussion of a paper by McKerrow, Lambert & Chamberlain on the Ordovician, Silurian and Devonian time-scale. *Earth & Planet. Sci. Lett.* **51**, 9–17.
- Holdsworth, R. E. 1989. Late brittle deformation in a Caledonian ductile thrust wedge: new evidence for gravitational collapse in the Moine Thrust sheet, Sutherland, Scotland. *Tectonophysics* **170**, 17–28.
- Høeg, O. A. 1936. Norges fossile fauna. *Naturen* **7–21**, 47–64.
- Kvale, A. 1960. The nappe area of the Caledonides in western Norway. *Norges geol. unders. Bull.* **212e**, 21–43.
- McClay, K. R., Norton, M. G., Coney, P. & Davis, G. H. 1986. Collapse of the Caledonian orogen and the Old Red Sandstone. *Nature* **323**, 147–149.
- Milnes, A. G. & Koestler, A. G. 1985. Geological structure of Jotunheimen, southern Norway (Sognefjell–Valdres cross-section). In: *The Caledonide Orogen—Scandinavia and Related Areas* (edited by Gee, D. G. & Sturt, B. A.). J. Wiley & Sons, New York, 457–474.
- Milnes, A. G., Dietler, T. N. & Koestler, A. G. 1988. The Sognefjord northshore log—a 25 km section through Caledonized basement in western Norway. *Spec. Publ. Norges Geol. Unders.* **3**, 114–121.
- Naterstad, J., Andresen, A. & Jorde, K. 1973. Tectonic succession of the Caledonian nappe front in the Haukeli–Røldal area, southwest Norway. *Norges geol. unders. Bull.* **292**, 1–20.
- Norton, M. 1986. Late Caledonian extension in western Norway: a response to extreme crustal thickening. *Tectonics* **5**, 192–204.
- Norton, M. 1987. The Nordfjord-Sogn Detachment, W. Norway. *Norsk Geol. Tidsskr.* **67**, 93–106.
- Olesen, N. Ø. 1986. Three thrust sheets on Hornsnipa, Jotun Nappe Complex, West Norway. *Norges geol. unders. Bull.* **404**, 55–66.
- Platt, J. P. 1986. Dynamics of orogenic wedges and the uplift of high-pressure metamorphic rocks. *Bull. geol. Soc. Am.* **97**, 1037–1053.
- Powell, D. & Glendinning, N. R. W. 1990. Late Caledonian extensional reactivation of a ductile thrust in NW Scotland. *J. geol. Soc. Lond.* **147**, 979–987.
- Rykkelid, E. 1992. Contractional and extensional structures in the Caledonides. Dr. scient thesis, University of Oslo.
- Rykkelid, E. & Andresen, A. In press. Late-Caledonian extension in the Ofoten area, northern Norway. *Tectonophysics*.
- Séranne, M. & Séguret, M. 1987. The Devonian basins of western Norway: Tectonics and kinematics of an extending crust. In: *Continental Extensional Tectonics* (edited by Coward, M. P., Dewey, J. F. & Hancock, P. L.). *Spec. Publs geol. Soc. Lond.* **28**, 537–548.
- Steel, R., Siedlecka, A. & Roberts, D. 1985. The Old Red Sandstone basins of Norway and their deformation: a review. In: *The Caledonide Orogen—Scandinavia and Related Areas* (edited by Gee, D. G. & Sturt, B. A.). J. Wiley & Sons, New York, 293–315.