

Structural evolution of different tectonic units across the Austroalpine–Penninic boundary in the middle Orco Valley (Western Italian Alps)

Ivano Gasco^{a,*}, Marco Gattiglio^a, Alessandro Borghi^b

^a Earth Sciences Department, University of Torino, Via Valperga Caluso 35, I-10125 Torino, Italy

^b Department of Mineralogical and Petrological Sciences, University of Torino, Via Valperga Caluso 35, I-10125 Torino, Italy

ARTICLE INFO

Article history:

Received 30 November 2007

Received in revised form

27 October 2008

Accepted 20 November 2008

Available online 3 December 2008

Keywords:

Structural evolution

Tectonic coupling

Exhumation

Shear zone

Western Italian Alps

ABSTRACT

A structural study along the Orco Valley, which represents a geological section through the Alpine belt (western Italy), is reported. A Lower Tectonic Element, composed of units with eclogite facies relics (Gran Paradiso and Zermatt–Saas-like Unit) and an Upper Tectonic Element, made of greenschist facies units (Combin-like Unit and Gneiss Minuti Complex of the Sesia–Lanzo Zone) have been distinguished. These two elements show a different main Alpine tectono-metamorphic event (D_1) and are separated by a mylonitic contact (D_2) developed under greenschist facies conditions (Orco Shear Zone). This regional scale shear zone was responsible for the exhumation of the deeper eclogitic units, and has been inferred as the continuation of the Combin Fault south of the Aosta Valley. From D_3 onward, the same tectono-metamorphic evolution was recorded in all units. The following extensional crenulation cleavage (D_4) occurs mostly along a NNE–SSW deformation zone about 1 km thick at the base of the Upper Tectonic Element which leads to a further exhumation of the Lower Tectonic Element.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

The axial portion of the Western Italian Alps (Austroalpine and Penninic Domains) represents a fossil subduction complex developed during the Alpine orogenic cycle. In the last 20 years, many geological transects (e.g. Pognante et al. 1987; Inger and Ramsbotham, 1997; Bucher et al., 2004) have improved our knowledge of the structural and metamorphic framework of the Western Alps. This has brought a better insight into the geodynamic evolution of the Alpine mountain belt and has led many workers to suggest different exhumation mechanisms for HP rocks (Platt, 1993; Bal-lèvre and Merle, 1993; Reddy et al., 1999). However, some questions about the timing of tectonic coupling between oceanic and continental units still remain controversial. Lapen et al. (2007), on the basis of geochronological data and PTt path analysis, have proposed that the Monte Rosa–Gran Paradiso Massifs and the Zermatt–Saas Unit were coupled during eclogite-facies tectono-metamorphic stage and have been exhumed together by a buoyancy-driven process. Instead Kassem and Ring (2004) proposed that the Gran Paradiso has been coupled to the Zermatt–Saas Unit early during subduction under brittle conditions. Finally Pleuger et al. (2005)

suggested tectonic coupling of the Monte Rosa Massif and the Zermatt–Saas Unit before the greenschist facies metamorphic re-equilibration.

In this paper we present the tectono-metamorphic evolution of a geological transect through different palaeogeographic units along the middle Orco Valley, Western Italian Alps (Fig. 1). The Orco Valley represents a poorly studied key-area to unravel the structural and metamorphic evolution of oceanic versus continental tectonic units because it is a natural section through the Alpine mountain belt. Indeed, the valley trends E–W and is perpendicular to both tectonic and lithological boundaries, and to the regional metamorphic foliation.

This work is based on structural analysis of an area of 35 km² and takes into consideration three tectonic units differing in palaeogeographic position and Alpine metamorphic evolution. From W to E and in ascending structural order, they are: the continental crust of the Gran Paradiso Unit, generally attributed to the European palaeo-margin, the Piedmont Zone that represents the remnants of the Tethys Ocean, and the Sesia–Lanzo Zone belonging to the African palaeo-margin, also known as Adria microplate (Dal Piaz, 1999; Schmid et al., 2004). Detailed structural and petrographic analysis established new insights into the relative timing of the tectonic contacts between the different studied units. In particular, the time of coupling between oceanic and continental crust units and of eclogite versus greenschist facies units has been established.

* Corresponding author. Tel.: +39 011 670 5335; fax: +39 011 670 5339.
E-mail address: ivano.gasco@unito.it (I. Gasco).




Fig. 1. (a) Tectonic map of the Western Alps (modified after Schmid et al., 2004). (b) Simplified geological map of the Gran Paradiso Massif and surrounding units (modified after Compagnoni and Lombardo, 1974).

2. Geological background

2.1. The Gran Paradiso Unit (GP)

The Gran Paradiso Unit (Fig. 1) is mainly composed of augen-gneisses (Gneiss Occhiadini Complex, Compagnoni et al., 1974) derived from Permian (270 ± 5 Ma) porphyric granitoids (Bertrand et al., 2000; Ring et al., 2005), intruded into a metasedimentary high-T Variscan complex (Gneiss Minuti Complex, Compagnoni et al., 1974). Intrusive contacts are reported at the Lake of Teleccio, where relicts of the intrusive Variscan structures and sillimanite paragneisses have been preserved from Alpine reworking (Callegari et al., 1969; Compagnoni and Prato, 1969).

The metasedimentary rocks have been reworked by Alpine orogeny that transforms them into a polymetamorphic complex, made of paragneisses and micaschists with lenses of eclogites, interpreted as pre-Alpine amphibolites (Compagnoni and Lombardo, 1974) or Variscan gabbros (Biino and Pognante, 1989). Eclogitic assemblages in the metabasites have been used to estimate minimum PT conditions for the eclogitic event at $P > 12$ – 14 kbar and T of 500 – 550 °C (Ballèvre, 1988; Brouwer et al., 2002). Within orthogneisses, layers of whiteschists that developed in metasomatic shear zone during the eclogitic event also occur (Compagnoni and Lombardo, 1974). For these rocks PT conditions of 21 – 23 kbar and 540 – 570 °C have been reported by Wei and Powell (2003, 2004) and Meffan-Main et al. (2004). Conditions of 18 – 20 kbar and 490 °C are reported for the Gneiss Minuti Complex (Le Bayon et al., 2006). The high- P event is followed by a late stage of re-equilibration at 4 – 6 kbar and 500 – 550 °C, interpreted as a re-heating stage after an initial cooling during decompression (Borghini et al., 1996; Brouwer et al., 2002, 2004).

Geochronological data, obtained with Rb–Sr microsampling on white mica, suggest that the eclogite-facies event occurred at

43 ± 0.5 Ma (Meffan-Main et al., 2004), whereas the greenschist-facies metamorphism occurred at about 34 – 38 Ma (Rb–Sr on white mica, Freeman et al., 1997; Meffan-Main et al., 2004). The final stages of exhumation are recorded by U–Pb fission tracks on zircon (30 ± 1 Ma at 225 ± 25 °C) and apatite (20 – 24 Ma at 100 ± 20 °C) (Hurford and Hunziker, 1989).

2.2. The Piedmont Zone

The Piedmont Zone, representing the remnant of the Tethys Ocean, consists of two main zones, the lower and the upper one, respectively known as the Zermatt–Saas Unit and the Combin Unit in the southern Valais and Aosta Valley (Beaerth, 1967; Dal Piaz, 1974, 1999; Dal Piaz and Ernst, 1978). The Zermatt–Saas Unit is composed of slivers of oceanic crust, mainly represented by serpentinites with minor peridotites, including discontinuous bodies of Mg–Al to Fe–Ti rich metagabbros and overlaid by metabasalts. Locally a thin metasedimentary cover composed of impure quartzites, micaschists, marbles and calcschists is preserved. Mafic and metasedimentary rocks display a well-preserved eclogite facies metamorphism.

The Combin Unit is made of calcschists with intercalations of metabasalts and serpentinites; all lithologies show a mainly greenschist facies metamorphic overprinting, with blueschist facies relicts.

PT conditions for different tectonic slivers belonging to Zermatt–Saas decrease from north ($P > 15$ – 20 kbar and 550 – 600 °C) to south (10 – 18 kbar and 450 – 500 °C; for review see: Spalla et al., 1996; Desmons et al., 1999a,b). The occurrence of coesite is reported for the Cignana Unit, in Valtournenche, indicating P of 26 – 28 kbar and T of 600 °C (Reinecke, 1991, 1998). Messiga et al. (1999) reported conditions of 24 kbar at 600 ± 50 °C for the Monviso ophiolites. The greenschist facies metamorphic overprinting

developed at 425–450 °C and 4–5 kbar and is concentrated along top to SE shear zones (Barnicoat et al., 1995; Cartwright and Barnicoat, 2002). The high-*P* event is dated at 40–50 Ma (Rb–Sr white mica, Duchêne et al., 1997; U–Pb zircon with SHRIMP, Rubatto et al., 1998; Sm–Nd garnet, Amato et al., 1999; Rb–Sr and ⁴⁰Ar–³⁹Ar white mica, Dal Piaz et al., 2001; Rb–Sr white mica, Cartwright and Barnicoat, 2002) while greenschist re-equilibration spread from 36 to 42 Ma (Rb–Sr whole rock-white mica, Amato et al., 1999; Cartwright and Barnicoat, 2002).

In the Combin Unit the high-*P* event developed under blueschist facies conditions (12–13 kbar and 425–475 °C, Cartwright and Barnicoat, 2002) while the re-equilibration event developed under greenschist facies conditions (*P* < 9 kbar and *T* of 300–450 °C, Reddy et al., 1999) at around 34–38 Ma (Rb–Sr white mica, Inger and Ramsbotham, 1997).

2.3. The Sesia–Lanzo Zone (SL)

The Sesia–Lanzo Zone consists of a pre-Alpine basement thrust above the Piedmont Zone and involved in the orogenic accretionary wedge. It is made of three tectonic units that, from top to bottom and from E to W, consist of the Eclogitic Micaschists Complex (EMC), the II Dioritic-Kinzigitic Zone (II DK) and the Gneiss Minuti Complex (GMC) (Compagnoni et al., 1977; Pognante, 1989). The EMC is a high-*T* Variscan complex made of paragneisses, marbles, amphibolites and granulites intruded by late-Variscan gabbros and granitoids, that developed widespread eclogite facies assemblages during the Alpine orogenic cycle (Lardeaux and Spalla, 1991). The II DK is a high-*T* Variscan metasedimentary complex little affected by high-*P* and greenschist facies Alpine overprinting. The GMC also comprises pre-Alpine rocks pervasively transformed into greenschist facies lithologies: mainly orthogneisses, subordinate micaschists and metabasites, with minor Mesozoic calcschists. On the east side of the GMC some relicts of blueschist facies conditions are preserved (Spalla et al., 1996).

The EMC reached conditions of 15–18 kbar and 550–620 °C (Pognante et al., 1980, 1987; Lardeaux et al., 1982; Pognante, 1989; Lardeaux and Spalla, 1991) dated 65–80 Ma (Rb–Sr white mica, Inger et al., 1996; Duchêne et al., 1997; Rubatto et al., 1999). The GMC, instead, has been pervasively equilibrated at *P* < 13 kbar and 450–500 °C (Williams and Compagnoni, 1983; Pognante et al., 1987; Pognante, 1989; Spalla et al., 1996) at about 37–39 Ma (Hunziker et al., 1993; Inger et al., 1996).

3. Tectono-stratigraphic setting

According to their different structural and metamorphic evolution, the tectonic units outcropping in the study area (Fig. 2) have been grouped in two main Tectonic Elements. From NW to SE they are: (i) a Lower Tectonic Element (LTE) preserving relicts of eclogite facies metamorphism and consisting of the Gran Paradiso Unit and oceanic tectonic slices referred to the Zermatt–Saas Unit according to lithostratigraphic and metamorphic affinities; (ii) an Upper Tectonic Element (UTE) made of greenschist facies units devoid of eclogitic relicts consisting of a Combin-like unit and the Gneiss Minuti Complex of the Sesia–Lanzo Zone. These two Tectonic Elements are separated by a tectonic contact of regional importance. A detailed summary of the metamorphic assemblages and the tectono-metamorphic relationships of the different studied units are respectively reported in Tables 1 and 2.

3.1. Lower Tectonic Element (LTE)

In the Lower Tectonic Element the GP continental crust represents the deepest tectonic unit of the study area, over-thrusted by

a tectonic mélange of oceanic affinity referred to as the Zermatt–Saas Unit.

The Gran Paradiso Unit consists of orthogneisses, a poly-metamorphic basement and rare Mesozoic cover (quartzites and dolomitic marbles) preserved near the tectonic contact with oceanic units. The orthogneisses are heterogeneously deformed ranging from isotropic rocks with magmatic structures perfectly preserved, to augen-gneisses and flaser-gneisses. Locally thin layers of whiteschists which grade to augen-gneisses are present. The polymetamorphic basement is mainly composed of micaschists hosting metabasites bodies. They consist of up to 1 m-sized lenses or layers of fine grained and felsic eclogites, and up to 100–150 m thick fine to medium-grained foliated metagabbros, that locally preserve the eclogite facies assemblage. A late metamorphic greenschist facies overprinting developed under static conditions in both micaschists and metabasites.

The oceanic units consist of a tectonic mélange formed by ultrabasic and basic rocks with minor carbonatic–pelitic cover sequences. The ultrabasic bodies mainly consist of serpentinites that envelop 10–100 m-thick lenses of metamorphosed spinel-lherzolites, showing a well developed tectonic foliation. Small bodies of metagabbros rarely occur within the ultrabasic rocks. The basic bodies consist of epidote-albite amphibolites and of metagabbros, which locally preserve eclogitic relicts.

Cover sequences are subordinate with at most 5–10 m thickness and are represented by marbles, silicate marbles, calcschists, micaschists and impure quartzites. These lithologies are transitional to each other and locally are interlayered at the decimetre-scale.

3.2. Upper Tectonic Element (UTE)

The Upper Tectonic Element consists of a lower unit showing Combin affinity and an upper unit of continental basement rocks referred to the Gneiss Minuti Complex of the Sesia–Lanzo Zone.

The Combin-like unit is mainly composed of calcschists with up to 50–100 m-thick bodies of greenstones and serpentinites/ophicalcites. The Gneiss Minuti Complex represents an heterogeneous continental crust unit with subordinate Mesozoic cover. It mainly consists of orthogneisses whose protholites range from monzodiorite to granodiorite and minor leuco-gneisses, greenstones, metagabbros and micaschists. The latter have been interpreted as a pre-Alpine basement from the presence of relict red biotite. The Mesozoic cover is made of rare quartzites and of calcschists interlayered with thin meta-carbonatic and meta-pelitic levels.

3.3. Mylonitic rocks along tectonic contacts

Mylonitic rocks occur along the main tectonic contacts between the units. Their peculiar mineral assemblages are indicative of the metamorphic conditions under which they were developed and were of great importance in reconstructing the tectono-metamorphic evolution of the area.

The tectonic contact between the Gran Paradiso and the Zermatt–Saas-like Unit is characterized by the presence of sheared Mesozoic cover belonging to both units. Where metasediments are absent, serpentinites become mylonitic (Srp + Mg–Chl + Ol + Di; mineral abbreviations according to Kretz, 1983), eclogites are re-equilibrated under blueschist facies conditions (Gln + Ep + Bt + Spn), while orthogneisses are transformed into micaschists (Qtz + Wm + Mg–Chl + Ab + Mg–Cld). On the southern side of the Orco Valley, a mylonitic foliation that envelops eclogite boudins has been observed. This foliation is made of Act–Chl–Wm–schists and preserves portions of an older foliation defined by blue Am + Chl + Wm which suggests that the tectonic contact developed under blueschist facies conditions and was reworked during later decreasing *P*. The thickness of the deformation zone varies between 1




Fig. 2. Simplified geological map of the study area located at the western border of the Gran Paradiso Unit.

and 5 m where competent rocks are in contact, to 10–50 m where Mesozoic cover is present.

The tectonic contact between the Gneiss Minuti Complex of the Sesia–Lanzo Zone and the Combin-like unit is marked by mylonitic rocks deriving from both units and developed under greenschist facies conditions. Locally Chl-schists (at the expense of greenstones) and Tlc-schists (at the expense of metabasites or serpentinites) are also present.

Finally, the regional tectonic contact between eclogitic LTE and greenschist UTE is marked by a mylonitic zone which ranges from some meters up to 50 m in width. Along this shear zone mylonitic

calcschists and gneisses are widespread and locally Chl–Act-schists (derived from metabasites and serpentinites) are developed, which envelop blocks of eclogites, metasediments and Sesia–Lanzo gneisses. This regional scale shear zone represents the boundary between the eclogite facies units in the footwall and the greenschist facies units in the hangingwall and has been defined as Orco Shear Zone.

4. Structural and metamorphic evolution

Three different Alpine tectono-metamorphic events at the regional scale (D_1 , D_2 , D_3), followed by two ductile to brittle events

Table 1
Evolution of the mineral assemblages respect to the structural and metamorphic setting in the main lithologies occurring in the different tectonic units.

Structural evolution	Metamorphic conditions	Lower tectonic unit (eclogite facies relics)			Metamorphic conditions		Upper tectonic unit (only greenschist facies assemblages)		
		Gran Paradiso Unit		Zermatt–Saas Unit		Piedmont Zone	Combin Unit	Piedmont Zone	Sesia–Lanzo Zone
Pre-Alpine		Micaschists	Metabasites	Orthogneisses (whiteschists)	Serpentinities (metaperidotites)	Metagabbros	Metabasites	Greenstones	Gneisses (metagranitoids)
Pre-S ₁	Eclogite facies	Enclaves in orthogneisses Grt, Gln, Wm, Qtz, Rt	Magmatic structures Omp, Grt, Gln, Zo, Wm, Rt	Kfs, Qtz, Bt, Pl (Wm, Qtz, Chl, Cld, Ky, Tlc, Grt)	(Ol, Cpx, Opx, Spl) (Ol, Chl, Am, Mag)	Magmatic structures Omp, Gln, Grt	–	–	Magmatic structures Wm, Qtz, Chl, Ep, Act; (Zo, Rt)
S ₁	Middle-P greenschist to epidote–albite amphibolite facies	Wm, Qtz, Hbl, Ab, Grt, Zo, Rt	Ca–Na Am, Zo, Wm, Chl, Ab, Rt	Wm, Qtz, Ab, Ep, Spn (Qtz, Wm, Chl)	Srp, Chl, Ol, Di	Am, Zo, Wm, Chl, Rt	Greenschist facies	Act, Ab, Ep, Chl, Spn	Wm, Qtz, Ab, Ep/Czo, Chl, Act, Spn
S ₂	Greenschist facies	Qtz, Wm, Chl, Ab, Spn	–	–	Srp	–	–	–	Wm, Chl, Act
S ₃	Low-grade metamorphic conditions	–	–	–	Srp	–	–	–	–
S ₄ (ECC)	Ductile to brittle	–	–	–	Srp	–	–	–	–
		Qtz, Wm, Bt	–	–	–	–	–	–	Qtz, Wm, Bt, Spn

(D₄, D₅) have been distinguished. The study area is marked by a regional transposed foliation concordant with the lithological boundaries, dipping 40–50° E to SE. However, the regional foliation developed under different metamorphic conditions (Table 2) and at a different time to the two main tectonic elements. In the following sections, we will refer to S₁ as the main regional foliation of each tectonic element, which is not synchronous across the whole area. The mineral assemblages related to the different deformation events are summarized in Table 1 for the main rock types.

4.1. D₁ structures and metamorphism in the Lower Tectonic Element

In the Gran Paradiso Unit S₁ foliation dips 35–70° SE with some poles in the opposite quadrant (Fig. 3a). D₁ folds are isoclinal to rootless in the paragneisses, isoclinal and asymmetric in metaplates within orthogneisses. At the microscale, D₁ folds transpose an early pre-S₁ foliation rarely preserved in relict hinges. Fold axes (A₁; we refer to A_n as fold axes of D_n) show similar orientation to stretching lineations (L₁) that plunge mainly 10–45° towards 050–095° with some scattering due to subsequent reorientation. L₁ is defined by oriented K-feldspar porphyroclasts in the augen-gneisses, by quartz + white mica + hornblende in paragneisses and by Ca–Na amphibole + zoisite in metabasites. D₁ is associated with a top to W sense of shear, which is proved by widespread σ-shaped K-feldspar porphyroclasts in orthogneisses (Fig. 4a). The occurrence of non-linear intersection lineations on S₁ foliation planes according to the parallelism between L₁ and A₁ on the regional scale suggests that the D₁ event mainly developed under a simple-shear regime, causing non-cylindrical folding (Fig. 4b).

In the main lithologies of the GP (paragneisses, metabasites, orthogneisses) the S₁ regional foliation (Fig. 5a,b) is defined by metamorphic assemblages (Table 1) indicative of upper-greenschist to epidote–albite amphibolite facies conditions (subsequently referred to as medium PT conditions), following pre-S₁ event/events (Fig. 5b) developed under eclogite facies conditions.

In the Zermatt–Saas-like Unit, the main foliation S₁ (Fig. 3b) dips towards the SE quadrant with a more obvious scattering with respect to the Gran Paradiso Unit, probably due to the presence of great volumes of serpentinites strongly and easily deformed by subsequent events. Folds are isoclinal in eclogites and rootless in serpentinites (Fig. 5c) and metasediments. The mineral lineations are defined by Ca–Na amphibole + zoisite in metagabbros and in re-equilibrated eclogites and by magnetite aggregates in serpentinites. Both fold axes and stretching lineations plunge to E and show a substantial parallelism to those in the Gran Paradiso Unit (Fig. 3b). Locally a well-preserved pre-S₁ eclogitic foliation occurs in metabasites and metagabbros. In meta-peridotites a relic foliation defined by olivine + chlorite + amphibole + magnetite is preserved. The following D₁ event developed a new axial surface defined by medium PT conditions.

The tectonic contact between the Gran Paradiso and the Zermatt–Saas units was deformed by D₁ top to W non-cylindrical folds (Figs. 6–8, cross-section B–B'). The presence of Mg–chlorite + albite + Mg–chloritoid micaschists and glaucophane + epidote + rutile blueschists along the contact suggests tectonic coupling under blueschist facies conditions.

4.2. D₁ structures and metamorphism in the Upper Tectonic Element

In both units of the Upper Tectonic Element the main regional foliation S₁ (Fig. 3d–f) dips 25–65° SE with some planes dipping in the opposite direction owing to the following deformation phases. D₁ folds (Fig. 4c) are isoclinal to rootless with axes plunging 25–60° towards 060–110° which are sub-parallel to the stretching

Table 2
Sketch of the tectono-metamorphic evolution of the studied units.

	Lower Tectonic Element (eclogite facies relics)		Upper Tectonic Element (only greenschist facies assemblages)	
	Gran Paradiso Unit	Zermatt-Saas Unit	Combin Unit	Sesia-Lanzo GMC
pre-D ₁	eclogite facies	eclogite facies	greenschist facies	greenschist facies
old tectonic contacts	blueschist facies		greenschist facies	
D ₁	middel-P greenschist to epidote-albite amphibolite facies		greenschist facies	
D ₂	greenschist facies		greenschist facies	
OSZ late-D ₂	greenschist facies			
D ₃	low-grade metamorphic conditions			
D ₄ (ECC)	ductile (low-grade metamorphic conditions) to brittle			




Fig. 3. Equiareal projection (lower hemisphere) of the main structural elements characterizing the four deformation phases identified. The mean surface (e.g.: 127/34) of foliation or axial plane is indicated. A_n: fold axis; L_n: stretching lineation; S_n: foliation or axial surface. For discussion see text.



Fig. 4. Meso-scale structures representing the four deformation phases distinguished in the study area: (a) σ -shaped Kfs porphyroclasts in orthogneisses of the Gran Paradiso Unit showing a top to W sense of shear; (b) S_1 surface in the Gran Paradiso paragneisses with a curved A_1 (intersection lineation), indicating a non-cylindrical folding; (c) D_1 isoclinal folds in the Sesia–Lanzo gneisses transposing leuco-gneisses layers; (d) D_2 fold in the Sesia–Lanzo gneisses folding a D_1 isoclinal fold; (e) D_3 open folds in the Gran Paradiso paragneisses folding a tight to isoclinal D_2 fold; (f) D_3 box fold in the Combin-like calcschists with S_3 surfaces defined by a spaced and discontinuous reverse-slip C' types shear bands; (g) discontinuous Extensional Crenulation Cleavage in the Zermatt–Saas serpentinites; (h) S–C type shear bands showing a top to E sense of shear in mylonitic calcschists along the tectonic contact between the Upper Tectonic Element and the Lower one.

lineations, defined by quartz + white mica in calcschists and by quartz + albite + white mica \pm actinolite in gneisses.

In both the Combin-like Unit (Fig. 5d,e) and the Gneiss Minuti Complex of the Sesia–Lanzo Zone (Fig. 5f,g), the main foliation S_1 is underlined by greenschist facies conditions. Pre- S_1 relic foliations

are preserved both in the Sesia–Lanzo gneisses and in the Combin calcschists and are also defined by greenschist facies assemblages. The D_1 event deforms the mylonitic contacts between the two tectonic units of the UTE and is mainly responsible for their lithological repetition at the regional scale (Figs. 2, 7 and 8).




Fig. 5. Microstructures representing the four deformation phases distinguished in the study area: (a) D_1 rootless fold in the Gran Paradiso whiteschists with a well preserved pre- S_1 foliation; (b) D_2 asymmetric fold in the Gran Paradiso paragneisses which developed an S_2 foliation; (c) D_1 rootless fold in the Zermatt–Saas serpentinites defined by a magnetite layer deformed by a D_2 closed fold; (d) D_2 discrete crenulation cleavage in the Combin calcschists; (e) D_4 spaced Extensional Crenulation Cleavage (ECC) in the Combin-like calcschists showing a millimetric displacement; (f) rootless D_1 isoclinal folds in the Sesia–Lanzo gneisses preserving a pre- S_1 foliation; (g) D_1 rootless fold preserving a pre- S_1 foliation in the Sesia–Lanzo micaschists, S_1 is cut by D_4 ECC; (h) mylonitic foliation in the Sesia–Lanzo gneisses along the tectonic contact between the LTE and the UTE cut by D_4 ECC.

4.3. D_2 structures and metamorphism

The D_2 event deforms D_1 structures in all tectonic units, but its structural imprinting affected the LTE and UTE differently. Indeed, in the former, D_2 structures are developed only at the mesoscale

while, in the latter, macroscale folds are also present and D_2 structures are very common. D_2 develops polyharmonic and asymmetric folds that show close to isoclinal profiles, but an axial plane foliation rarely occurs at the mesoscale (Fig. 4d,e). Axial planes dip 30–60° E to SE while the fold axes are scattered on the




Fig. 6. Structural and metamorphic relationships between the Gran Paradiso orthogneisses and the Zermatt–Saas eclogites: (a) orthogneisses show a well-developed S_1 foliation while metabasites preserve a pre- S_1 foliation; (b) eclogite facies pre- S_1 foliation is preserved in metabasites and is cut by mylonites.

average maximum circle of the foliation planes (Fig. 3g,h) showing the presence of non-cylindrical structures developed during non-coaxial deformation. Therefore, A_2 fold axes show a variable angle respect to the stretching lineations L_2 which plunge concordantly E. The asymmetric Z folds (looking to N) present on the long limbs have an enveloping surface dipping to the E more than S_2 surfaces. These features of the meso- to macroscale D_2 structures indicate a top to E sense of shear (Fig. 8 cross sections A–A', C–C') according to the structural relationship described by Inger and Ramsbotham (1997) for the Eclogitic Micaschists Complex of the Sesia–Lanzo Zone outcropping to the E of the study area.

At the microscopic scale S_2 developed under greenschist facies conditions and is defined by a crenulation cleavage (Fig. 5b–d) in all lithologies of the studied units (Table 1).

4.4. D_3 structures and metamorphism

The D_3 phase developed box folds with open to close profiles which indicate a prevalent top to SE asymmetry. Axial planes dip mainly NW by a range of inclination from 30° to sub-vertical with some data in the opposite quadrant, indicating polyclinal profile typical of box folds (Figs. 3i and 4f). Rarely C'-type conjugate shear bands, with reverse-slip movement were developed at the mesoscale in calcschists (Fig. 4f), serpentinites and subordinately in the Sesia–Lanzo gneisses, with stretching lineations L_3 defined by quartz and calcite fibres and by striations. L_3 on C'-type surfaces and shear band geometry indicate both top to NW and top to SE sense of shear pointing to a NW–SE directed compression. Axes plunge 10 – 30° NE and SW while the scattering of at least 60° is due to D_5 deformation phase.

At the microscale, C'-type shear bands developed under low-grade metamorphic conditions. The D_3 event deforms the tectonic boundary between LTE and UTE and all other previous structures (Fig. 8, cross-sections B–B', C–C').

4.5. D_4 Extensional Crenulation Cleavage (ECC)

Extensional Crenulation Cleavage (as defined by Platt and Vissers, 1980) is well developed in calcschists and locally in serpentinites at the mesoscale (Fig. 4g). It is emphasised by centimetric to millimetric spaced surfaces which mostly occur along a NE–SW belt less than one kilometre thick within the calcschists and subordinately in the Sesia–Lanzo gneisses of the UTE. S_4 surfaces dip mainly to SE at 60° to sub-vertical with down-dip lineations defined by quartz and calcite fibres and by striations indicating top to SE normal shear. Axes plunge less than 30° NE and SW (Fig. 3i). Locally at the contact between calcschists and gneisses, ECC steps gradually into frictional structures in the latter lithologies, suggesting that this deformation event took place under the brittle–ductile

transition regime. ECC shows a significant grain-size reduction (Fig. 5e,g,h) and is defined by low-grade metamorphic conditions.

Field observations at the western border of the Gran Paradiso Unit, on the watershed separating the Orco Valley from the Isère Valley, confirm the presence of ECC dipping to NW with top to NW normal shear.

4.6. D_5 structures and frictional deformation regime

The D_5 event is only detected at the macroscale by analysing variation in orientation of previous structures. It is responsible for undulations of S_1 with 10–200 m wavelengths. Axial planes strike between 100 – 160° and dip both NE and SW with high angle, while fold axes scatter from 080° to 150° with variable inclination. The scattering of geometric features indicate the presence of conjugate and anastomosing axial surfaces that coincide with a conjugate high angle fracture system striking 120 – 150° .

5. Discussion

5.1. Coupling of the Gran Paradiso versus the Zermatt–Saas

In the Gran Paradiso Unit the development of the S_1 at the expense of garnet + omphacite + glaucophane assemblages in metabasites indicates that the main foliation formed during exhumation under medium PT conditions (Table 1). An eclogitic-peak pre- D_1 event in orthogneisses was recorded only along thin shear bands where whiteschists developed, while surrounding rocks remained in metastable conditions probably because of H_2O deficiency that prevents metamorphic reactions and recrystallization into denser rocks (Austrheim et al., 1997; Proyer, 2003). On the basis of bibliographic data, a maximum age of 43 ± 0.5 Ma (Meffan-Main et al., 2004) at 21–23 kbar and 540 – 570°C (Wei and Powell, 2003, 2004) can be assigned to the eclogite facies pre- D_1 event for the Gran Paradiso Unit.

In the Zermatt–Saas-like Unit an eclogitic pre- S_1 foliation is locally well preserved while metagabbros and most metabasites are extensively re-equilibrated during the D_1 event, which is characterised by medium PT conditions.

The tectonic contact between the Gran Paradiso and the Zermatt–Saas-like units is deformed by the D_1 event (Figs. 8, 9 and 10a,b) and is defined by blueschist assemblages (see Section 3.5). This tectonic contact is deformed by a macro-scale fold showing an anvil-shaped section (Mies, 1993) as reported in the structural map of Fig. 7. This geometry, linked to the parallelism between L_1 and A_1 (Fig. 3a–c), is referred to the development of sheath folds whose structure is sketched in Fig. 9. The structural evidence that eclogite facies pre- S_1 foliation of the Zermatt–Saas metabasites is truncated by the tectonic contact and is folded by the D_1 event (Fig. 6) implies that the coupling between the Gran Paradiso and the Zermatt–Saas-like




Fig. 7. Simplified structural map of the study area with some structural features of the D_1 event. Only the different palaeo-geographic units are distinguished to show their structural relationship (geochronological data from Inger et al., 1996; Inger and Ramsbotham, 1997; Meffan-Main et al., 2004).

units developed after the pre- S_1 eclogite facies event but before the D_1 event, suggesting nappe emplacement during exhumation.

The D_1 event developed during a top to W compressional regime (Figs. 4c and 8) and could be related to the top to NW shearing event of Inger and Ramsbotham (1997), dated at 38–40 Ma, within Gran Paradiso Unit W of Locana. On the base of structural and petrographic features, the main foliation S_1 of the Gran Paradiso Unit can be related to the regional fabric described by Brouwer et al. (2002) west of the study area, for which they suggest metamorphic conditions of 10–12 kbar and 500–550 °C.

5.2. Coupling of the Sesia–Lanzo GMC and the Combin Unit

Both the greenschist facies units were pervasively deformed during D_1 event that developed the main regional foliation under greenschist facies conditions (Figs. 7 and 10b). The presence of rare pre- D_1 , zoisite–rutile-bearing assemblages, in little deformed lithons within Sesia–Lanzo orthogneisses indicates an older metamorphic event, but there is no evidence of eclogitic or blueschist facies assemblages. Inger et al. (1996) reported an age of 37.5 ± 1 Ma for the contact between the Gneiss Minuti Complex




Fig. 8. Cross-sections representing the structural relationships between the different tectonic units (see text for explanation). GP, Gran Paradiso Unit; ZS, Zermatt–Saas Unit; CO, Combin Unit; SL, Sesia–Lanzo Gneiss Minuti Complex; OSZ, Orco Shear Zone. Legend as in Fig. 7.

and the Combin-like calcschists in Soana Valley north of the study area.

5.3. Coupling of LTE vs. UTE: The Orco Shear Zone

Constraining the timing of the tectonic contact between Upper and Lower Tectonic Element (Orco Shear Zone) is not simple. The geometric and structural relationships show that this tectonic contact cuts D₁ structures both in LTE (HP units) and UTE (LP units) and is folded by the D₃ event (see Figs. 2, 7 and 8). The evidence of a top to E sense of shear (Fig. 4h) in the mylonitic foliation along the contact between LTE and UTE is in agreement with D₂ tectonic transport. D₂ structures are slightly different in LTE and UTE: the main axial plane surface differs by at least 25° (Fig. 3g,h) and moreover D₂ structures are better developed in the UTE, suggesting that the contact can be related to the late-D₂ event (Fig. 10c). Therefore, the Orco Shear Zone represents a metamorphic gap between the two Tectonic Elements accommodating exhumation of the Lower one, and stopped working before the onset of the D₃ event (see Figs. 2, 7 and 8). This regional scale structure is responsible for the absence of Zermatt–Saas-like oceanic units towards the N, in Soana Valley (see figure 8 of Inger and Ramsbotham, 1997) where GP orthogneisses are directly in contact with the calcschists of the Combin-like Unit. The absence of the oceanic crust can be attributed to a delamination process during exhumation along this tectonic contact, for which Inger and Ramsbotham (1997) suggest an age of 34–39 Ma.

In summary the Orco Shear Zone shows the following features: (i) it consists of a late ductile tectonic contact developed under




Fig. 9. Three dimension sketch of the structural relationships between the Gran Paradiso and the Zermatt–Saas Units showing a D₁ non-cylindrical synform-antiform pair folding their tectonic contact. Axial plane surface (S₁) and mean direction of stretching lineation (L₁) are indicated. The structure outcrops north of Schiaroglio (see Fig. 7). Sketch not to scale.




Fig. 10. W–E cross-sections representing the structural history of the Orco Valley geological transect related to its possible geodynamic evolution. OSZ, Orco Shear Zone; EMC-SL, Eclogitic Micaschists Complex, Sesia–Lanzo; GMC-SL, Gneiss Minuti Complex, Sesia–Lanzo; CO, Combin Unit; ZS, Zermatt–Saas Unit; GP, Gran Paradiso Unit. LTE, Lower Tectonic Element; UTE, Upper Tectonic Element. (a) pre-D₁ stage and HP metamorphism in the LTE. The position of the OSZ is indicated and probably accommodated reverse-slip movement. (b) D₁ stage represents the main folding event resulting in a well-developed regional foliation in both the UTE and the LTE. (c) D₂ stage is linked to minor folding (not shown) and to the exhumation of the LTE along the OSZ that cuts D₁ structures. (d) D₃ stage folded the OSZ and caused the doming of the Gran Paradiso Unit. (e) D₄ stage is associated to the Extensional Crenulation Cleavage (ECC) on both side of the LTE and probably represents the onset of gravitational collapse.

greenschist facies conditions; (ii) it juxtaposes eclogite-bearing units to greenschist ones; (iii) at present, it dips towards ESE and shows top to E normal sense of shear. In the Western Alps, similar regional scale shear zones have been reported by some authors

(Combin Fault: Ballèvre and Merle, 1993; Gressoney Shear Zone; Reddy et al., 1999). Both the Combin Fault and the Gressoney Shear Zone represent the tectonic contact between HP and overlying LP units outcropping in the northern area of the Aosta Valley. Since the Orco Shear Zone occurs in the same structural position (HP–LP boundary), it can be considered the southern continuation of these structures. According to Ballèvre and Merle (1993), the Combin Fault is mainly an Early Tertiary detachment fault reactivated as thrust during Oligocene. Ring (1995) interpreted the Combin Fault as an out-of-sequence NW directed thrust (Late Eocene) subsequently reactivated as SE directed backthrust (Early Oligocene). Reddy et al. (1999, 2003) suggest for the Gressoney Shear Zone an early compressional history followed by pervasive SE extension between 45 and 36 Ma. Therefore the interpretation of these shear zones is very ambiguous. On the basis of our structural and metamorphic data we suggest for the Orco Shear Zone a main top to SE extensional component at a shallow crustal level, even if an older top to NW compressional displacement (before D₂) cannot be excluded.

5.4. Late structural and metamorphic evolution

After the D₂ event a metamorphic stage under static conditions was recorded in all the studied units. It is characterized by albite + biotite in Gran Paradiso paragneisses and Sesia–Lanzo gneisses and by albite + green amphibole + epidote ± biotite in Gran Paradiso and Zermatt–Saas metabasites.

Afterward the D₃ event developed and locally caused activation of reverse-slip C′-type shear bands only in less competent lithologies such as calcschists, serpentinites and Sesia-gneisses. Microstructural observations indicate recrystallization coupled with grain-size reduction, suggesting high strain deformation in low-grade metamorphic conditions. The D₃ event deformed the contact between the LTE and the UTE (Figs. 7 and 10d) and may be responsible for the doming of the Gran Paradiso.

The following ECC caused a further differentiated exhumation between the Lower Tectonic Element (the footwall) and the Upper one (the hangingwall) (Fig. 10e). ECC developed in ductile conditions in phyllosilicate-rich rocks while in quartz-feldspathic rocks ECC is defined by fractures and faults indicating a plastic-frictional transition regime in agreement with deformation at a shallow crustal level. The presence of ECC at the western border of the Gran Paradiso Unit (this work; Rolland et al., 2000; Bucher et al., 2003) can be related to an axial culmination of the Alpine mountain belt in this area, corresponding to the outcrop of the deepest eclogitic units of the Western Alps. D₄ probably represents the onset of the gravitational collapse of the Alpine-orogen linked to further exhumation of the Lower Tectonic Element.

6. Conclusions

This paper provides new structural and petrographic data in the Orco Valley and allows a better constraint of the tectonic framework within this portion of the Alpine orogen. It also gives more detailed information about the tectono-metamorphic evolution of a key-area in the Western Italian Alps. Two main Tectonic Elements have been distinguished (Fig. 7): the Lower one composed of eclogite facies units (Gran Paradiso Unit and Zermatt–Saas-like oceanic units) and the Upper one made of greenschist facies units (the Combin-like Unit and the Gneiss Minuti Complex of the Sesia–Lanzo Zone). In both Tectonic Elements, the units were independently coupled before the occurrence of the regional schistosity (S₁) which developed under different metamorphic conditions and probably at different times across the two Tectonic Elements.

In the Lower Tectonic Element (LTE), Gran Paradiso and Zermatt–Saas-like units were coupled during exhumation after pre-D₁

eclogite facies conditions and then were folded together by the D₁ event under medium PT conditions (upper-greenschist to epidote–albite amphibolite facies). Pre-D₁ structural and mineralogical HP relicts are still present at every scale, particularly in the Zermatt–Saas-like Unit where mesoscale eclogitic foliations and macroscale structures are preserved. Additionally this unit consists of different tectonic slices in agreement with the structural setting proposed by Battiston et al. (1984) towards N in Val Soana. In the Gran Paradiso orthogneisses eclogite facies assemblages occur only along discrete shear zones where whiteschists developed.

In the Upper Tectonic Element (UTE), structural and mineralogical evidence of a pre-D₁ evolution is also preserved. However, in these units both the pre-D₁ metamorphic assemblages and the regional foliation (D₁) developed under greenschist facies conditions.

The LTE and the UTE are separated by a normal-slip tectonic contact represented by a thin mylonitic top to E shear zone that occurred during the late-D₂ event. This first order structure (Orco Shear Zone) led to the exhumation of the HP units (LTE) and coupled them with LP ones (UTE). This contact defines the boundary between the Zermatt–Saas-like Unit and the Combin-like Unit; therefore it can be considered as the continuation of the Gressoney Shear Zone (Reddy et al., 1999) and of the Combin Fault (Ballèvre and Merle, 1993) further S of the Aosta Valley.

From D₃ onward all units followed the same tectono-metamorphic history and the dome structure of the Gran Paradiso took its final shape. Finally the D₄ event (ECC) occurred along a thin belt mainly within Combin-like calcschists, implying a further differential exhumation of LTE (footwall) with respect to UTE (hangingwall) at the brittle-plastic transition regime.

The occurrence of a W-dipping normal-slip shear zone between Zermatt–Saas-like Unit (footwall) and Combin-like Unit (hangingwall) at the western border of the Gran Paradiso Unit (Rolland et al., 2000), corresponding to our late-D₂ Orco Shear Zone, suggests the presence of a core complex made of eclogite-bearing units beneath greenschist facies ones. This core complex occurs in the axial position within the Western Alps orogenic wedge and can be responsible for the late exhumation stage of the deeper subducted units.

Acknowledgements

This work was financially supported by Ministero dell'Università e della Ricerca Scientifica e Tecnologica (M.U.R.S.T.). J. Platt and two anonymous reviewers are thanked for providing substantial improvement of the manuscript.

References

- Amato, J.M., Johnson, C.M., Baumgartner, L.P., Beard, B.L., 1999. Rapid exhumation of the Zermatt–Saas ophiolite deduced from high-precision Sm–Nd and Rb–Sr geochronology. *Earth and Planetary Science Letters* 171, 425–438.
- Austrheim, H., Erambert, M., Engvik, A., 1997. Processing of crust in the root of the Caledonian continental collision zone: the role of eclogitization. *Tectonophysics* 273, 129–153.
- Ballèvre, M., 1988. Collision continentale et chemins P–T: L'unité pennique du Grand Paradis (Alpes occidentales). *Mémoires et Documents du Centre Armoricain d'Etudes Structurales des Socles* 19, 352.
- Ballèvre, M., Merle, O., 1993. The Combin fault: Compressional reactivation of a Late Cretaceous–Early Tertiary detachment fault in the western Alps. *Schweizerische Mineralogische und Petrographische Mitteilungen* 73, 205–227.
- Barnicoat, A.C., Rex, D.C., Guise, P.G., Cliff, R.A., 1995. The timing of and nature of greenschist facies deformation and metamorphism in the upper pennine Alps. *Tectonics* 14 (2), 279–293.
- Battiston, P., Benciolini, L., Dal Piaz, G.V., De Vecchi, G., Marchi, G., Martin, S., Polino, R., Tartarotti, P., 1984. Geologia di una traversa dal Gran Paradiso alla Zona Sesia–Lanzo in alta Valle Soana, Piemonte. *Memorie della Società Geologica Italiana* 29, 209–232.
- Bearth, P., 1967. Die Ophiolithe der Zone von Zermatt–Saas Fee. *Beiträge Geologische Karte Schweiz*, Neue Folge 132, 1–130.
- Bertrand, J.M., Pidgeon, R.T., Leterrier, J., Guillot, F., Gasquet, D., Gattiglio, M., 2000. SHRIMP and IDTIMS U–Pb zircon ages of the pre-Alpine basement in the internal Western Alps (Savoy and Piedmont). *Schweizerische Mineralogische und Petrographische Mitteilungen* 80, 222–248.
- Biino, G., Pognante, U., 1989. Palaeozoic continental-type gabbros in the Gran Paradiso nappe (Western Alps, Italy): Early-Alpine eclogitization and geochemistry. *Lithos* 24, 3–19.
- Borghi, A., Compagnoni, R., Sandrone, R., 1996. Composite P–T paths in the internal Penninic massifs of the Western Alps: petrological constraints to their thermomechanical evolution. *Eclogae Geologicae Helveticae* 89, 345–367.
- Brouwer, F.M., Vissers, R.L.M., Lamb, W.M., 2002. Structure and metamorphism of the Gran Paradiso massif, Western Alps, Italy. *Contributions to Mineralogy and Petrology* 143, 450–470.
- Brouwer, F.M., Van De Zedde, D.M.A., Wortel, M.J.R., Vissers, R.L.M., 2004. Late-orogenic heating during exhumation: Alpine PT trajectories and thermomechanical models. *Earth and Planetary Science Letters* 220, 185–199.
- Bucher, S., Schmid, S.M., Bousquet, R., Fügenschuh, B., 2003. Late stage deformation in a collisional orogen (Western Alps): nappe refolding, back thrusting or normal faulting? *Terra Nova* 15, 109–117.
- Bucher, S., Ulardic, C., Bousquet, R., Ceriani, S., Fügenschuh, B., Gouffon, Y., Schmid, S.M., 2004. Tectonic evolution of the Briançonnais units along a transect (ECORS-CROP) through the Italian–French Western Alps. *Eclogae Geologicae Helveticae* 97, 321–345.
- Callegari, E., Compagnoni, R., Dal Piaz, G.V., 1969. Relitti di strutture intrusive erciniche e scisti a sillimanite nel Massiccio del Gran Paradiso. *Bollettino della Società Geologica Italiana* 88, 59–69.
- Cartwright, I., Barnicoat, A.C., 2002. Petrology, geochronology, and tectonics of shear zones in the Zermatt–Saas and Combin Zones of the Western Alps. *Journal of Metamorphic Geology* 20, 263–281.
- Compagnoni, R., Lombardo, B., 1974. The Alpine age of the Gran Paradiso eclogites. *Rendiconti della Società Geologica Italiana di Mineralogia e Petrografia* 30, 223–237.
- Compagnoni, R., Prato, R., 1969. Paramorfosi di cianite su sillimanite in scisti pregranitici del Massiccio del Gran Paradiso. *Bollettino della Società Geologica Italiana* 88, 547–549.
- Compagnoni, R., Elter, G., Lombardo, B., 1974. Eterogeneità stratigrafica del complesso degli “Gneiss Minuti” nel massiccio cristallino del Gran Paradiso. *Memorie della Società Geologica Italiana* 13, 227–239.
- Compagnoni, R., Dal Piaz, G.V., Hunziker, J.C., Lombardo, B., Williams, P.F., 1977. The Sesia–Lanzo Zone, a slice of continental crust with Alpine high pressure–low temperature assemblages in the Western Italian Alps. *Rendiconti della Società Geologica Italiana di Mineralogia e Petrografia* 33 (1), 281–334.
- Dal Piaz, G.V., 1974. Le métamorphisme alpin de haute pression et basse température dans l'évolution structurale du bassin ophiolitique alpine-appenninique. 1^e Partie: *Bollettino della Società Geologica Italiana*, 93, 437–468, 2^e Partie: *Schweizerische Mineralogische und Petrographische Mitteilungen*, 54, 399–424.
- Dal Piaz, G.V., 1999. The AustroAlpine–Piedmont nappe stack and the puzzle of Alpine Tethys. *Memorie della Società Geologica Italiana* 51 (1), 155–176.
- Dal Piaz, G.V., Ernst, W.G., 1978. Areal geology and petrography of eclogites and associated metabasites of the Piedmont ophiolite nappe, Breuil–St. Jacques area, Italian Western Alps. *Tectonophysics* 51, 99–126.
- Dal Piaz, G.V., Cortiana, G., Del Moro, A., Martin, S., Pennacchioni, G., Tartarotti, P., 2001. Tertiary age and paleostructural inferences of the eclogitic imprint in the Austroalpine outliers and Zermatt–Saas ophiolite, Western Alps. *Journal of Earth Sciences* 90, 668–684.
- Desmons, J., Compagnoni, R., Cortesogno, L., with the collaboration of Frey, M., Gaggero, L., Della Giovanna, G., Seno, S., Radelli, L., 1999a. Alpine metamorphism of the Western Alps: I. Middle to high P/T metamorphism. *Schweizerische Mineralogische und Petrographische Mitteilungen* 79, 89–110.
- Desmons, J., Compagnoni, R., Cortesogno, L., with the collaboration of Frey, M., Gaggero, L., Della Giovanna, G., Seno, S., Radelli, L., 1999b. Alpine metamorphism of the Western Alps: II. Middle to high P/T metamorphism. *Schweizerische Mineralogische und Petrographische Mitteilungen* 79, 111–134.
- Duchêne, S., Lardeaux, J.M., Albarède, F., 1997. Exhumation of eclogites: insights from depth–time path analysis. *Tectonophysics* 280, 125–140.
- Freeman, S.R., Inger, S., Butler, R.W.H., Cliff, R.A., 1997. Dating deformation using Rb–Sr in white mica: Greenschist facies deformation ages from the Entrelor shear zone, Italian Alps. *Tectonics* 16 (1), 57–76.
- Hurford, A.J., Hunziker, J.C., 1989. A revised thermal history for the Gran Paradiso massif. *Schweizerische Mineralogische und Petrographische Mitteilungen* 69, 319–329.
- Hunziker, J.C., Desmons, J., Hurford, A.J., 1993. Thirty-two years of geochronological work in the Central and Western Alps: a review on seven maps. *Mémoires de Géologie*, Lausanne, 13, 52 pp.
- Inger, S., Ramsbotham, W., 1997. Syn-convergent exhumation implied by progressive deformation and metamorphism in the Valle dell'Orco transect, NW Italian Alps. *Journal of the Geological Society of London* 154, 667–677.
- Inger, S., Ramsbotham, W., Cliff, R.A., Rex, D.C., 1996. Metamorphic evolution of the Sesia–Lanzo Zone, Western Alps: time constraints from multi-system geochronology. *Contributions to Mineralogy and Petrology* 126, 152–168.
- Kassem, O.K., Ring, U., 2004. Underplating-related finite-strain patterns in the Gran Paradiso massif, Italian Western Alps: Heterogeneous ductile strain superimposed on a nappe stack. *Journal of the Geological Society*, London 161, 875–884.

- Kretz, R., 1983. Symbols for rock-forming minerals. *American Mineralogist* 68, 277–279.
- Lapen, T.J., Johnson, C.M., Baumgartner, L.P., Dal Piaz, G.V., Skore, S., Beard, B., 2007. Coupling of oceanic and continental crust during Eocene eclogite-facies metamorphism: evidence from the Monte Rosa nappe, western Alps. *Contributions to Mineralogy and Petrology* 153, 139–157.
- Lardeaux, J.M., Spalla, M.I., 1991. From granulites to eclogites in the Sesia Zone (Italian Western Alps): a record of the opening and closure of the Piedmont ocean. *Journal of Metamorphic Geology* 9, 35–59.
- Lardeaux, J.M., Gosso, G., Kienast, J.R., Lombardo, B., 1982. Relations entre le métamorphisme et la déformation dans la Zone de Sesia–Lanzo (Alpes occidentales) et le problème de l'éclogitisation de la croûte continentale. *Bulletin de la Société Géologique de France* 24 (7), 793–800.
- Le Bayon, B., Pitre, P., Ballèvre, M., Bohn, M., 2006. Reconstructing P-T Paths during continental collision using multi-stage garnet (Gran Paradiso nappe, Western Alps). *Journal of Metamorphic Geology* 24, 477–496.
- Meffan-Main, S., Cliff, R.A., Barnicoat, A.C., Lombardo, B., Compagnoni, R., 2004. A Tertiary age for Alpine high-pressure metamorphism in the Gran Paradiso massif, Western Alps: a Rb–Sr microsampling study. *Journal of Metamorphic Geology* 22, 267–281.
- Messiga, B., Kienast, J.R., Rebay, G., Piccardi, M.P., Tribuzio, R., 1999. Cr-rich Magnesio-Chloritid from the Monviso ophiolites (Western Alps, Italy). *Journal of Metamorphic Geology* 17, 287–299.
- Mies, J.W., 1993. Structural analysis of shear folds in the Sylacauga Marble Group, Talcadegate slate belt, southern Appalachians. *Journal of Structural Geology* 15, 983–993.
- Platt, J.P., 1993. Exhumation of high-pressure rocks: a review of concept and processes. *Terra Nova* 5, 119–133.
- Platt, J.P., Vissers, R.L.M., 1980. Extensional structures in anisotropic rocks. *Journal of Structural Geology* 2, 397–410.
- Pleuger, J., Froitzheim, N., Jansen, E., 2005. Folded continental and oceanic nappes on the southern side of Monte Rosa (western Alps, Italy): Anatomy of a double collision suture. *Tectonics*, 24, TC4013, doi:10.1029/2004TC001737.
- Pognante, U., 1989. Lawsonite, blueschist and eclogite formation in the Southern Sesia Zone Western Alps, Italy. *European Journal of Mineralogy* 1, 89–104.
- Pognante, U., Compagnoni, R., Gosso, G., 1980. Micro-mesostructural relationships in the continental eclogitic rocks of the Sesia–Lanzo Zone (Italian Western Alps): a record of subduction cycle. *Rendiconti della Società Italiana di Mineralogia e Petrografia* 36, 169–186.
- Pognante, U., Talarico, F., Rastelli, N., Ferrati, N., 1987. High pressure metamorphism in the nappes of the Valle dell'Orco traverse (Western Alps collisional belt). *Journal of Metamorphic Geology* 5, 397–414.
- Proyer, A., 2003. The preservation of high-pressure rocks during exhumation: metagranites and metapelites. *Lithos* 70, 183–194.
- Reddy, S.M., Wheeler, J., Cliff, R.A., 1999. The geometry and timing of orogenic extension: an example from the Western Italian Alps. *Journal of Metamorphic Geology* 17, 573–589.
- Reddy, S.M., Wheeler, J., Butler, R.W.H., Cliff, R.A., Freeman, S., Inger, S., Pickels, C., Kelley, S.P., 2003. Kinematic reworking and exhumation within the convergent Alpine Orogen. *Tectonophysics* 365, 77–102.
- Reinecke, T., 1991. Very-high-pressure metamorphism and uplift of coesite bearing metasediments from the Zermatt–Saas Zone, Western Alps. *European Journal of Mineralogy* 3, 7–17.
- Reinecke, T., 1998. Prograde high- to ultrahigh-pressure metamorphism and exhumation of oceanic sediments at Lago di Cignana, Zermatt–Saas Zone, Western Alps. *Lithos* 42, 147–189.
- Ring, U., 1995. Horizontal contraction or horizontal extension? Heterogeneous Late Eocene and Early Oligocene general shearing during blueschist and greenschist facies metamorphism at the Pennine–Austroalpine boundary zone in the Western Alps. *Geologische Rundschau* 84, 843–859.
- Ring, U., Collins, A.S., Kassem, O.K., 2005. U–Pb SHRIMP data on the crystallization age of the Gran Paradiso augengneiss, Italian Western Alps: Further evidence for Permian magmatic activity in the Alps during break-up of Pangea. *Eclogae Geologicae Helveticae* 98, 363–370.
- Rolland, Y., Lardeaux, J.M., Guillot, S., Nicollet, C., 2000. Extension syn-convergence, poinçonnement vertical et unités métamorphiques contrastées en bordure ouest du Gran Paradis (Alpes Franco-Italiennes). *Geodynamica Acta* 13, 133–148.
- Rubatto, D., Gebauer, D., Fanning, M., 1998. Jurassic formation and Eocene subduction of the Zermatt–Saas-Fee ophiolites: implications for the geodynamic evolution of the Central and Western Alps. *Contributions to Mineralogy and Petrology* 132, 269–287.
- Rubatto, D., Gebauer, D., Compagnoni, R., 1999. Dating of eclogite-facies zircon: the age of Alpine metamorphism in the Sesia–Lanzo Zone (Western Alps). *Earth and Planetary Science Letters* 167, 141–158.
- Schmid, S.M., Fügenschuh, B., Kissling, E., Schuster, R., 2004. Tectonic map and overall architecture of the Alpine orogen. *Eclogae Geologicae Helveticae* 97, 93–117.
- Spalla, M.I., Lardeaux, J.M., Dal Piaz, G.V., Gosso, G., Messiga, B., 1996. Tectonic significance of Alpine eclogites. *Journal of Geodynamics* 21, 257–285.
- Wei, C., Powell, R., 2003. Phase relations in high-pressure metapelites in the system KFMASH (K₂O–FeO–MgO–Al₂O₃–SiO₂–H₂O) with application to natural rocks. *Contributions to Mineralogy and Petrology* 145, 301–315.
- Wei, C., Powell, R., 2004. Calculated phase relations in high-pressure metapelites in the system NKFMAH (Na₂O–K₂O–FeO–MgO–Al₂O₃–SiO₂–H₂O). *Journal of Petrology* 45, 183–202.
- Williams, P.F., Compagnoni, R., 1983. Deformation and metamorphism in the Bard area of the Sesia–Lanzo Zone, Western Alps, during subduction and uplift. *Journal of Metamorphic Geology* 1, 117–140.