

EQUILIBRIUM STUDY BETWEEN CONDENSED PHASES BY ISOPLETHIC THERMAL ANALYSIS WHEN A MISCELLIBILITY GAP IS OBSERVED

E. Labarthe*, A. J. Bougrine, H. Delalu, J. Berthet and J. J. Couniou

Université de Lyon, Université Lyon 1, Laboratoire Hydrazines et Procédés – UMR CNRS 5179, F-69622 Villeurbanne Cedex

Isoplethic thermal analysis was used to determine the solid-liquid-liquid equilibria in the ternary system water-sodium sulfate-piperidine. The changes in state observed on the thermogram recorded during the displacement in a quasi-binary section permit the identification of the different phases and the delimitation of the corresponding equilibrium domains. Two isotherms were established at 25°C and 40°C because these temperatures frame the peritectic decomposition of the sodium sulfate dehydrate. Miscibility gaps were completely delimited and each critical point was calculated. This study permitted to determine the optimal conditions for separating the organic phase by liquid-liquid extraction.

Keywords: critical point, isoplethic thermal analysis, miscibility gap, piperidine, sodium sulfate, ternary diagram, water

Introduction

Studying a high order system requires long and painstaking experimentations when using traditional analysis techniques. Therefore, the laboratory has developed automated isoplethic methods that consist in progressively modifying the global composition of the system, monitoring the evolution of an intensive variable and using the information obtained. The first works [1], which were based on successive and then continuous conductimetric measurements, demonstrated how it was possible to determine all the characteristics of an isotherm: solubility curves, invariant points, nature and composition of solid phases.

Direct or differential thermal analysis rests on the exploitation of the thermal effects that occur when a sample is subjected to a regular variation of its heat energy. The progressive addition of a given solution in a mixture modifies its global composition. The molar and mass compositions of each constituent vary linearly through time and thus define an isoplethic section of the system formed by all the independent constituents. Isoplethic Thermal Analysis [2, 3], or ITA, consists in using the thermal phenomena observed under almost constant pressure and temperature conditions.

Experimental

In an invariant isotherm-isobar system, the number of phases in equilibrium is equal to the number of independent constituents c . If an infinitesimal quantity da

(molar or mass unit) of a mixture A , with composition vector \vec{A} , is added to an invariant system composed of c phases j of composition vector \vec{F}_j in quantity f_j , only the quantities df_j of these phases will be modified. According to the properties of the isobarycentric transformations [4], these thermodynamic variables satisfy the relation (1):

$$da \vec{A} = \sum_{j=1}^{i=c} df_j \vec{F}_j \quad (1)$$

which can also be written as:

$$\sum_{j=1}^{i=c} df_j \vec{A} \vec{F}_j = \vec{0} \quad (2)$$

since

$$\sum_{j=1}^{i=c} df_j = da \quad (3)$$

The resolution of this linear equation system permits the calculation of each df_j for a given value of da . If H_A is the enthalpy of the mixture added and H_j is the enthalpy of the phase j , the enthalpy difference (molar or in mass) of the system is given by:

$$dH = \sum_{j=1}^{i=c} df_j H_j - da H_A \quad (4)$$

In the case of an invariant system, dH depends only on the added quantity da . In particular, this variation of enthalpy is not a function of the mass and

* Author for correspondence: emilie.labarthe@univ-lyon1.fr

composition of the initial system but it is directly proportional to da .

If the mixture A is introduced at a constant flow rate, $q=da/dt$, the heat quantity involved per time unit is constant:

$$dH/dt=k_1q \quad (5)$$

In addition, if the measurement cell is of isoperibolic type, the heat flow Φ , to which the system is subjected, is proportional to the variation of temperature, according to Newton's law:

$$\Phi=\frac{\delta Q}{dt}=k_2\Delta T \quad (6)$$

where ΔT is the temperature difference between the mixture temperature T_M and the thermostatic chamber temperature T_C .

$$\Delta T=T_M-T_C \quad (7)$$

Hence the relation:

$$\Delta T=(k_1/k_2)q=Kq \quad (8)$$

Finally, for an invariant isotherm-isobar system, a given mixture and injection speed, the variation of temperature between the mixture and the thermostatic chamber is constant and a temperature plateau should be observed. The displacement of an indifferent equilibrium is therefore expressed by ITA as a plateau of invariance, as in direct thermal analysis.

In practice, the variation of the temperature ΔT is recorded as a function of the volume v of solution added. Each equilibrium domain has a corresponding specific variation law, allowing the determination of phase transformations.

An experimental set-up was built [5] according to the principle defined previously and adjusted in particular to a temperature measurement sensitivity in the region of mK. This set-up is completely automated and is composed of a platinum probe Pt100, a thermostat Hetofrig type CB11, a reactor, an automatic injection system and a magnetic agitator.

The injection of the solvent in the stirred mixture is performed by a computer controlled automatic injection




Fig. 1 Example of thermogram registered during addition of pure water in the ternary mixture A

system. The injection range used is within 1 and 4 mL h⁻¹. The temperature values are stored at regular time intervals. The phase transformations are determined by computer processing of the numerical data by linear extrapolation of the variation laws.

This device was applied to the study of equilibria between the solid-liquid phases and liquid-liquid phases of the ternary system water-sodium sulfate-piperidine, used as the basis for extracting organic compounds in a complex synthetic mixture.

The ternary system studied is delimited by three binary systems. The Na₂SO₄-C₅H₁₀NH system is unknown, while in the temperature field studied the piperidine is totally miscible in water. On the other hand, two hydrates are present in the binary mixture H₂O-Na₂SO₄ [6, 7]: a metastable heptahydrate and a stable decahydrate (Glauber salt) which undergoes a peritectic decomposition at 32.4°C.

In order to analyse the impact of the decahydrate on the equilibria, we established two isotherms at 25 and 40°C, that frame its decomposition temperature.

Figure 2 gives an example of a thermogram obtained from a ternary mixture whose composition point A is located in Fig. 3. It shows the evolution of the temperature mixture when a volume of pure water is added.

Segment AB corresponds to the progressive hydration of anhydrous sodium sulfate into dehydrated salt. Reaching steady state was terminated with a short plateau of invariance. From B to C, crossing the two phases equilibrium domain liquid+Na₂SO₄·10H₂O is monotonous. After reaching steady state C at C', crossing the triphasic domain L1+L2+Na₂SO₄·10H₂O is shown by a plateau of invariance up to point D. Point D, beyond which no solid phase can be observed, corresponds to the limit conode of the miscibility gap. By using this technique, the different equilibrium domains are clearly defined and permit the full determination of the isotherm.




Fig. 2 Ternary system H₂O-Na₂SO₄-C₅H₁₀NH : isotherm 25°C (mass fractions). Solid phase I: Na₂SO₄, Solid phase II: Na₂SO₄·10H₂O




Fig. 3 Ternary system $\text{H}_2\text{O}-\text{Na}_2\text{SO}_4-\text{C}_5\text{H}_{10}\text{NH}$: isotherm 40°C (mass fractions), Solid phase I: Na_2SO_4 , Solid phase II: $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$

Results and discussion

Figures 2 and 3 represent isotherms 25 and 40°C respectively of the ternary system $\text{H}_2\text{O}-\text{Na}_2\text{SO}_4-\text{C}_5\text{H}_{10}\text{NH}$ while Tables 1 and 2 give the associated experimental values, expressed in mass fractions.

The 25°C isotherm is characterized by the crystallization of two solid phases, the anhydrous and the sodium sulfate decahydrate, and the existence of a miscibility gap in the crystallization area of the decahydrate. The liquid-liquid equilibrium is delimited by the conode L1-L2 and the critical point κ . The solubility of the anhydrous sulfate in concentrated aqueous solutions of piperidine is very weak and the liquidus curve is almost confounded with the limit binary system $\text{H}_2\text{O}-\text{C}_5\text{H}_{10}\text{NH}$.

At 40°C, the decahydrate is no more stable and the liquidus curve of the anhydrous salt is interrupted

Table 1 Isotherm 25°C determined by ATI

w($\text{C}_5\text{H}_{10}\text{NH}$)	w(Na_2SO_4)	Phenomenon
0	0.207	Liquidus of $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$
0.0144	0.2011	"
0.0272	0.1811	"
0.0345	0.1731	"
0.045	0.170	Invariant point L1*
0.0350	0.2333	Limit conode L1- $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$
0.054	0.151	binodal
0.0680	0.136	"
0.0995	0.1161	"
0.1304	0.1012	"
0.16	0.09	Critical point κ^{**}
0.198	0.073	binodal
0.3085	0.0383	"
0.336	0.036	Limit conode L1-L2
0.0770	0.1539	"
0.0956	0.1439	"
0.1214	0.1417	"
0.1602	0.1243	"
0.1830	0.1161	"
0.2273	0.0835	" (point D)
0.3160	0.0551	"
0.3267	0.0406	Invariant point L2*
0.385	0.026	Limit conode L2- $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ (point C)
0.3438	0.0708	Liquidus of $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$
0.495	0.0243	"
0.60	0.0039	Limit conode L3- $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$
0.5694	0.0711	"(point B)
0.5152	0.1061	Invariant point L3*
0.67	<0.001	

chemical analysis(*) or generalized diameter method (**)

Table 2 Isotherm 40°C determined by ATI

$w(C_5H_{10}NH)$	$w(Na_2SO_4)$	Phenomenon
0.0009	0.314	Invariant point L1*
0.008	0.272	binodal
0.008	0.259	"
0.022	0.182	"
0.039	0.142	"
0.078	0.118	"
0.12	0.10	Critical point κ^{**}
0.137	0.092	binodal
0.291	0.041	"
0.375	0.027	"
0.465	0.011	"
0.563	0.006	Limit conode L1-L2
0.04	0.30	"
0.272	0.181	"
0.474	0.067	"
0.61	0.014	Invariant point L2*
0.628	0.002	

chemical analysis (*) or generalized diameter method (**)

by the demixing curve. One of the parts of the liquidus is almost confounded with the point L1.

The liquid-liquid area is strongly dependent on temperature. The critical line, that separates the two half surfaces of the binodal, is a fundamental characteristic of the miscibility gap. However, even at a given temperature, its composition is difficult to define experimentally. Ideally, when the two branches of the binodal curve are almost symmetrical, determining the coordinates of the critical point by using the diameter method is possible when a series of conodes is known. Its application to the two isotherms studied shows, unfortunately, that the curvatures are too important to allow reliable extrapolation and we have adapted this method to non-symmetrical curves by barycentric

weighting of two liquid phases. For each conode (i), the point P_i obtained satisfies the relation:

$$\vec{P}_i = w_{1i} \vec{L}_{1i} + w_{2i} \vec{L}_{2i} \quad (9)$$

where w_{1i} and w_{2i} represent respectively the mass fractions of the liquid phases of compositions \vec{L}_{1i} and \vec{L}_{2i} . The correlation coefficients R^2 of the regression lines are calculated as a function of the parameter $w=w_{2i}$, then the curve $R^2=f(W)$ is plotted as shown in the example in Fig. 4.

The value of w corresponding to the best correlation coefficient is chosen to then define the coordinates of the critical point by intersection with the binodal.

Conclusions

After a theoretical analysis of the bases of the Isoplethic Thermal Analysis intended to improve the use of thermograms, this technique was applied to the study of the water-sodium sulfate-piperidine system. The two isotherms 25 and 40°C, framing the peritectic decomposition of the sodium sulfate decahydrate, were both plotted in full and, in particular, the miscibility gap which appears was completely delimited: the critical point of the binodal and the limit conode, corresponding to the appearance of anhydrous and dehydrated sodium sulfate were identified.




Fig. 4 Correlation coefficient R^2 as a function of the mass fraction of liquid phase (L_{2i})

Lastly, the experimental results obtained constitute fundamental data for treating more or less complex solutions containing the three constituents taken into account. In particular, this study highlighted that increasing temperature favours extraction potential by liquid-liquid separation and determines the associated conditions.

Acknowledgements

Thanks to the Région Rhône-Alpes for the study funding (EMERGENCE project).

References

- 1 R. Tenu, J. J. Courniou and R. Cohen-Adad, Bull. Soc. Chim. Fr., 3–4, I (1979), 82.
- 2 R. Tenu and J. J. Courniou, Bull. Soc. Chim., 5–6 (1979) 155.
- 3 J. Berthet, R. Tenu and J. J. Courniou, Thermochim. Acta, 68 (1983) 145.
- 4 J. Berthet and J. J. Courniou, ISBN2-9504481-5-1, JEEP XX (March 1994).
- 5 J. Berthet and J. J. Courniou, Brevet F9313402 (4 November 1993).
- 6 H. C. Dickenson and E. F. Mueller, J. Am. Chem. Soc., 29 (1907) 1381.
- 7 T. W. Richards and V. Yugve, J. Am. Chem. Soc., 40 (1918) 164.

Received: January 16, 2008

Accepted: August 17, 2008

DOI: 10.1007/s10973-008-9000-8