The experiments already made show that a good quantitative separation of thorium from these elements can be based upon this reaction. When thorium is precipitated in this manner in the presence of either cerium, lanthanum, or didymium, traces of the latter elements are carried down, but may be removed by a single reprecipitation. It is intended to compare this method with those at present in use for the analysis of monazite.

A number of other weak organic acids are being investigated along the same lines, and several of these show interesting results.

QUANTITATIVE CHEMICAL LABORATORY, COLUMBIA UNIVERSITY.

## NOTE.

Reinsch's Test for Arsenic.—To obtain a black deposit upon a piece of copper is a comparatively easy matter; to prove that this deposit is due to the presence of arsenic is perhaps more difficult. In place of the usual glass tube, Sheridan Delépine<sup>1</sup> recommends a "thimble-shaped copper cone" half an inch high, inserted through a hole in a thin iron plate, the sublimate of arsenious oxide being formed on a microscope cover glass. The following simplified form of the above arrangement I have found extremely convenient for the detection of very minute quantities of arsenic.

In the center of a piece of sheet copper an inch or so square, a small depression is punched, an eighth of an inch deep and of about the same diameter. The arsenic is deposited from solution on a little piece of copper a few millimeters in area. After being washed and dried, the little piece of blackened copper is placed in the depression in the sheet metal. A microscope cover glass, having a drop of water on its upper surface, is now placed over the miniature copper crucible, and the latter is then gently heated over a very small flame. When examined under a high power of the microscope, the crystals of arsenious oxide may sometimes be seen even when no sublimate is visible to the naked eye. Careful attention to the illumination is, of course, necessary.

UNIVERSITY OF MANITOBA, WINNIPEG. EDGAR B. KENRICK.

<sup>1</sup> Abstract in J. Soc. Chem. Ind., March, 1901.

276