

[CONTRIBUTION FROM THE CHEMICAL LABORATORY OF IOWA STATE COLLEGE]

THE REACTION BETWEEN CITRONELLAL AND SOME ORGANOMAGNESIUM HALIDES

BY HENRY GILMAN AND W. F. SCHULZ

RECEIVED APRIL 28, 1930


PUBLISHED SEPTEMBER 5, 1930

Introduction

A very good case has been made for the non-addition, under usual experimental conditions, of organomagnesium halides to an ethylenic linkage.¹ There is, however, an apparent addition to such a linkage in compounds having the olefinic bond conjugated with another, and a different double bond, such as the carbonyl group in benzalacetophenone, $C_6H_5CH=CH-COC_6H_5$. Kohler and co-workers have demonstrated in an unequivocal manner that such conjugated systems can undergo 1,4-addition with the Grignard reagent. The enolic compound obtained subsequent to hydrolysis of the 1,4-addition intermediate then rearranges to give as an ultimate product a compound which might have been interpreted (and was interpreted incorrectly by earlier investigators) as involving direct addition to the ethylenic linkage.

The author of a recent book on organomagnesium halides,² in emphasizing the probability of direct addition to an ethylenic linkage, directs attention to a study by Rupe³ on the purported addition of benzylmagnesium chloride to the ethylenic linkage in citronellal and makes the pointed observation that in this compound "1,4-addition is absolutely excluded."

The following reactions illustrate Rupe's interpretation of the mechanism of reaction.



¹ Gilman and Harris, *Rec. trav. chim.*, **49**, June, (1930). References to earlier studies may be traced from this article.

² Courtot, "Le Magnésium en Chimie Organique," published in 1926 by the author at Nancy (see pages 317 and 324).

³ Rupe, *Ann.*, **402**, 161 (1914). The present controversy on the constitution of citronellal does not affect the discussion in our paper because neither the double bond in the methylene form nor the double bond in the isopropylidene form is conjugated with the carbonyl group.

not, up to the present time, any case of even a slow addition of RMgX compounds to a non-conjugated ethylenic linkage.⁶ Second, the acid (B) might actually have been formed in accordance with Reactions II and III and yet not revealed itself in the alkali extraction because of intramolecular esterification to give a ten-membered lactone. This also is highly unlikely *per se*, and particularly under our experimental conditions, because the lactone even though formed quantitatively should have been hydrolyzed in part, at least, by the alkali treatment to give the acid (B). This acid might subsequently have been converted to lactone, in part. Either or both of these products should have revealed themselves as impurities in the phenylacetic acid, but the high purity of this acid belied admixture, in any significant quantity, of contaminating lactone or acid.

Experimental Part

A solution of 23.1 g. (0.15 mole) of citronellal in 50 cc. of ether was added dropwise to a well-stirred solution containing 0.45 mole of benzylmagnesium chloride in 250 cc. of ether. Stirring and gentle refluxing were continued for fifteen minutes after addition of the aldehyde. The reaction mixture was then carbonated in a customary manner,⁷ poured into iced dilute sulfuric acid, and the ether layer separated, washed with water and extracted with 10% sodium hydroxide. The alkaline extract was washed with ether, acidified with hydrochloric acid, and the precipitate so obtained filtered, washed with a minimum of water and dried in the air. Its melting point and mixed melting point with phenylacetic acid were identical with that of phenylacetic acid. The neutralization equivalent was 137.1, and the calculated value for phenylacetic acid is 136.1.

The reaction undoubtedly proceeded in accordance with that carried out by Rupe³ because we also obtained the oil described by him (the product from hydrolysis of compound (A) in Reaction II) and its boiling point ($244^\circ/9$ mm.) agreed with that determined by him. We have not investigated the constitution of this compound.

In a corresponding experiment with phenylmagnesium bromide the benzoic acid obtained melted at the correct melting point for this acid and showed no depression in a mixed melting point determination with an authentic specimen. The neutralization equivalent was 121.2, and the calculated equivalent for benzoic acid is 122.

The *n*-valeric acid obtained from a similar reaction between citronellal and an excess of *n*-butylmagnesium bromide had the following constants: b. p. 186° ; d_{20}^{20} 0.9362; n_D^{20} , 1.404; and its neutralization equivalent was 108.8 (calcd. 102.1).

Summary

Contrary to the interpretations of others, benzylmagnesium chloride does not add to the ethylenic linkage in citronellal.

AMES, IOWA

steric hindrance about the carbonyl group will react more slowly than aldehydes without such steric hindrance.

⁶ An interesting case of 1,4-addition to a conjugated system that is part aliphatic and part aromatic has been described recently by Gilman, Kirby and Kinney, *THIS JOURNAL*, **51**, 2252 (1929). There is a somewhat remote possibility that even though there is no formal conjugated system in citronellal, there still might be addition to a conjugated system resulting from the spatial contiguity of the aldehyde and ethylenic groups.

⁷ Gilman and Parker, *ibid.*, **46**, 2816 (1924).