

hydrochloride show that there is but one break in the titration curve when all the chloride ion has been neutralized, the formula for the vitamin is $C_8H_{11}NO_8 \cdot HCl$.

The absorption spectrum studies of the vitamin hydrochloride in acid, alkaline and neutral solutions indicate that we are dealing with a compound with tautomeric properties.

Vitamin B-6 shows pronounced and well-defined absorption in the spectral region from 2300 to 3300 Å. In dilute hydrochloric acid (pH 2) there is a single band with maximum absorption at 2925 Å. The hydrochloride dissolved in water (pH 4.5) shows that the band at 2925 Å. diminished in intensity along with a new band with maximum absorption at 3275 Å. At pH 6.75 the band at 3275 Å. has increased markedly while that at 2925 Å. has disappeared and simultaneously a new band at 2560 Å. has appeared. At pH 10.2 the two bands which were present at pH 6.75 have increased in intensity and shifted toward the shorter wave lengths.

RESEARCH LABORATORY
MERCCK & CO., INC.
RAHWAY, NEW JERSEY

JOHN C. KERESZTESY
JOSEPH R. STEVENS

RECEIVED APRIL 22, 1938

HYDROLYSIS OF SUCROSE

Sir:

In the literature on the hydrolysis of sucrose by acids no mention seems to have been made of the following effect, which we have observed experimentally.

The rate of inversion of 2% sucrose solutions by hydrochloric acid at various concentrations has been measured carefully at temperatures from 0 to 35°, using the dilatometric method. It has been observed that the energy of activation varies both with temperature, as already has been shown, and with the concentration of hydrochloric acid, which has not, we believe, been noted previously.

Choosing, for example, the temperature range from 0 to 10°, we find that the energy of activation varies from 24.2 kcal. at 4.842 molar hydrochloric acid to 26.2 kcal. at 1.1235 molar hydrochloric acid.

A possible explanation based on the theory of

the transition state has been suggested for the concentration effect [private communication from Harold F. Walton, Princeton University].

DEPT. OF CHEM. & CHEM. ENG. PAUL M. LEININGER
TOWNE SCIENTIFIC SCHOOL MARTIN KILPATRICK
UNIVERSITY OF PENNSYLVANIA
PHILADELPHIA, PENNA.

RECEIVED MARCH 25, 1938

AROMATIC AMINES AS CATALYSTS FOR THE DEHYDROGENATION OF GLYCERALDEHYDE

Sir:

Transformation of glyceraldehyde into pyruvic aldehyde (methyl glyoxal) by the catalytic action of aromatic amines in dilute acetic acid solution [Strain and Spoehr, *J. Biol. Chem.*, **89**, 527 (1930)] may be regarded as an intramolecular hydrogenation-dehydrogenation reaction. In the presence of amines and hydrogen acceptors (oxygen, methylene blue and indigo carmine) hydrogen is removed from the glyceraldehyde. This dehydrogenation yields a mixture of carbonyl compounds as is shown by the formation of phenylhydrazine and 3-nitrobenzohydrazine derivatives and their chromatographic adsorption. Reaction of glyceraldehyde with oxygen in the presence of amines is accompanied by the formation of peroxides and colored insoluble products. Pyruvic aldehyde itself does not reduce methylene blue in the presence of amines, and dihydroxyacetone exhibits only a slow reducing action. Substances which might be expected to occur as dehydrogenation products of glyceraldehyde such as those produced by oxidation of dihydroxyacetone with cupric acetate [Evans and Waring, *THIS JOURNAL*, **48**, 2678 (1926)] exhibit very strong reduction. Intermolecular oxidation and reduction of glyceraldehyde and its intermediate transformation products may account for the variations in the yields of pyruvic aldehyde previously observed (Ref. 1, p. 532). These observations also indicate that aromatic amines may be regarded as catalysts (analogous to dehydrogenases) for the dehydrogenation of glyceraldehyde as well as for the molecular rearrangement of glyceraldehyde to pyruvic aldehyde.

DIVISION OF PLANT BIOLOGY HAROLD H. STRAIN
CARNEGIE INSTITUTION OF WASHINGTON
STANFORD UNIVERSITY, CALIFORNIA

RECEIVED FEBRUARY 21, 1938