

It is believed that this latter oxidation also takes place through the medium of a selenium complex. This is supported by the fact that the reaction mixture becomes yellow, then red and finally deposits selenium, and upon steam distillation leaves a non-volatile selenium compound.

Since carvopinone yields *trans*-pinocamphone upon reduction it may be seen, by setting up a model, that the carbonyl group of carvopinone is close to the *gem*-dimethyl group. This proximity would tend to hinder the rate of reaction of the carbonyl group with large molecules. The slowness of the reactivity of the carbonyl group was pointed out in the previous paper² in respect to the formation of the semicarbazone. This steric relation may also be the cause of the relative ease of the rupture of the cyclobutane ring to form carvone.

The pinocarveol formed by the action of sele-

mium dioxide on β -pinene is probably the *trans* form with respect to the hydroxyl group and the *gem*-dimethyl group. It is believed that the method of formation and purification would tend to give rise to the more stable form, which is *trans*.

Summary

1. The oxidation of β -pinene with selenium dioxide in acetic acid and acetic anhydride gave pinocarvyl acetate, pinocarveol, pinocarvone and carvopinone as steam volatile products. The main product was the acetate.

2. The oxidation product of pinocarveol by selenium dioxide is largely carvopinone.

3. The catalytic reduction of carvopinone yields 1-*trans*-pinocamphone.

4. The method of formation and the structure of some of these compounds have been proposed.

GAINESVILLE, FLORIDA

RECEIVED APRIL 10, 1942

[CONTRIBUTION FROM THE GEOCHEMICAL SECTION OF THE ILLINOIS STATE GEOLOGICAL SURVEY]

The Oxidizing Power of Illinois Coal. II. The Effects of Extended Time¹

BY G. R. YOHE² AND MYRON H. WILT³

Previous work³ has shown that Illinois coals take on the ability to act as oxidizing agents toward titanous chloride upon exposure to air or oxygen, even for very short periods of time. In the present work, a similar study has been applied to another group of coals with emphasis upon the changes in oxidizing power over longer times of exposure to air.

Preparation of Samples.—The coal samples were obtained in the form of large fresh blocks at the mines, brought to the laboratory as soon as possible, and ground to pass a 100-mesh sieve. No precautions were taken to prevent atmospheric oxidation during the preparation of the samples. After grinding, portions were taken for analyses, and the remaining stocks were placed in 2-quart mason jars and deliberately exposed to air by leaving the covers loose and by occasionally mixing the contents of each jar. The time of exposure to air was measured in days from the date of grinding.

Determination of Oxidizing Power.—The determinations of oxidizing power were made by the titanous chloride-ferric chloride method previously described.³ Parallel determinations of soluble ferric iron extractable

from the coals under similar conditions were made, and the oxidizing power data reported here are corrected for soluble ferric iron. As a rule, the soluble ferric iron values were low compared to total oxidizing power. The results obtained with five whole coal samples are shown in Fig. 1 expressed in milliequivalents per gram of coal dried in contact with laboratory air, but not calculated to a moisture- and ash-free basis. Analytical data for the coals are given in Table I.

Measurement of Specific Surface.—In order to determine whether the variations in magnitude of oxidizing power were due entirely to differences in surface exposed, specific surface measurements were made. The procedure used is an adaptation of the air-permeability method of Lea and Nurse⁴ with apparatus constructed according to specifications obtained from Battelle Memorial Institute.⁵ Table II gives the results obtained and a comparison of the maximum oxidizing power values per unit surface.

Discussion

The mechanism of the reaction of oxygen with coal is obscure and will doubtless remain so until the chemical nature of coal is more clearly revealed. A number of workers in this field have considered that the initial step is the formation of a "coal-oxygen complex" or an unstable surface

(1) Presented before the Division of Gas and Fuel Chemistry at the Memphis meeting of the American Chemical Society, April 22, 1942. Published with permission of the Chief, Illinois State Geological Survey.

(2) Associate Chemist and Research Assistant, respectively.

(3) Yohe and Harman, *THIS JOURNAL*, **63**, 555 (1941).

(4) Lea and Nurse, *J. Soc. Chem. Ind.*, **68**, 277-83 (1939).

(5) Private communication from R. A. Sherman to O. W. Rees.

TABLE I
 ANALYTICAL DATA, ILLINOIS COALS

Laboratory number	County	Seam	"As received" basis		Ash- and moisture-free basis					Calorific value, cal. per g. ^a	Bituminous coal rank (A. S. T. M.), high volatile
			Moisture	Ash	C	H	N	O	S		
C-2413	LaSalle	6	10.2	9.2	78.38	5.62	1.22	9.25	5.53	7182	C
C-2414	Fulton	5	11.7	7.4	79.93	5.57	1.47	10.43	2.60	7003	C
C-2415	Sangamon	5	11.3	6.8	78.02	5.49	1.48	9.70	5.31	7040	C
C-2422	Vermilion	6	13.1	3.2	81.05	5.65	1.76	10.05	1.49	7035 ^b	C
C-2423	Gallatin	5	2.6	10.4	81.36	5.78	1.80	7.78	3.28	7832 ^c	A

^a Moist mineral-matter-free basis. Multiply by 1.8 to obtain the B. t. u. per lb. values used in A. S. T. M. classification by rank. ^b Calculated from ultimate analysis. ^c Value obtained on another sample (C-2416) from the same source.


 TABLE II
 SPECIFIC SURFACE DATA

Coal number	Specific surface sq. cm./gram	Maximum O. P. × 10 ⁹
		Specific surface
C-2413	3270	5.84
C-2414	5450	2.86
C-2415	3510	4.37
C-2422	3410	3.30
C-2423	3850	1.96

combination.⁶⁻⁹ Although some assume this to be adsorbed oxygen,⁷ others believe it to be chemi-

cally combined, since evacuation does not lead to oxygen recovery in the way such treatment of an adsorption complex should do.¹⁰ This "complex" has been referred to as "peroxide,"^{8,11,12} yet little is known of its nature or properties.¹³ Nevertheless, this loosely bound oxygen is commonly mentioned as an intermediate stage in the atmospheric oxidation of coal.^{6,7,8}

In general form, the curves of Fig. 1 are suggestive of the existence of an intermediate such as B in consecutive reactions of the type



Although the atmospheric oxidation of coal is undoubtedly a complex process, the data presented herein may be considered as evidence supporting the hypothesis that loosely held oxygen (perhaps of a peroxide type) constitutes an intermediate stage in a part, at least, of the total change occurring in the oxidative degradation of these coals. It is of interest to note that data for the change in peroxide number of glyceryl trilinolen-




Fig. 1.—Change in oxidizing power with time of exposure to air.

(6) Wheeler, *J. Chem. Soc.*, **113**, 945-55 (1918).

(7) Parr and Milner, *Ind. Eng. Chem.*, **17**, 117 (1925).

(8) Davis and Byrne, *ibid.*, **17**, 125 (1925).

(9) Porter and Ralston, *U. S. Bur. Mines Tech. Paper*, **65**, 21-6 (1914).

(10) Winmill, *Trans. Inst. Mining Engrs.* (London), **48**, 503-7 (1914-15).

(11) Bunte and Brückner, *Angew. Chem.*, **47**, 84-6 (1934).

(12) F. Fischer, *Ges. Abhandl. Kennitus Kohle*, **4**, 454-5 (1920).

(13) Attempts to demonstrate the presence of peroxides in coal in this Laboratory have given inconclusive results. The common potassium iodide test is rendered valueless because of the avidity with which coal combines with iodine. The chemiluminescent oxidation of magnesium phthalocyanine [Cook, *J. Chem. Soc.*, 1845 (1938); Helberger and Hever, *Ber.*, **72B**, 11-15 (1939)] was tried with magnesium phthalocyanine kindly provided by Dr. R. P. Linstead of Harvard University. This reaction is apparently inhibited by coal; a chlorobenzene solution of tetralin which produced red luminescence ceased to do so when a small amount of powdered coal was added. The use of powdered coals as catalysts in the peroxide-catalyzed side chain chlorination of toluene and the addition of chlorine to the ethylenic double bond [Kharasch and Brown, *This Journal*, **61**, 2142-60, 3432 (1939)] gave results which could be explained without assuming the presence of peroxides. A modification of the Yule-Wilson ferrous sulfate test [Yule and Wilson, *Ind. Eng. Chem.*, **23**, 1254-9 (1931)] gave positive results essentially like those obtained with the titanous chloride method. Although proof of the presence of peroxide structures is lacking, such groups must still be considered as possible causes of the oxidizing power observed.

(14) Getman and Daniels, "Outlines of Theoretical Chemistry," 6th Edition, John Wiley and Sons, New York, N. Y., 1937, pp. 329-31.

ate¹⁵ and of a soybean oil¹⁶ give curves similar to those shown in Fig. 1.

There is no apparent correlation between the magnitude of the oxidizing power and properties within the group of coals having high-volatile C bituminous rank. It will be seen, however, that the one coal of higher rank (C-2423) exhibits a definitely lower oxidizing power per unit surface. It seems obvious from Table II that the magnitude of oxidizing power attained is not simply a function of the extent of the surface, but that the nature of the surface (which undoubtedly varies with the rank and source of the coal) is of utmost importance in this respect. It is hoped that this investigation may be extended to cover a series of coals of widely differing ranks.

It is possible that the different petrographic or "banded" ingredients of the coal may differ in

(15) Elm, *Ind. Eng. Chem.*, **23**, 882 (1931).

(16) Clark and Rugg, *Ind. Eng. Chem., Anal. Ed.*, **13**, 243 (1941).

their ability to take on oxidizing power. Such differences, as well as errors in sampling, might contribute to the irregularities in the curves. The present study is being continued with samples of separated banded ingredients of an Illinois coal.

Acknowledgment.—The authors wish to extend their thanks to Drs. F. H. Reed and O. W. Rees for valuable suggestions; to Mr. C. C. Boley for assistance in collecting samples and to Mr. K. F. Bursack and Mr. W. F. Wagner who made the coal analyses and specific surface measurements under the direction of Dr. Rees.

Summary

The oxidizing power exhibited by five Illinois coals has been shown to reach a maximum value and then decrease, suggesting that this "reactive oxygen" may play the role of an intermediate in the oxidative degradation of these coals.

URBANA, ILLINOIS

RECEIVED MAY 1, 1942

[CONTRIBUTION FROM THE MORLEY CHEMICAL LABORATORY OF WESTERN RESERVE UNIVERSITY]

The Oxidation of *n*-Butylboron Oxide¹

BY OLIVER GRUMMITT

The reported² auto-oxidation of *n*-butaneboronic acid (*n*-BuB(OH)₂) to the mono-*n*-butyl ester of boric acid (*n*-BuO-B(OH)₂) suggested a parallel study of the auto-oxidation of *n*-butylboron oxide, (*n*-BuBO)₃, which is the trimeric anhydride of the boronic acid. This oxidation, because of the six-electron configuration of the boron atom, offered the possibility of proceeding through the intermediate formation of an oxygen coordination compound (or peroxide) comparable to those postulated for many oxidation processes.³

Preliminary measurements⁴ on several alkylboron oxides showed that oxidation occurred readily but at widely different rates depending upon the nature of the alkyl group (Table I).

Apparently the ease of oxidation decreases with an increasing chain length of the alkyl group and increases sharply in the order primary, secondary,

(1) Presented before the Division of Organic Chemistry at the Detroit meeting of the American Chemical Society, September 12, 1940.

(2) Snyder, Kuck and Johnson, *THIS JOURNAL*, **60**, 105 (1938).

(3) Bailey, "Retardation of Chemical Reactions," Longmans, Green and Co., New York, 1937.

(4) The author wishes to thank Professor J. R. Johnson for his permission to use these data which were obtained during a du Pont Post-Doctorate Fellowship at Cornell University, 1936-1938, and for his suggestions on this problem.

TABLE I

R of R-BO	Time required for the consumption of 2.0 cc. oxygen, 25°, min.
Et	33
<i>n</i> -Pr	36
<i>n</i> -Bu	71
<i>s</i> -Bu	3
<i>t</i> -Bu	2
<i>n</i> -Hex	105

and tertiary. The relative stability of the corresponding boronic acids in air is also of this order.^{2,5}

Experimental Procedure

Apparatus.—The reaction flask is shown in Fig. 1. The total volume of this two-bulb flask is approximately 125 cc. and since none of the dimensions is of critical importance, they are not shown. Connected to the side arm of the flask are two mercury-filled gas burets in series. The first of these burets was made of a 10-cc. delivery pipet calibrated in divisions of 0.05 cc. and a three-way stop-cock and was used to measure the consumption of oxygen during the earlier stages of the reaction. The second buret was the orthodox 100-cc. type with 0.2-cc. divisions. Each gas buret was fitted with a leveling bulb and a jacket through which thermostated water could be circulated. The use of two burets permitted accurate readings at the start of the auto-oxidation when an induction

(5) Johnson, Van Campen and Grummitt, *THIS JOURNAL*, **60**, 111 (1938); and unpublished work of Johnson and Grummitt.