

chloride,³ warmed until no more nitric oxide was evolved (five minutes), and then distilled to dryness under reduced pressure. The residue was dissolved in 8 ml. of water and treated with excess sodium hydroxide. The mixture was boiled, copper oxide was removed, and the filtrate was shaken with 100 mg. of benzoyl chloride. Acidification gave a white precipitate which crystallized from toluene in the form of colorless prisms (100 mg., 71%), m. p. 149–150° alone or mixed with an authentic sample¹ of 1-benzoylpiperidine-4,β-propionic acid.

The author thanks Mr. Stanley T. Rolfson for the analyses reported in this paper.

(3) Jones and Kenner, *J. Chem. Soc.*, 711 (1932).


SCHOOL OF CHEMISTRY
UNIVERSITY OF MINNESOTA
MINNEAPOLIS, MINNESOTA

RECEIVED MARCH 25, 1944

Preparation of 21-Pyridinium-3-β-hydroxy-5-pregnene-20-one Halides and 3-β-Hydroxy-5-androstene-17-carboxylic Acid

BY L. CARROLL KING

In a recent paper from this Laboratory the preparation of substituted β-ketoalkylpyridinium iodides by reaction of ketones with iodine and pyridine was reported.¹ This reaction has now been extended to 3-β-hydroxy-5-pregnene-20-one (I), which gave 21-pyridinium-3-β-hydroxy-5-pregnene-20-one iodide (II) in 50% yield. Alkaline decomposition of (II) or of the corresponding perchlorate yielded 93% of crude 3-β-hydroxy-5-androstene-17-carboxylic acid (III) from which the pure methyl ester was obtained (83% yield from II). The main reactions may be formulated as



(1) King, *This Journal*, **66**, 894 (1944).

Reich and Reichstein² obtained compounds of type II when a 21-halo-3-β-hydroxy-5-pregnene-20-one was treated with pyridine.

Experimental³

21-Pyridinium-3-β-hydroxy-5-pregnene-20-one Iodide (II).—A solution containing 3.12 g. of 3-β-hydroxy-5-pregnene-20-one in 75 ml. of pyridine was treated with 2.56 g. of iodine in a manner previously described.¹ The resulting relatively insoluble pyridinium iodide was separated from pyridine hydroiodide and crystallized from methanol; yield 3.053 g.; m. p. 248–250°.

Anal. Calcd. for $\text{C}_{26}\text{H}_{36}\text{INO}_2$: C, 59.91; H, 6.91. Found: C, 59.50; H, 6.83.

21-Pyridinium-3-β-hydroxy-5-pregnene-20-one Perchlorate.—This substance was prepared from the above pyridinium iodide as previously described.¹ The product was crystallized from methanol and apparently contains water of crystallization; m. p. 255–260° dec.

Anal. Calcd. for $\text{C}_{26}\text{H}_{36}\text{ClNO}_6 \cdot \frac{1}{2}\text{H}_2\text{O}$: C, 62.07; H, 7.37. Found: C, 62.12; H, 7.04.

Methyl 3-β-Hydroxy-5-androstene-17-carboxylate.—To a suspension of 656 mg. of the above pyridinium iodide (II), or a corresponding quantity of the perchlorate, in water or 50% ethyl alcohol, 0.5 g. of sodium hydroxide was added. The mixture was heated for two hours on the steam-bath, acidified and the acidic fraction isolated. The crude 3-β-hydroxy-5-androstene-17-carboxylic acid, yield 400 mg., was of a light tan color and melted at 255–270°. On recrystallization from methanol or dioxane the melting point increased to 270–275°. The total crude product was then dissolved in methanol, a few drops of acetyl chloride added and the reaction mixture heated one hour. The methyl ester was recovered and crystallized from methanol; yield 332 mg.; m. p. 174–177°.

Anal. Calcd. for $\text{C}_{21}\text{H}_{32}\text{O}_3$: C, 75.83; H, 9.70. Found: C, 75.69; H, 9.37.

The above methyl 3-β-hydroxy-5-androstene-17-carboxylate was converted to the benzoate, m. p. 204–206°, and to the acetate, m. p. 153–154°.

(2) Reich and Reichstein, *Helv. Chim. Acta*, **22**, 1124 (1939).

(3) All melting points were observed on a Fisher-Jones melting point block. Carbon and hydrogen analyses by T. S. Ma.

CHEMISTRY DEPARTMENT
NORTHWESTERN UNIVERSITY
EVANSTON, ILLINOIS

RECEIVED JUNE 5, 1944

NEW COMPOUNDS

ω-Nitro-3-vinyl-pyrene

In the manner described for the preparation of nitrostyrene,¹ equimolecular amounts of 3-pyrene-aldehyde and nitromethane were condensed in the presence of methylalcoholic sodium hydroxide solution. Hydrochloric acid precipitated the ω-nitro-3-vinyl-pyrene as red crystals, which after recrystallization from a mixture of acetone and benzene had m. p. 177°.

Anal. Calcd. for $\text{C}_{18}\text{H}_{11}\text{O}_2\text{N}$: C, 79.1; H, 4.0. Found: C, 78.6; H, 3.9.

Alcoholic alkali gives a yellow solution, but the substance is not satisfactory as an indicator.

DEPARTMENT OF ORGANIC CHEMISTRY
HEBREW UNIVERSITY
JERUSALEM, PALESTINE

ELIAHU BOGRACHOV

RECEIVED JUNE 12, 1944

(1) "Organic Syntheses," Coll. Vol. I, p. 413.