

cance. The constitutional significance of the results obtained in this investigation is under consideration.

Experimental

Numerous variations of procedure were employed but the following is cited as typical. The electrolysis cell was constructed of 3-inch Pyrex piping.¹ The assembled, 3-compartment cell consisted of two 90° L's for ends, connected by a T. The 3 compartments were assembled with the usual gaskets and metal joint flanges, and with the compartments separated by a membrane of parchment paper. During reaction, the entire cell assembly was immersed in running water to control temperature and both anolyte and catholyte were stirred vigorously. The anode consisted of 99.9% pure lead sheet 6 × 70 × 100 mm. The cathode was a pool of mercury in a beaker 50-mm. in diameter. The distance from anode center to cathode center was about 52 cm.

Twenty grams of dry butanol lignin² from western hemlock (*Tsuga heterophylla* Sarg.) was dissolved in 1500 ml. of 1% sodium hydroxide solution. Half of this solution was placed in the anode compartment and half in the cathode compartment. The center compartment was filled to the same level with 1% sodium hydroxide solution.

An e. m. f. of 60 volts was applied which caused a passage of 3 amperes. As the current increased, a small amount of solution from the center compartment was withdrawn and replaced with water, maintaining the current approximately constant at 3 amperes. The center compartment served the important functions of controlling current, preventing undue heating, preventing foaming and providing a visual check on undesirable migration. The temperature of the electrolyte was 30 to 35°, and the time of electrolysis was forty-eight hours. Both anolyte and catholyte became straw-colored in this time, the latter more rapidly, and produced maximum yields of products.

Anolytes and catholytes were acidified and extracted with ether, the volume of each reduced under a 34-plate column, re-extracted, and rough yields determined. The aqueous solution of anolyte was then evaporated to dryness

(1) Corning Glass Works, Pyrex Brand Piping and Heat Exchangers, Bulletin 814, Corning, N. Y., 1940.

(2) A. Bailey, *Paper Trade J.*, **111**, 63-6 (1940); *Tech. Assoc. Papers*, **24**, 608-11 (1941), entitled "Preparation and Properties of Butanol Lignin."

and the solids left were extracted with acetone. Eight or ten cycles of grinding the solids and re-extracting with acetone were necessary to recover all soluble material.

Anolyte-ether extracts from a number of identical electrolyses were combined and fractionated directly (without bicarbonate extraction, etc.) in efficient columns of small size of the helices and twisted-screen types. Low pressure distillation was employed on fractions not sufficiently volatile to ascend the columns. Molecular distillation was used on the fractions of lowest volatility. A pressure of 10⁻⁶ mm. of mercury at the distilling gap was established by means of a three-stage, four-compartment oil diffusion pump using Octoil-S. Anolyte-acetone extracts were treated separately and similarly.

Identification of Products

1. **Ethyl Methyl Ketone.**—The 2,4-dinitrophenylhydrazone melted at 113.1° and the semicarbazone at 134.3°.
2. **Acetone.**—The 2,4-dinitrophenylhydrazone melted at 126° and the *p*-nitrophenylhydrazone at 147.2°.
3. **Acetic Acid.**—The *p*-nitrobenzyl ester melted at 75-76° and the *p*-phenylphenacyl ester at 109.5°.
4. **β -Resorcylic Acid.**—The *p*-nitrobenzyl ester melted at 187-188°. When methylated with dimethyl sulfate and alkali and purified, yielded acid, m. p. 106.0-106.3° (2,4-dimethoxybenzoic acid, m. p. 108°).
5. **Protocatechuic Acid.**—The *p*-nitrobenzyl ester melted at 186°, the amide at 209-210° and the anilide at 164°.
6. ***p*-Hydroxybenzoic Acid.**—The *p*-nitrobenzyl ester melted at 180-181° and the *p*-phenylphenacyl ester at 238.5°.
7. ***m*-Toluic Acid.**—The *S*-benzylthiuronium ester melted at 162.6° and the *p*-bromophenacyl ester at 107°.
8. **Oxalic Acid.**—Sublimed below 100°, the *p*-phenylphenacyl ester melted at 164° and the *S*-benzylthiuronium ester at 192-193°.
9. **Isobutyl Methyl Ketone.**—The semicarbazone melted at 130.0-130.5° and the 2,4-dinitrophenylhydrazone at 131°.

Summary

Lignin was electrolytically oxidized and reduced. Oxidation products were obtained in good yield and identified as ketones and acids.

SEATTLE 5, WASHINGTON RECEIVED SEPTEMBER 24, 1945

[CONTRIBUTION FROM THE MINER LABORATORIES]

Diglycerol by a New Ether Synthesis¹

BY H. J. WRIGHT² AND R. N. DU PUIS

Diglycerol has been prepared by a new series of reactions between glycerol, calcium hydroxide or calcium oxide, and carbon dioxide. The course of these reactions, which may be carried out to yield diglycerol with substantial absence of higher polyglycerols, has not been definitely established. Theories concerning the mechanism of the reactions have been devised from experimental data and are presented in the following discussion.

The reactions between an alcohol, calcium oxide or calcium hydroxide, and carbon dioxide to yield the ether of the alcohol have never been reported

(1) This research was sponsored by the Association of American Soap and Glycerine Producers, Inc.

(2) Present address: 700 Desnoyer Street, Kaukauna, Wisconsin.

in the literature so far as we are aware. It was thought, therefore, that although such a reaction has been applied only to glycerol in the Laboratory and the data are admittedly incomplete, publication of the procedures used and results obtained might be of interest to workers in related fields.

Reactions between glycerol and calcium hydroxide to form calcium glyceroxides have been studied previously.^{3,4} Briefly, it has been found that heating calcium oxide or calcium hydroxide with glycerol results in formation of calcium monoglyceroxide, I, Ca[C₃H₅(OH)O₂] or calcium diglyceroxide, II, Ca[C₃H₅(OH)₂O]₂ and water.

(3) Grün and Bockisch, *Ber.*, **41**, 3476 (1908).

(4) Wheeler, *Chem. News*, **142**, 241-243 (1931).

Acknowledgment.—The authors wish to acknowledge the assistance of E. E. Langdon and H. L. Wehrmeister in various phases of this work and the literature search made by Miss V. M. King.

Summary

Diglycerol has been prepared by a new series

of reactions involving treatment of glycerol with calcium oxide or calcium hydroxide and carbon dioxide.

Some possible mechanisms of the reaction are suggested and experimental procedures and results obtained are reported.

CHICAGO 6, ILLINOIS

RECEIVED SEPTEMBER 19, 1945

[CONTRIBUTION FROM THE COLLEGE OF PHARMACY OF THE UNIVERSITY OF MINNESOTA]

4-Thio and 2,4-Dithio 5,5-Dialkylthiobarbituric Acids

BY JAMES H. BOOTHE¹ AND CHARLES O. WILSON

Early in the history of barbituric acid synthesis, a few 2-thiobarbiturates were prepared for the purpose of converting them into the corresponding oxygen analogs. In fact, Fischer and Dilthey² reported the preparation of 5,5-diethyl-2-thiobarbituric acid in their original paper in 1904. Until 1935, however, no systematic attempt was made to prepare a series of these compounds and evaluate them pharmacologically. This was probably due to the fact that Fischer and Mering³ reported that the administration of 120 mg. per kg. of the sulfur analog of barbital to a dog produced a deep sleep, followed by death. This fact was interpreted by Fraenkel⁴ as conclusive evidence that sulfur imparted a pronounced toxic character to barbital. Dox and Hjort⁵ also reported in 1927 that this compound caused tremors during anesthesia as well as pre-anesthetic excitement and cyanosis.


In spite of these adverse indications of the toxicity of the thiobarbiturates, two separate groups of workers, namely, Volwiler and Tabern⁶ and Miller, Munch and Crossley^{7,8} in 1935, reported independently systematic studies of the preparation and pharmacological properties of a number of 2-thiobarbiturates. They found that these compounds had, in general, a rapid onset of action and were destroyed much more rapidly in the body than were their oxygen analogs. Because of these properties, the 2-thiobarbiturates are proving very successful as intravenous anesthetics.

Until recently, no substituted barbituric acids in which more than one of the oxygen atoms were replaced by sulfur were known, although the preparation of 2,4,6-trithiobarbituric acid by the action of potassium hydrosulfide on 2,4,6-trichloropyrimidine had been described.⁹ During the course of this study, there appeared a paper

by Henze and Smith¹⁰ dealing with polythiobarbiturates in which was reported the preparation of 5,5-diethyl-2,4,6-trithiobarbiturate and 5-phenyl-5 ethyl-2,4,6-trithiobarbiturate. It was stated that these possessed no hypnotic properties. Carrington¹¹ has prepared the 2-thio, 2,4-dithio and 2,4,6-trithio derivatives of a series of barbiturates. He used those barbiturates that are well-known hypnotics, but reported no pharmacological data.

In view of the fact that no synthesis of 4-thiobarbiturates has been reported in the literature and that the 2,4-dithiobarbiturates reported by Carrington have not been tested pharmacologically, and since such thio compounds might exhibit interesting physiological activity, it was decided to synthesize, if possible, several typical examples of these thio compounds.

Several workers¹²⁻¹⁶ have replaced an imino



(1) Lederle Laboratories, Inc., Research fellow, 1942-1943. Present address: Lederle Laboratories, Inc., Pearl River, New York.

(2) Fischer and Dilthey, *Ann.*, **335**, 334 (1904).

(3) Fischer and Mering, *Therap. Gegen.*, **101**, 97 (1903).

(4) Fraenkel, "Die Arzneimittel Synthese," 6th ed., 1927, p. 510.

(5) Dox and Hjort, *J. Pharmacol.*, **31**, 455 (1927).

(6) Volwiler and Tabern, *THIS JOURNAL*, **57**, 1961 (1935).

(7) Miller, Munch and Crossley, *ibid.*, **58**, 1090 (1936).

(8) Miller, Munch and Crossley, *Science*, **81**, 615 (1935).

(9) Büttner, *Ber.*, **36**, 2234 (1903).

(10) Henze and Smith, *THIS JOURNAL*, **65**, 1090 (1943).

(11) Carrington, *J. Chem. Soc.*, 124 (1944).

(12) Hofmann, *Ber.*, **2**, 460 (1869).

(13) Hobrecker, *ibid.*, **2**, 689 (1869).

(14) Bernthsen, *ibid.*, **10**, 1240, 38 (1877).

(15) Matsui, *Mem. Coll. Sci. Eng. Kyoto Imp. Univ.*, **1**, 285 (1908).

(16) Delépine, *Comp. rend.*, **153**, 281 (1911).