

yielding the desired 3,4-bis-(3'-methyl-4'-hydroxy-5'-allylphenyl)-2,4-hexadiene, as a red gum. This tetraalkyl substituted analog of dienestrol was characterized by the preparation of the corresponding diacetate and dipropionate.

The estrogenic assays on these compounds were carried out by administering orally the test sample to ovariectomized female rats. Vaginal smears were examined and only full cornification was regarded as a positive oestrus response. The results of preliminary measurements of gross estrogenic activity indicate that at the 100 microgram dose level; the 3,4-bis-(3'-methyl-4'-propiony-5'-allylphenyl)-2,4-hexadiene shows a 20% oestrus response; the 3,4-bis-(3'-methyl-4'-acetoxy-5'-allylphenyl)-2,4-hexadiene gives a 25% response, and the 3,4-bis-(3'-methyl-4'-hydroxy-5'-allylphenyl)-2,4-hexadiene shows a 35% oestrus response. Control rats injected with 0.65 microgram of U.S.P. Reference Estrone produced a 40% oestrus response.

Acknowledgment.—The authors desire to express their appreciation to Dr. C. T. Van Meter of Reed and Carnrick, Jersey City, N. J., for the estrogenic assays.

Experimental⁵

3,4-bis(3'-Methyl-4'-allyloxyphenyl)-2,4-hexadiene.—A mixture of 15 g. (0.05 mole) of 3,4-bis-(3'-methyl-4'-hydroxyphenyl)-2,4-hexadiene,³ 15 g. (0.11 mole) of allyl bromide, 15 g. (0.1 mole) of anhydrous potassium carbonate, and 200 ml. of dry acetone were refluxed for six hours, and then set aside at room temperature for twenty-four hours. Water was added to the reaction mixture, it was extracted with ether, and the ether layer was extracted with ice cold 10% aqueous sodium hydroxide solution to remove any unreacted dihydroxy compound. The excess alkali was removed by washing with water, the neutral ether solution was dried over anhydrous potassium carbonate, filtered, and the solvent removed by distillation. Crystallization and re-crystallization from dilute ethanol yielded 11.2 g. (59%) of a white crystalline solid; m. p. 67.5–68°.

Anal. Calcd. for $C_{26}H_{30}O_2$: C, 83.42; H, 8.02. Found: C, 83.57; H, 8.31.

3,4-bis-(3'-Methyl-4'-hydroxy-5'-allylphenyl)-2,4-hexadiene.—The procedure used by Kaiser and Svarz⁴ for the preparation of 3,4-bis-(3'-allyl-4'-hydroxyphenyl)-hexene-3 was employed. A solution of 6.8 g. of 3,4-bis-(3'-methyl-4'-allyloxyphenyl)-2,4-hexadiene in 40 ml. of diethylaniline was refluxed for five hours in a nitrogen atmosphere. The desired product was isolated as a red gum (2.2 g. or 32%) which could not be crystallized, but gave proper analytical results.

Anal. Calcd. for $C_{26}H_{30}O_2$: C, 83.42; H, 8.02. Found: C, 83.20; H, 8.26.

The diacetate, prepared in 68% yield by refluxing in acetic anhydride for four hours, formed white crystals from dilute ethanol, m. p. 125–126°.

Anal. Calcd. for $C_{30}H_{34}O_4$: C, 78.60; H, 7.42. Found: C, 78.31; H, 7.29.

The dipropionate was prepared in 61% yield in a similar manner, m. p. 114–115°.

Anal. Calcd. for $C_{32}H_{38}O_4$: C, 79.01; H, 7.81. Found: C, 78.83; H, 7.71.

CHEMICAL LABORATORIES
WASHINGTON SQUARE COLLEGE
NEW YORK UNIVERSITY
NEW YORK, N. Y.

RECEIVED OCTOBER 26, 1948

Closed-Circuit Seals in Glassblowing

BY ROGER K. TAYLOR

In fabricating glass apparatus in which one part is multiply connected to the rest, the customary procedure is to make the seals after the first by using a thin rod to pull together or bridge over the edges of the open-ended immobilized tubes that are to be joined. Difficulties occasionally arise, especially when the seal is not readily accessible from all directions, in manipulating the rod and obtaining a completely tight seal.

The writer for some years has on occasion used an alternative method which avoids these troubles. Although it is difficult to believe that the same idea has not occurred to others, neither the method nor a description of it has been seen elsewhere; and it is in the hope that the scheme may be of some general assistance that the following exposition is offered.




Fig. 1.—Closed-circuit seals in glassblowing.

The innovation consists in closing the ends of the tubes where the joint is to be made (a). Then, when both ends are hot, blowing brings together the bulbs formed, producing a septum (b). When the enlarged portion is permitted to shrink (c), a portion of the glass of the septum is picked up by the wall; alternate blowing and shrinking (d, c) ultimately reduces the thickness of the septum to the point that it breaks (e), after which the joint is readily smoothed out (f). Glass can of course be added or removed with a thin rod as necessary, if the wall thickness at the seal does not match that of the rest of the tubing.

An advantage is that the annoyance of air leakage is avoided in the blowing, during making the seal; and this is particularly important when several connections have to be made between immobilized parts. It is also unnecessary to fit the parts so precisely, to give the small clearances required for easy working when the tubes are open. Furthermore there is no chance of leaving pinholes in the seals. The chief disadvantage is that prolonged working of the glass may lead to devitrification.

THE DAVISON CHEMICAL CORPORATION
RESEARCH DIVISION
BALTIMORE 3, MD.

RECEIVED NOVEMBER 18, 1948

(5) All melting points given are uncorrected.