

Propargylglycine.—The diethyl propargylformamidomalonic acid was hydrolyzed by refluxing 2.4 g. (0.01 mole) of the ester with 24 ml. of 8% sodium hydroxide for four hours. The mixture was diluted to 100 ml. with water and passed through a column of Duolite C-10H which removed all the base. The effluent was subsequently passed through a column of Duolite A-2 which removed the formic acid produced. The resulting solution was concentrated *in vacuo*, decolorized with Darco G-60, and on addition of acetone, the amino acid crystallized. The yield was 88.5%. An analytical sample of propargylglycine was obtained from water and acetone, m. p. 243° with decomposition.

Anal. Calcd. for $C_6H_7O_2N$: C, 53.09; H, 6.24; N, 12.39. Found: C, 53.24; H, 6.20; N, 12.47.

Norvaline.—Hydrogenation of 113 mg. (0.001 mole) of propargylglycine in the presence of Adams catalyst, at 28° and 1 atmosphere, required 0.00207 mole (104% of theoretical) of hydrogen and produced 100 mg. of norvaline, which melted with decomposition at 297° in a sealed tube; the benzoyl derivative prepared as above, m. p. 150–151°.

Microbiological Tests.—Propargylglycine was tested as an inhibitor of the growth of *Escherichia coli*, strain 9723 and *Saccharomyces cerevisiae*, strain 139.⁶ To inhibit the growth of these microorganisms, to 50% of normal growth, required 4 micrograms per 7.5 ml. of medium for *S. cerevisiae* and 65 micrograms for *E. coli*. According to these data propargylglycine is more active than allylglycine for yeast but less active for the inhibition of *E. coli*.

(6) The methods employed and the test organisms were the same as those previously described.^{3a}

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF COLORADO
BOULDER, COLORADO

HERMAN GERSHON
JOHN S. MEEK
KARL DITTMER

RECEIVED AUGUST 5, 1949

N¹⁰-NITROPTEROYLGLUTAMIC ACID

Sir:

The action of nitrous acid upon 2-aminopteridines has led to destruction of the ring system,¹ desamination in the 2-position,² and simultaneous desamination of the 2-position and nitrosation on the nitrogen atom in the 10-position of pteric acid.³ "Folic acid" concentrates from natural sources⁴ were inactivated by nitrous acid under the conditions of the Van Slyke determination.⁵ These experiments involved the use of excess nitrous acid and temperatures ranging from "room temperature" to 100° or higher.

We wish to report that pteroylglutamic acid in cold hydrochloric acid solution reacts quantitatively with one mole of nitrous acid to form N¹⁰-nitropteroylglutamic acid, which precipitates from the reaction mixture as a white solid.

N¹⁰-Nitropteroylglutamic acid gives a positive Liebermann nitroso reaction. The nitroso group can be removed by treatment with phenol and hydrochloric acid, and pteroylglutamic acid thus regenerated.

Under substantially the same conditions, the

- (1) Schopf and Kottler, *Ann.*, **539**, 134 (1939).
- (2) Wieland, *et al.*, *ibid.*, **507**, 245 (1933); Wittle, *et al.*, *THIS JOURNAL*, **69**, 1780 (1947); Taylor and Cain, *ibid.*, **71**, 2538 (1949).
- (3) Wolf, *et al.*, *ibid.*, **69**, 2758 (1947).
- (4) Mitchell and Williams, *ibid.*, **66**, 272 (1944).
- (5) Van Slyke, *J. Biol. Chem.*, **16**, 121 (1913).

following give N¹⁰-nitroso compounds: pteroyl- α -glutamylglutamic acid⁶; pteroyl- γ -glutamyl- γ -glutamylglutamic acid,⁷ 9-methylpteroylglutamic acid,⁸ 4-aminopteroylglutamic acid,⁹ 2-dimethylaminopteroylglutamic acid,¹⁰ and 4-(1-piperidyl)-pteroylglutamic acid.¹⁰

Neither 2-amino-4-hydroxy-6-methylpteridine,¹¹ 2,4-diamino-6-methylpteridine,^{9b} nor N¹⁰-methylpteroylglutamic acid¹² react appreciably with nitrous acid under these conditions.

In a typical experiment, 4.4 g. of pteroylglutamic acid (90% purity,¹³ 0.5% *p*-aminobenzoylglutamic acid, 8% H₂O) was dissolved in 50 ml. of concentrated hydrochloric acid, and cooled to 5–10° by the addition of ice. Then 0.7 g. of sodium nitrite dissolved in a little water was added slowly. A white precipitate formed, which was filtered, washed, and dried to give 3.2 g. of N¹⁰-nitropteroylglutamic acid. One gram of this dissolved in 25 ml. of 5 *N* sodium hydroxide was clarified with activated charcoal. On standing the sodium salt crystallized. It was collected, dissolved in water and precipitated with acid, filtered, washed, and dried two hours at 100° (2 mm.). *Anal.* Calcd. for $C_{19}H_{18}N_8O_7$: C, 48.5; H, 3.83; N, 23.8. Found: C, 48.9; H, 4.77; N, 24.05 (corrected for 3.8% ash).

Through the courtesy of our colleagues, Dr. B. L. Hutchings, and Dr. J. J. Oleson, of the Lederle Laboratories Division, American Cyanamid Company, N¹⁰-nitropteroylglutamic acid has been tested in a preliminary way in the nutrition of *S. faecalis* R and the chick, and in both cases appears to be equivalent to pteroylglutamic acid.

(6) Mowat, *et al.*, *THIS JOURNAL*, **70**, 1096 (1948).

(7) Boothe, *et al.*, *ibid.*, **70**, 1099 (1948).

(8) Hultquist, *et al.*, *ibid.*, **71**, 619 (1949).

(9) (a) Seeger, Smith and Hultquist, *ibid.*, **69**, 2567 (1947)

(b) Seeger, *et al.*, *ibid.*, **71**, 1753 (1949).

(10) Roth, Smith and Hultquist, in press.

(11) Mowat, *et al.*, *THIS JOURNAL*, **70**, 17 (1948).

(12) Cosulich and Smith, *ibid.*, **70**, 1922 (1948).

(13) Hutchings, *et al.*, *J. Biol. Chem.*, **168**, 705 (1947).

CALCO CHEMICAL DIVISION
AMERICAN CYANAMID COMPANY DONNA B. COSULICH
BOUND BROOK, NEW JERSEY JAMES M. SMITH, JR.

RECEIVED AUG. 15, 1949


STEROLS. VIII.¹ 17 α -HYDROXYPROGESTERONE AND 17 α -HYDROXY-11-DESOXYCORTICOSTERONE

Sir:

Recently we reported¹ the facile preparation of 16,17-oxidoprogesterone from 16,17-oxido-5-pregnene-3 β -ol-20-one acetate (I). We now wish to record the use of I as an intermediate for a new and very simple partial synthesis of both 17 α -hydroxyprogesterone and 17 α -hydroxy-11-desoxycorticosterone acetate (V) (acetate of Reichstein's Compound S). The reactions as applied to the latter are schematically presented by formulas I \rightarrow V. A treatment of 16,17-oxidoprogesterone

(1) For paper VII in this series see *THIS JOURNAL*, **71**, 756 (1949)

similar to that given IV produced 17 α -hydroxyprogesterone.



The smooth cleavage of the oxido compounds by hydrogen bromide, the clean reductive removal of bromine from the resulting bromohydrins by Raney nickel, and the good yields encountered at every step make these hitherto difficultly accessible cortical hormones readily available.

16,17-Oxido-5-pregnene-3 β ,21-diol-20-one 21-Acetate (III).—From 16,17-oxido-5-pregnene-3 β -ol-20-one acetate by treating first with one molar equivalent of bromine in acetic acid-carbon tetrachloride, then with hydrogen bromide in acetic acid followed by a second molar equivalent of bromine. Then hydrolysis with hydrogen bromide in methanol-benzene, treatment with sodium iodide in benzene-ethanol and finally with potassium acetate in acetone gave needles from acetone-petroleum ether, m. p. 190–192°; $[\alpha]^{25\text{D}} +15^\circ$ (chloroform). *Anal.* Calcd. for $\text{C}_{23}\text{H}_{32}\text{O}_5$: C, 71.11; H, 8.30. Found: C, 70.86; H, 8.33.

16,17-Oxido-4-pregnene-21-ol-3,20-dione Acetate (IV).—By the Oppenauer oxidation of III; prisms from ether-petroleum ether, m. p. 170–172°; $[\alpha]^{25\text{D}} +167^\circ$ (chloroform). *Anal.* Calcd. for $\text{C}_{23}\text{H}_{30}\text{O}_5$: C, 71.48; H, 7.82. Found: C, 71.82; H, 8.23.

17 α -Hydroxy-11-desoxycorticosterone Acetate (V).—From IV with hydrogen bromide in acetic acid, followed by reduction with Raney nickel in ethanol; needles from

methanol, m. p. 235–238° (sinters 230°), no depression with material isolated from a natural source²; $[\alpha]^{25\text{D}} +114^\circ$ (acetone). *Anal.* Calcd. for $\text{C}_{23}\text{H}_{32}\text{O}_5$: C, 71.11; H, 8.30. Found: C, 70.94; H, 8.28.

(2) We are indebted to Dr. Marvin H. Kuizenga of The Upjohn Laboratories for a sample of this material.

THE GLIDDEN COMPANY
RESEARCH LABORATORIES
SOYA PRODUCTS DIVISION
CHICAGO 39, ILLINOIS

PERCY L. JULIAN
EDWIN W. MEYER
WILLIAM J. KARPEL
ISABELLE RYDEN

RECEIVED AUGUST 22, 1949

ELECTRON DEFICIENT COMPOUNDS. IV. THE STRUCTURE OF HEXAMETHYLDIPLATINUM

Sir:

The reaction $(\text{CH}_3)_3\text{PtI} + \text{K}$ was presumed to yield $(\text{CH}_3)_3\text{Pt}-\text{Pt}(\text{CH}_3)_3$.¹ A recent explanation of electron deficient bonding² suggests that since Pt^{IV} has six low energy orbitals, d^2sp^3 , the configuration about platinum should be octahedral, requiring polymerization through electron deficient bonds.

"Hexamethyldiplatinum" is monoclinic, $a_0 = 17.35$, $b_0 = 18.99$, $c_0 = 17.79$ kX., $\beta = 116^\circ$, space group from Weissenberg diagrams, $\text{P}2_1/c$. The observed density, 3.65, requires 48 $\text{Pt}(\text{CH}_3)_3$ per unit cell. There can be only 2 or 4 crystallographically equivalent atomic aggregates per unit cell. In molecular crystals these aggregates are almost invariably the molecules, so that the unit cell and density data indicate a minimum molecular size of $[(\text{CH}_3)_3\text{Pt}]_{12}$. Our data do not exclude continuous chains.

Patterson functions, $P(x,z)$ and $P(x,y)$, have been calculated, each involving about 125 reflections. All important peaks of $P(x,z)$ (Fig. 1) are interpreted, even as to heights, in terms of 16 platinum positions. Since h is even except for 13 weak reflections, there is a pseudo eight-fold set of positions: $x, z; \bar{x}, \bar{z}; x, \frac{1}{2} + z; \bar{x}, \frac{1}{2} - z; \frac{1}{2} + x, z; \frac{1}{2} - x, \bar{z}; \frac{1}{2} + x, \frac{1}{2} + z; \frac{1}{2} - x, \frac{1}{2} -$




Fig. 1.—Patterson projection, $P(xz)$ for "Hexamethyldiplatinum."

(1) H. Gilman and W. Lichtenwalter, *THIS JOURNAL*, **61**, 957 (1939).

(2) R. Rundle, *ibid.*, **69**, 1327 (1947); *J. Chem. Phys.*, **17**, 671 (1949).