

TABLE I

Compound	M.p., ^a °C.	Carbon, % ^b		Hydrogen, % ^b		Nitrogen, % ^b	
		Calcd.	Found	Calcd.	Found	Calcd.	Found
4-(Benzamidoacetyl)-biphenyl	182-183	79.98	79.61	5.44	5.41
4-(2-Furamidoacetyl)-biphenyl	145	74.75	74.84	4.92	4.99
2-(Myristamidoacetyl)-naphthalene	105-107	78.94	79.02	9.43	9.40	3.54	3.55
2-(Benzamidoacetyl)-naphthalene	143-144	78.87	78.76	5.23	5.13	4.84	4.77
2-Phenyl-5-(4-biphenyl)-thiazole	191-192	80.48	80.06	4.82	4.96
2-(2-Furyl)-5-(4-biphenyl)-thiazole	152	79.43	79.29	4.56	4.51	4.88	4.82
2-Phenyl-5-(4-biphenyl)-oxazole	158	84.85	84.79	5.06	5.19	4.71	4.45
2-Tridecyl-5-(2-naphthyl)-oxazole	56-57	82.71	82.29	9.34	9.50
2-Phenyl-5-(2-naphthyl)-oxazole	106	84.11	83.75	4.83	5.02

^a All melting points taken on a Fisher-Johns melting point block. ^b Microanalyses performed by Micro-Tech Laboratories, Skokie, Illinois.

obtained by reaction with sulfuric acid³ at room temperature or heating at 200-300° with phosphorus pentoxide. Thiazoles were obtained by heating with phosphorus pentasulfide⁴ at 200-300°. The oxazoles and thiazoles described in the table are being investigated for their use as phosphors in scintillation counters.⁵

(3) J. Lister and R. Robinson, *J. Chem. Soc.*, **101**, 1297 (1912).

(4) S. Gabriel, *Ber.*, **43**, 137 (1910).

(5) H. Kallman and M. Furst, *Phys. Rev.*, **79**, 857 (1950).

DEPARTMENT OF CHEMISTRY
ILLINOIS INSTITUTE OF TECHNOLOGY
CHICAGO, ILLINOIS

DEPARTMENT OF CHEMISTRY
NORTHWESTERN UNIVERSITY
EVANSTON, ILLINOIS

F. N. HAYES

L. C. KING

D. E. PETERSON

RECEIVED AUGUST 13, 1951

Chlorination of 4-Dibenzothiophenol

A solution of 3.4 g. (0.025 mole) of sulfuryl chloride in 50 ml. of chloroform was added dropwise to a well-stirred suspension of 5 g. (0.025 mole) of 4-dibenzothiophenol¹ in 100 ml. of chloroform. The resulting dark brown solution was allowed to stand for 12 hours. The mixture was then refluxed until gases were no longer evolved. On cooling, 1.5 g. (22.4%) of a light brown solid (m.p. 182-185°) separated. Two recrystallizations from methanol yielded 0.65 g. of material melting at 195-196°.

Anal. Calcd. for C₁₂H₇OClS: Cl, 15.02. Found: Cl, 14.91.

This analysis checks for a monochlorinated 4-dibenzothiophenol. The position of the chlorine was not determined, but it may be 1-chloro-4-dibenzothiophenol. This structure is suggested on the basis that bromination of 4-acetaminodibenzothiophene or of the closely related 4-hydroxydibenzofuran, or 4-methoxydibenzofuran² or 4-acetaminodibenzofuran³ yields an isomer-free product with the halogen atom in the position ortho to the biphenyl bridge and para to the other substituent.

(1) H. Gilman and A. L. Jacoby, *J. Org. Chem.*, **3**, 108 (1938).

(2) H. Gilman and P. R. Van Ess, *THIS JOURNAL*, **61**, 1365 (1939).

DEPARTMENT OF CHEMISTRY
IOWA STATE COLLEGE
AMES, IOWA

HENRY GILMAN
KENNETH E. LENTZ

RECEIVED AUGUST 3, 1951

3-Hydroxy-2-naphthalenemethanol

A solution of 9 g. (0.24 mole) of lithium aluminum hydride in 280 ml. of dry ether was placed in a 1-l., 3-necked flask

equipped with a dropping funnel, a mechanical stirrer, and a reflux condenser, and protected from moisture by calcium chloride tubes. A solution of 37 g. (0.197 mole) of 3-hydroxy-2-naphthoic acid in 675 ml. of dry ether was added at such a rate that the ether refluxed gently. Fifteen minutes after the last addition, water was added cautiously with cooling and stirring to decompose the excess lithium aluminum hydride. This was followed by the addition of 300 ml. of 10% sulfuric acid with cooling. The orange-red mixture was extracted with ether. The ether extracts were washed with water and 5% sodium bicarbonate solution and then dried over barium oxide. Evaporation of the ether gave 10 g. of material melting at 178-185°. After recrystallization from alcohol there was obtained 8 g. (23%) of 3-hydroxy-2-naphthalenemethanol, m.p. 190-191°.

Anal. Calcd. for C₁₁H₁₀O₂: C, 75.85; H, 5.79. Found: C, 75.64; H, 5.66.

CHEMISTRY LABORATORY
STATE UNIVERSITY OF IOWA
IOWA CITY, IOWA

WALTER T. SMITH, JR.
LOUIS CAMPANARO

RECEIVED NOVEMBER 8, 1951

Bromo-2-chlorokojic Acid¹

2-Chlorokojic acid (2-chloromethyl-5-hydroxy-4-pyrone) was found to participate in the Wohl-Ziegler reaction when 10 g. of the acid was refluxed with 12 g. of N-bromosuccinimide in 100 ml. of benzene for one hour.

At the end of the reaction period the whole mass was placed in a crystallizing dish and the benzene was removed by surface evaporation. The residual mixture was treated with 50 ml. of distilled water and filtered. The red crystalline product was recrystallized twice from absolute ethanol. The yield was 6.2 g.; the melting point of the white crystals obtained by sublimation was 163°. The compound was exceedingly sensitive to traces of iron salts, giving an intense red coloration.

*Anal.*² Calcd.: C, 30.07; H, 1.67. Found: C, 29.78; H, 1.75.

The compound is probably 2-chloromethyl-5-hydroxy-6-bromo-4-pyrone. When the same reaction as described above was tried with kojic acid only tars resulted.

DEPARTMENT OF CHEMISTRY
TEXAS SOUTHERN UNIVERSITY
HOUSTON 4, TEXAS

L. L. WOODS

RECEIVED SEPTEMBER 28, 1951

(1) Acknowledgment: The author is indebted to the Corn Products Company for the kojic acid used in this experiment and to the Research Corporation for a grant-in-aid on this project.

(2) Analyses by Dr. Carl Tiedcke.