

The m.p. of either product fell about 10° when the sample was allowed to stand in a tightly stoppered vial in the refrigerator. Repeated recrystallization from ethanol or methanol caused rearrangement to the ketone (VI).

Rearrangement of VII.—When 0.500 g. (0.00175 mole) of *trans*-V was brominated in methanol containing 0.30 g. of suspended sodium bicarbonate at -35° and then to the cold solution was added 3.00 g. of freshly prepared silver oxide, an oil was obtained which crystallized from petroleum ether to give 0.42 g. (69%) of *p*-methoxybenzhydryl phenyl ketone dimethyl ketal (VIII), which on further recrystallization from ethanol melted at 77.4–78°.

Anal. Calcd. for C₂₃H₂₄O₄: C, 79.3; H, 6.9. Found: C, 79.5; H, 7.0.

When the same conditions were used for *cis*-V, a 66% yield of VIII was obtained as shown by m.p. and mixed m.p.


The structure of VIII was established by its hydrolysis with 2 drops of 10% hydrochloric acid in 5 cc. of 95% ethanol to VI in 70% yield. It also gave VI after standing at room temperature for one month.

DEPARTMENT OF CHEMISTRY
COLUMBIA UNIVERSITY
NEW YORK 27, NEW YORK

Some Observations on Phosphinic Anhydrides

BY GENNADY M. KOSOLAPOFF
RECEIVED MAY 28, 1952

Some time ago we reported the synthesis of several dialkylphosphinic anhydrides from alkyl dialkylphosphinates and dialkylphosphinyl chlorides, as¹



At that time we also reported that the reaction of dimethylphosphinyl chloride with ethyl di-*n*-butylphosphinate gave not the expected mixed anhydride but a mixture of the two symmetric anhydrides.

In order to explore the possibly general nature of this observation we attempted the preparation of several mixed phosphinic anhydrides by this reaction. Equimolar mixtures of chlorides and ethyl esters of dimethyl-, diethyl-, di-*n*-propyl- and di-*n*-butylphosphinic acids were heated as described previously¹; substantially the theoretical amounts of ethyl chloride were obtained from all possible combinations of the above-mentioned compounds at 140–150°. In all instances, however, the distillation of the reaction mixtures, at 0.05–0.1 mm. pressure, gave only the corresponding pairs of the symmetric phosphinic anhydrides, which were characterized by comparison with the known specimens. The work was done in standard Pyrex apparatus. It is conceivable that the transformation of the expected unsymmetrical substances might have been catalyzed by the glass composition and might have been avoided by the use of fused quartz apparatus. However, it appears that the existence of unsymmetric phosphinic anhydrides at moderately elevated temperatures is purely transient. This points out a similarity of behavior to that of mixed carboxylic anhydrides and acetals, which tend to symmetrize during thermal treatment.

After this work was completed there appeared

(1) G. M. Kosolapoff and R. M. Watson, *THIS JOURNAL*, **73**, 4101 (1951).

the paper of Todd and co-workers² which reports the preparation of unsymmetrical tetraesters of pyrophosphoric acid. Although the unheated crude reaction mixtures indicate the presence of such unsymmetrical esters, the latter could not be isolated by distillation owing to symmetrization similar to the one reported here.

Phosphinic anhydrides are unique among the known anhydrides of various organic acids of phosphorus, in that they have definite boiling points and, therefore, they can be positively identified and purified by conventional techniques of organic chemistry. Tetraesters of pyrophosphoric acid, probably the most studied anhydrides at this time, on the other hand are prone to decomposition and rearrangement reactions at elevated temperatures which make impossible a true fractional distillation of such compounds. These esters are commonly purified by distillation in molecular stills in which boiling points, in the true sense of the word, have very little significance. Identification of individuals in a mixture of closely related substances under such circumstances becomes dubious.

As a result of our work and that reported by Todd, *et al.*,² it appears quite possible that at this time no positively characterized unsymmetrical neutral derivatives of phosphorus acid anhydrides are known. While such compounds have been undoubtedly obtained in low temperature reactions, no methods of isolation at low temperatures have been developed and all distillation techniques, used thus far, require thermal treatment which causes the symmetrization discussed above.³

(2) N. S. Corby, G. W. Kenner and A. R. Todd, *J. Chem. Soc.*, 1234 (1952).

(3) The material was presented at the 121st National Meeting of the A.C.S., March, 1952.

THE ROSS CHEMICAL LABORATORY
ALABAMA POLYTECHNIC INSTITUTE
AUBURN, ALA.

Some Salts of Symmetric Phosphinic Acids

BY WILLIAM C. DRINKARD AND GENNADY M. KOSOLAPOFF
RECEIVED MAY 28, 1952

Many phosphinic acids, *i.e.*, compounds of the general type R₂P(O)OH, are known at this time. Qualitative descriptions of a number of metallic salts of a few such acids appear in the literature. However, the only quantitative report dealing with solubility of such salts appears to be that dealing with the copper and lead salts of di-pseudocumylphosphinic acid.¹

Information about the solubilities of various salts of these acids is of obvious interest for the development of techniques of separation of mixtures of the acids themselves and of mixtures with other substances. We prepared, accordingly, several salts of four phosphinic acids. Two of the latter were aliphatic and two were aromatic, thus encompassing the relatively water-soluble and insoluble acids. Solubilities in water, benzene and 96% ethanol were determined at three temperatures (25, 35 and 45°) by conventional technique.

It is of interest to note the relatively high solu-

(1) A. Michaeils, *Ann.*, **204**, 1 (1896).