

the procedure described by Rietz, Chapman and Fernandez.¹

The tetradecyl derivative, m.p. 73°, was purified by crystallization from ethyl ether. *Anal.* Calcd. for C₂₂H₄₂S₄: C, 74.47; H, 12.71; S, 12.81. Found: C, 74.37; H, 12.69; S, 12.66.

The octadecyl derivative, m.p. 82°, was purified by crystallization from isopropyl ether. *Anal.* Calcd. for C₂₈H₅₄S₄: C, 76.52; H, 13.00; S, 10.49. Found: C, 76.53; H, 12.83; S, 10.59.

(1) Rietz, Chapman and Fernandez, *THIS JOURNAL*, **70**, 3486 (1948).

ORGANIC CHEMICAL LABORATORIES
UNIVERSITY OF FLORIDA
GAINESVILLE, FLA.

RICHARD D. CHAPMAN
JAMES B. FERNANDEZ
ROBERT MILLER
STANLEY D. MITCHEL
EDWARD G. RIETZ

RECEIVED JUNE 9, 1952

2,2-Dimethyl-5-dimethylammonium-1,3-benzodioxole Iodide and 2,2-Dimethyl-5-trimethylammonium-1,3-benzodioxole Iodide

A solution of 6.5 g. (0.033 mole) of 2,2-dimethyl-5-nitro-1,3-benzodioxole¹ in 200 ml. of absolute ethanol was hydrogenated in the presence of 1.5 g. of Raney nickel catalyst at 60° and 5 atmospheres of hydrogen for 3 hours. The resulting amine solution was separated from catalyst by filtration in an inert atmosphere, then evaporated under reduced pressure to half its volume, and mixed with 10 g. (0.07 mole) of methyl iodide. A precipitate which formed soon after addition of the methyl iodide was isolated after four days and washed with ethyl ether. The colorless precipitate was extracted with 30 ml. of hot chloroform, and the chloroform solution evaporated to dryness. The chloroform-insoluble fraction was recrystallized from absolute ethanol; yield 0.85 g., m.p. 264° (dec.). The chloroform-soluble fraction was purified in the same way; yield 1.5 g., m.p. 216° (dec.). The analysis of the higher-melting compound corresponded to that of 2,2-dimethyl-5-trimethylammonium-1,3-benzodioxole iodide. *Anal.* Calcd. for C₁₂H₁₈NO₂I: C, 43.0; H, 5.4; N, 4.2. Found: C, 42.8; H, 5.6; N, 3.9. The lower-melting compound gave an analysis corresponding to that calculated for 2,2-dimethyl-5-dimethylammonium-1,3-benzodioxole iodide. *Anal.* Calcd. for C₁₁H₁₆NO₂I: C, 41.1; H, 5.0; N, 4.4. Found: C, 41.3; H, 5.2; N, 4.1.

Additional amounts of both products were isolated from the alcoholic mother liquor of the reaction mixture by precipitation with ether.

Attempted conversion of the two compounds to the corresponding catechol polymethylammonium chlorides by treatment with a hot mixture of aqueous hydrochloric acid and silver chloride led to the isolation of glasses which could not be crystallized. These glasses, when made basic, were subject to air oxidation, which causes them to become highly colored.

(1) G. Slooff, *Rec. trav. chim.*, **54**, 995 (1935).

CHEMICAL CORPS
MEDICAL LABORATORIES
ARMY CHEMICAL CENTER, MARYLAND

DAVID H. ROSENBLATT
THOMAS PARR, III
JOSEPH EPSTEIN

RECEIVED JULY 7, 1952

The γ -Lactone of 3-Tridecyl-4-hydroxy-2-butenic Acid

1-Acetoxy-2-pentadecanone.—A 70% yield of a white waxy solid, b.p. 153–163° (2 mm.), m.p. 49–52°, was obtained by treating myristoyl chloride with diazomethane and the azide thus obtained with glacial acetic acid.

3-Tridecyl-4-hydroxy-2-butenic Acid, γ -Lactone.—An anhydrous benzene-ether solution of 1-acetoxy-2-pentadecanone (14.0 g.) reacted with granular zinc (3.53 g.) and ethyl bromoacetate (7.35 g.). The zinc complex was hydrolyzed and the intermediate thus obtained was dehydrated and lactonized by a saturated solution of anhydrous hydrogen bromide in glacial acetic acid. An oil, b.p. 185–195° (3–4 mm.), was obtained and recrystallization from alcohol and water gave a white solid, m.p. 57–58°. The product gave a positive Legal test, a negative potassium ferricyanide test, a positive Tollens test and it did not add bromine. The Legal test became negative after treatment

with alcoholic potassium hydroxide. The melting point of a mixture with myristic acid was lowered 5°.

Anal. Calcd. for C₁₇H₃₀O₂: C, 76.6; H, 11.4. Found: C, 75.6; H, 11.4.

1-Acetoxy-2-heptadecanone.—A 68% yield of a white waxy solid, m.p. 62–63°, was obtained from palmitoyl chloride as described above. The product gave a positive ketone test with 2,4-dinitrophenylhydrazine. The melting point of a mixture with palmitic acid was not depressed. Two attempts to prepare a lactone from this compound failed.

Anal. Calcd. for C₁₉H₃₆O₃: C, 73.0; H, 11.6. Found: C, 73.0; H, 11.6.

SCHOOL OF PHARMACY
PURDUE UNIVERSITY
LAFAYETTE, INDIANA

DAVID A. SCHLICHTING
GLENN L. JENKINS

RECEIVED JUNE 9, 1952

α -(Benzylmercaptomethyl)-succinic Anhydride

Dimethyl α -(Benzylmercaptomethyl)-succinate.—A solution of 31.6 g. (0.2 mole) of dimethyl itaconate in 50 cc. of absolute ethanol was treated with 27.2 g. (0.22 mole) of benzylmercaptan after the addition of 5 drops of piperidine. The mixture was refluxed for 10 hr., then the solvent was removed and the residue heated at ca. 110–120° for two hours before fractionation. A 78% yield of golden oil, b.p. 153–160° (0.7–0.8 mm.) was obtained; redistillation gave pure dimethyl α -(benzylmercaptomethyl)-succinate, b.p. 154–156° (0.7 mm.), n_D^{20} 1.5250.

Anal. Calcd. for C₁₄H₁₈O₄S: S, 11.35. Found: S, 11.30.

α -(Benzylmercaptomethyl)-succinic Acid.—A mixture of 30.5 g. (0.107 mole) of dimethyl α -(benzylmercaptomethyl)-succinate and a solution of 15.5 g. of 85% potassium hydroxide (0.235 mole) in 50 cc. of water was refluxed five hours. The solution was charcoaled, extracted with ether and the aqueous layer acidified, extracted with ether and dried. A creamy white solid (21.5 g., 79% yield) remained after the removal of solvent and cooling of the residue. The crude product (m.p. 104–106°) was crystallized twice from 3:2 hexane-benzene to yield the pure acid in the form of white blades, m.p. 108.5–109° (cor.).

Anal. Calcd. for C₁₂H₁₄O₄S: C, 56.67; H, 5.55; S, 12.61. Found: C, 56.49; H, 5.33; S, 12.65.

α -(Benzylmercaptomethyl)-succinic Anhydride.—There was refluxed 15.26 g. (0.6 mole) of α -(benzylmercaptomethyl)-succinic acid and 50 cc. of acetic anhydride for 4 hours, and then the excess acetic anhydride removed. The residue was distilled, and 11.8 g. (83% yield) of golden oil collected at 150–155° (0.2 mm.). The oil solidified after some standing and crystallized from benzene-cyclohexane as fine white needles, m.p. 58–58.5° (cor.).

Anal. Calcd. for C₁₂H₁₄O₃S: C, 61.00; H, 5.12; S, 13.57. Found: C, 60.70; H, 5.36; S, 13.32.

STERLING-WINTHROP RESEARCH INSTITUTE

RENSELAER, NEW YORK

EDGAR A. STECK

RECEIVED MAY 15, 1952

Ethyl β -Benzylmercaptopropionate

A solution of 12.0 g. (0.52 mole) of sodium in 150 cc. of absolute ethanol was treated (reflux, stirring, 0.5 hr.) with 64.5 g. (0.52 mole) of benzylmercaptan at 30–40°, then refluxed for an hour. To the sodium benzylmercaptide was added (one hour) a solution of 90.5 g. (0.5 mole) of ethyl β -bromopropionate in 75 cc. of absolute ethanol and the mixture refluxed three hours. The mixture was quenched in considerable water, extracted with ether, then the extracts washed (saturated sodium chloride solution) and dried over sodium sulfate. Fractionation of the extract gave 92.7 g. (87.5% yield) of ethyl β -benzylmercaptopropionate; b.p. 101–103° (0.22 mm.), n_D^{25} 1.5305.

Anal. Calcd. for C₁₁H₁₆O₂S: C, 62.23; H, 9.05; S, 15.10. Found: C, 62.29; H, 9.37; S, 14.88.

STERLING-WINTHROP RESEARCH INSTITUTE

RENSELAER, NEW YORK

EDGAR A. STECK

RECEIVED MAY 15, 1952

(1) Analyses by Mr. M. E. Auerbach and Miss P. Curran.