

pyridine and 100 ml. of acetic anhydride were added slowly, with stirring and slight cooling.

After several hours a clear solution, free of undissolved particles, resulted. After 20 hours at room temperature, the solution was added to a large volume of vigorously stirred ice and water and the dingy-looking precipitate was washed free of acid with water. A second acetylation treatment^{1,3} resulted in no change in acetyl content or in intrinsic viscosity. To provide comparison with the results of others, several variations of this procedure are indicated in Table I.

For acetylation with pyridine as catalyst at 100°, 5 g. of oven-dried polysaccharide (60 mesh) was stirred for 3 hours with 20 ml. of pyridine and 17 ml. of acetic anhydride. The homogeneous solution was cooled, poured into vigorously agitated ethanol, and the colorless precipitate was washed free of acid and dried in the air.

For acetylation with fused sodium acetate as catalyst at 100°, 5 g. of air-dried corn amylose (60 mesh) was soaked in 75 ml. of acetic anhydride overnight, then 1.5 g. of fused sodium acetate was added and the mixture stirred at 100° for 6 hours. The homogeneous solution was cooled, poured into vigorously stirred ice-water, and the colorless product was washed free of acid with water.

Solubility of the Triacetates.—Corn amylose triacetates prepared by the formamide or sodium acetate methods were soluble in acetone; that prepared by the pyridine method at 100° was not. All corn amylose triacetate preparations, as well as the triacetates from corn starch, were soluble in chloroform and 1,1,2-trichloroethane. Preparations of waxy corn starch triacetate from both the pyridine method at 100° and the formamide method were soluble in acetone as well as in the chlorinated solvents. Corn amylopectin triacetate prepared by the pyridine method at 100° was insoluble in the chlorinated solvents, but that prepared by the formamide method was readily soluble.

Acknowledgment.—We wish to acknowledge our indebtedness to Dr. G. E. Hilbert for his interest in the early phases of this work, and to C. H. Van Etten and R. Tobin for the acetyl determinations.

NORTHERN REGIONAL RESEARCH LABORATORY⁸
PEORIA, ILLINOIS

(8) One of the laboratories of the Bureau of Agricultural and Industrial Chemistry, Agricultural Research Administration, U. S. Department of Agriculture. Article not copyrighted.

Alkylation of 2-Naphthol by Esters of Aliphatic Acids

BY SAUL PATAI AND M. BENTOV

RECEIVED JULY 17, 1952

In the course of a study on the preparation of aromatic methacrylates¹ the transesterification of methyl methacrylate with 2-naphthol in the presence of sulfuric acid was investigated. Instead of the expected ester, 2-methoxynaphthalene was formed. In order to elucidate this reaction, the transesterification of other aliphatic esters with other phenolic compounds was tried, using sulfuric acid, orthophosphoric acid or metallic sodium as the catalyst. Phenol, 1-naphthol, 4-hydroxybiphenyl and 4-nitrophenol did not react with methyl methacrylate or ethyl acetate. 2-Naphthol, on the other hand, reacted readily in the presence of sulfuric acid, but not of phosphoric acid or sodium, with methyl methacrylate, methyl acetate, ethyl acetate and ethyl butyrate, yielding 2-methoxynaphthalene with the methyl esters and 2-ethoxynaphthalene with the ethyl esters, but failed to

(1) S. Patai, M. Bentov and M. E. Reichmann, *THIS JOURNAL*, **74**, 845 (1952).

react with ethyl formate or dimethyl oxalate. With methyl benzoate the normal product of transesterification, 2-naphthyl benzoate, was obtained.²

2-Naphthol is known to be one of the very few phenols which can be alkylated by alcohols in the presence of acids.³ This fact may be related to the tautomerizability of 2-naphthol.⁴

As, however, our experiments were carried out under strictly anhydrous conditions, no alcohol could be formed as an intermediate in the reaction. It must, therefore, be assumed, that the alkylation is caused in both cases by the hydrogen alkyl sulfate, formed in the conditions of our experiments from the ester and sulfuric acid. The dissociation constant of formic acid is about ten times, and that of oxalic acid about a thousand times greater than that of either acetic, butyric or methacrylic acid: the failure of the reaction with the esters of formic and oxalic acid could then be due to the strength of these acids, which prevents the formation of the alkylsulfuric acids by interaction of the esters with sulfuric acid.

Acknowledgment.—This investigation was carried out under the auspices of the Scientific Department, Israeli Ministry of Defence, and is published with its permission. The authors wish to thank Dr. E. D. Bergmann for helpful advice and discussions.

Experimental

A mixture of 0.04 mole of 2-naphthol, 0.1 ml. of concentrated sulfuric acid and 0.08 mole of the ester was refluxed for four hours with the exclusion of moisture. (In the case of methyl methacrylate, some hydroquinone was added to the reaction mixture.) The product was poured into an excess of 5% aqueous sodium hydroxide solution and cooled for one hour in an ice-bath, and the resulting precipitate filtered and recrystallized from ethanol. 2-Methoxy- and 2-ethoxynaphthalene were identified by melting point (72 and 37°, respectively) and mixed melting point with authentic samples. Yield of 2-methoxynaphthalene from methyl acetate 20–25%; from methyl methacrylate 45–50%; of 2-ethoxynaphthalene from ethyl acetate 40–45%; from ethyl butyrate 20–25%.

(2) C. A. Bischoff and A. Hedenstrom (*Ber.*, **35**, 3449 (1902)) obtained di-2-naphthyl oxalate from diphenyl oxalate and 2-naphthol, and report the failure of the reaction with diethyl oxalate.

(3) C. Liebermann and A. Hagen, *ibid.*, **15**, 1427 (1882); L. Gattermann, *Ann.*, **244**, 72 (1888); W. A. Davis, *J. Chem. Soc.*, **77**, 33 (1900); L. F. Fieser and W. C. Lothrop, *THIS JOURNAL*, **57**, 1459 (1935).

(4) N. N. Voroshcov, *Bull. Inst. Polyt. Ivanovo-Voznesensk*, **6**, 125 (1922); C. A., **17**, 1637 (1923); J. C. Salfeld, *Ber.*, **73**, 376 (1940).

DEPARTMENT OF ORGANIC CHEMISTRY
THE HEBREW UNIVERSITY AND
LABORATORIES OF THE SCIENTIFIC DEPARTMENT
ISRAELI MINISTRY OF DEFENCE
JERUSALEM, ISRAEL

Cohumulone, a New Hop Constituent

BY F. L. RIGBY AND J. L. BETHUNE

RECEIVED JULY 16, 1952

In the course of an investigation to develop a quantitative method for the estimation of the isohumulone content of beer, petroleum ether extracts of several beers were distributed in a 100-tube all-glass countercurrent apparatus using a solvent system of 2,2,4-trimethylpentane and a phosphate-citrate buffer pH 4.9. The concentration in each