

[CONTRIBUTION FROM THE RADIATION LABORATORY AND DEPARTMENT OF CHEMISTRY, UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA]

Cyanide Effects on Carbon Dioxide Fixation in *Chlorella*¹

BY B. R. RABIN,² D. F. SHAW,³ N. G. PON, J. M. ANDERSON⁴ AND M. CALVIN

RECEIVED OCTOBER 15, 1957

Green algae have been treated with radioactive KCN in an investigation of the action of cyanide on photosynthesis. A multitude of products have been found to be formed in very short exposures (10-15 sec.). One of these products has been identified with the product formed when the algae are given radioactive CO₂ and non-radioactive KCN. The same product has been synthesized by a non-enzymatic cyanohydrin addition reaction on ribulose 1,5-diphosphate. It has been shown to be a 2-carboxy-pentitol (probably mostly ribitol) 1,5-diphosphate. Upon hydrolysis it gives an hydroxy acid (or mixture of isomers) closely related to hamamelonic acid. The significance of this and the other as yet unidentified products of cyanide interaction with a biological system is discussed with respect to the use of cyanide as an inhibitor.

Introduction

The effects of cyanide on biochemical systems, both *in vivo* and *in vitro*, have been studied under a large variety of conditions. Thus, the use of cyanide as an inhibitor for enzymes such as tyrosinase⁵ and catalase⁶ is well known. The mode of action appears to be due to metal inactivation.⁷ The possibility of cyanide acting as a carbonyl trap by cyanohydrin formation has been suggested in a limited way.⁸

Applications of cyanide to the problem of photosynthesis are also quite numerous. For example, in 1919 Warburg⁹ found that HCN administered at very low concentrations, decreased the rate of photosynthesis (O₂ evolution) and increased the rate of respiration in *Chlorella*. The experiment was done in the presence of strong light and abundant carbon dioxide. A similar result was observed by Ruben, *et al.*,¹⁰ when cyanide was added to *Chlorella* at high concentrations (10⁻² M KCN); both the photosynthesis rate (CO₂ uptake) and the carbon dioxide dark fixation rate were reduced to 0.3% of its normal value, but its respiration remained essentially unchanged. On the other hand, Gaffron¹¹ noted that cyanide affected *Scenedesmus* D1 in the opposite manner. Thus, an addition of 2 × 10⁻⁴ M KCN inhibited respiration nearly completely while photosynthesis (O₂ evolution) remained untouched. In a later work¹² it was found that the cyanide-suppressed CO₂ dark fixation was a function of the preillumination period. That is, the longer the KCN is allowed to stand in the light with the plant, before the introduction of C¹⁴O₂ and darkening, the greater the inhibition of CO₂ dark

fixation. A marked effect on the C¹⁴-content of the water-soluble fraction also was observed. Meanwhile, in this Laboratory, we¹³ had examined the effect of cyanide on C¹⁴O₂ fixation by *Scenedesmus* (Gaffron D-3 strain) during photosynthesis. The algae were preilluminated aerobically without CO₂ for 30 minutes. Cyanide (3 × 10⁻⁴ M) was added 1 minute before the addition of C¹⁴O₂, followed by 1 minute photosynthesis. A 95% inhibition of C¹⁴O₂ fixation was found. It is interesting to note that the ratio of 3-phosphoglyceric acid to hexose and triose phosphates varied from 1:4.5 under non-inhibited conditions to 3.2:1 under cyanide inhibited conditions. The conclusion from this experiment was that cyanide, under these conditions, does not prevent the initial carboxylation step with the subsequent formation of phosphoglyceric acid. Related to this work is the cyanide inhibition experiment on the *in vitro* system, carboxydismutase.¹⁴ CO₂ fixation appeared to be completely suppressed at 10⁻² M KCN while lower concentrations (10⁻³ M and 10⁻⁴ M) yielded about 50% and 2% inhibition, respectively.

An extension of the examination of the effect of CN⁻ on the pattern of CO₂ fixation in the alcohol-water soluble fractions of *Chlorella* was made in the early part of 1957 by O. Kandler,¹⁵ in this Laboratory. The algae were subjected to normal photosynthesis (10 sec.) with C¹⁴O₂. To the algal suspension was added KCN (final concentration M/80) followed by a 10 second light period and finally hot alcohol was injected (to a final concentration of 80%) to stop all of the enzyme reactions. The essential features of these experiments were the large decrease of C¹⁴ activity in phosphoglyceric acid and concurrent increase of C¹⁴ activity in the diphosphates. The latter, when treated with acid phosphatase, gave two new spots plus other known sugars. The new spots were characterized by means of paper electrophoresis and rechromatography. These spots were found to exhibit an acid-lactone behavior.

It is the purpose of this paper to establish in more detail the role of cyanide in the formation of this new compound in *Chlorella*. Beyond this it seemed useful to examine the fate of C¹⁴N⁻ in bio-

(1) The work described in this paper was sponsored in part by the United States Atomic Energy Commission and in part by the Chemistry Department, University of California, Berkeley, California.

(2) Rockefeller Foundation Fellow, 1956-1957. On leave from the Department of Biochemistry, University College, London, W.C. 1.

(3) Commonwealth Fund Fellow, 1956-1957. Imperial Chemical Industries, Ltd., Billingham, England.

(4) King George VI Memorial Fellow for New Zealand, 1956-1957.

(5) L. E. Tenenbaum and H. Jensen, *J. Biol. Chem.*, **147**, 27 (1943).

(6) K. Zeile and H. Hellström, *Z. physiol. Chem.*, **192**, 171 (1930).

(7) J. B. Neilands, P. K. Stumpf and R. Y. Stanier, "Outlines of Enzyme Chemistry," John Wiley and Sons, Inc., New York, N. Y., 1955, p. 91.

(8) E. A. Zeller, "The Enzymes," edited by J. B. Sumner and K. Myrback, Vol. II, part 1, Academic Press, Inc., New York, N. Y., 1951, p. 552.

(9) O. Warburg, *Biochem. Z.*, **100**, 230 (1919).

(10) S. Ruben, M. D. Kamen and W. Z. Hassid, *THIS JOURNAL*, **62**, 3443 (1940).

(11) H. Gaffron, *Biol. Zentr.*, **59**, 302 (1939).

(12) H. Gaffron, E. W. Fager and J. L. Rosenberg, *Symposia Soc. Exp. Biol.*, **5**, 262 (1951).

(13) M. Calvin, J. A. Bassham, A. A. Benson, V. H. Lynch, C. Ouellet, L. Schou, W. Stepka and N. E. Tolbert, *ibid.*, **5**, 284 (1951).

(14) H. L. Kornberg, J. R. Quayle and M. Calvin, University of California, Radiation Laboratory Report, UCRL 2885, May (1955).

(15) O. Kandler, University of California Laboratory Report, UCRL 3710, p. 8, March, 1957; *Arch. Biochem. Biophys.*, in press.



Fig. 4.—Radioautograph: chromatography of the nonenzymatic reaction product of $KC^{14}N$ and RuDP.




Fig. 5.—Radioautograph of chromatogram of ribulose diphosphate and hamamelonic acid diphosphate ($KC^{14}N$) in butanol propionic acid (48 hours). The circles are the sprayed areas of non-radioactive RuDP.





Fig. 6.—Co-chromatography of authentic hamamelonic acid lactone with the HCl hydrolysate of the nonenzymatic reaction product of $KC^{14}N$ and RuDP. (Note origin at lower left.) The circle on the paper (right) corresponds to the radioactivity of film (left).




Fig. 7.—Radioautograph of a chromatogram of the hot alcohol-water extract of *Chlorella* which were treated as follows: Normal photosynthesis for 2 min. with $C^{14}O_2$ followed by addition of KCN (final concentration 0.02 M) along with a further 15 sec. light period and injection of hot alcohol (end concentration 80%) to kill the algae. ($C^{14}O_2$, CN^- expt.).



Fig. 8.—Radioautograph of a chromatogram of the hot alcohol-water extract of *Chlorella* which were treated as follows: Normal photosynthesis for 2 min. with CO_2 followed by addition of $KC^{14}N$ (final conc. 0.02 M) along with a further 15 sec. light period and finally, injection of hot alcohol (end concentration 80%) to kill the algae. (CO_2 , $C^{14}N^-$ expt.).




Fig. 9.—Radioautograph: chromatography of the enzymically hydrolyzed diphosphate area of Fig. 7 ($C^{14}O_2$, CN^- expt.).



Fig. 10.—Radioautograph: chromatography of the enzymically hydrolyzed diphosphate area of Fig. 8 (CO_2 , $C^{14}N^-$ expt.).




Fig. 11.—Radioautograph: Chromatography of the HCl hydrolyzed diphosphate area of Fig. 7 ($C^{14}O_2$, CN^- expt.).




Fig. 12.—Radioautograph: Chromatography of the HCl hydrolyzed diphosphate area of Fig. 8 (CO_2 , $C^{14}N^-$ expt.).




Fig. 13.—Radioautograph: Parallel paper electrophoresis of the diphosphate area of Fig. 7 ($C^{14}O_2$, CN^- expt.) and the nonenzymic reaction product of $KC^{14}N$ and RuDP.

logical systems in order to try and understand its widespread biological effects.

Experimental

Photosynthesis Experiment.—The cultures of *Chlorella pyrenoidosa* used in the photosynthesis experiments were grown in the medium described by Holm-Hansen, *et al.*¹⁶ using a continuous culture apparatus with constant fluorescent lighting of approximately 1,000 f.c. After harvesting, the algae were centrifuged, washed and resuspended in distilled water to give a 1% suspension. Aliquots of 10 ml. were used in the photosynthesis experiments, which were performed in the vessel illustrated in Fig. 1.

Illumination was provided by reflector spotlight which gave a light intensity of approximately 7,000 f.c. on either side of the vessel; excessive heat production was avoided by using water-cooled infrared filters.

The algae were preilluminated for 20 minutes; a continuous stream of 1% CO₂ in air was bubbled through during this period. At the commencement of the experiment the air stream was shut off and sodium bicarbonate (1 ml. of 0.0013 M) was added. After 2 minutes photosynthesis with occasional stirring, KCN (1 ml. of 0.24 M) was added, and after a further 15 seconds the algae were killed by the addition of boiling ethanol (40 ml.) to produce an 80% alcoholic mixture. The experiments were of two types: (1) using a combination of NaHC¹⁴O₃ (20 μc./ml.) and KCN designated as C¹⁴O₂, CN⁻; (2) using a combination of NaHCO₃ and KC¹⁴N (3.6 mc./ml.)¹⁷; designated as CO₂, C¹⁴N⁻. Extracts for chromatography were prepared by centrifuging the mixture; the supernatant liquid was retained and the residue was extracted successively with 10 ml. of 20% ethanol and 10 ml. of boiling water. All the extracts were combined, concentrated to approximately 3 ml. in a rotating evaporator¹⁸ and chromatographed in two dimensions (phenol-water first dimension, butanol-propionic acid-water second dimension).¹⁹ The diphosphate was eluted with water and either used as such, or hydrolyzed. Two methods of hydrolysis were employed: (1) HCl (1 N) hydrolysis in a sealed tube at 120° for 16 hr. and (2) enzymic hydrolysis at pH 5.0 (0.2 M in acetate buffer and 0.01 M Mg⁺⁺) using purified Polidase-S²⁰ (Schwarz Laboratories, Inc.) or acid prostatic phosphatase.²¹

Chromatography and Electrophoresis.—Whatman No. 4 paper was used throughout, except where otherwise noted. For routine chromatography the paper was washed according to Benson¹⁹; for chromatography in ethanol-water-formic acid-sodium molybdate, it was washed according to Trudinger.²² (In the latter case Whatman No. 1 was used.) Unwashed paper was used for electrophoresis and for chromatograms which were subsequently sprayed with Tollens reagent. (AgNO₃ dissolved in a little water, concentrated NH₄OH added until the precipitate dissolves, then diluted with methanol to 5% AgNO₃ w./v.) The apparatus routinely used in this Laboratory for chromatography was employed. Electrophoresis was carried out in an apparatus similar to that described by A. B. Foster.²³ Glycine buffer (0.1 M) pH 9.75 was used and the experiment was run at 600 v. (10 to 20 ma.) for 3 hr. at room temperature. For chromatography with carrier hamamelonic acid diphosphate, the ethanol-water-formic acid-sodium molybdate system was used.

Phosphate esters were detected on the paper by spraying with the reagent described by Hanes and Isherwood,²⁴ dry-




Fig. 1.—Glass vessel for KCN-algae work.

ing in an oven with a forced air draft at about 60° and exposing to sunlight. Free sugar and hamamelonic acid were detected by spraying the paper, after thorough steaming, with Tollens reagent. The paper was washed with 5% ammonia to remove the excess of silver salts and to reduce the background color. Both hamamelonic acid and the lactone react poorly with the spray and prolonged heating (several minutes at 100°) is required for satisfactory results.

Radioactive areas on paper chromatograms were detected by exposing the paper to du Pont X-ray film type 507. For plate counting, Mylar²⁵ end-window, Q gas G.M. tubes were employed; the planchets were prepared in 2 N acetic acid and dried under an infrared lamp.

Preparation and Assay of Phosphoriboisomerase-Phosphoribulokinase Mixture.—The preparative steps are illustrated in the flow diagram, Fig. 2.^{26,27} It is essential that all operations are carried out at 0°.

The isomerase-kinase mixture was assayed by measuring the rate of acid production using a pH-stat.²⁸ The test solution contained 11 μmoles of adenosine triphosphate (ATP),²⁹ 10 μmoles of ribose 5-phosphate, 10 μmoles MgCl₂ and 5 μmoles of cysteine in 10 ml. The pH was adjusted to 7.40 and held constant during the reaction by addition of 0.1 M NaOH. The reaction was started by the addition of the isomerase-kinase mixture (150 μl.). In a typical assay 7.14 μmoles of NaOH was consumed in 25 minutes; the alkali uptake slowed down considerably after 10 minutes. A control without ribose 5-phosphate showed that the preparation contained very little ATPase activity (alkali consumption ceased after 10 minutes; a total of 1.5 μmoles was consumed).

Preparation of RuDP.—The method depends on the conversion of RMP to RuDP in the presence of ATP and the isomerase-kinase mixture. The barium salt of ribose 5-phosphate (4 mmoles, 1.83 g.) was dissolved in water (40 ml.) and potassium sulfate (40 ml. of 0.15 M) added. The mixture was centrifuged and the residue washed with water (20 ml.). To the combined washings and supernatant liquid 6 mmoles of ATP, 0.4 mmole of cysteine and 4 mmoles of MgCl₂ were added and the volume made up to 300 ml. The reaction was carried out in the pH-stat at room temperature under an atmosphere of N₂ gas; the pH was adjusted to 7.40 and held constant by the addition of CO₂-free NaOH (1.8 N). The reaction was started by the ad-

(16) O. Holm-Hansen, P. Hayes and P. Smith, University of California Radiation Laboratory Report, UCRL 3595, p. 56, October 1956.

(17) R. Lemmon, B. Fingerma and H. Simon, University of California Radiation Laboratory Report, UCRL 3836, p. 18, June, 1957. It was shown that the amount of C¹⁴O₂ present in KC¹⁴N could not be greater than 0.001%.

(18) Rinco Instrument Co., Greenville, Illinois.

(19) A. A. Benson "Moderne Methoden der Pflanzenanalyse" II, edited by K. Paech and M. V. Tracey, Springer Verlag, Berlin, 1955, p. 113.

(20) S. S. Cohen, *J. Biol. Chem.*, **201**, 71 (1953).

(21) G. Schmidt, "Methods of Enzymology," edited by S. P. Colowick and N. O. Kaplan, Vol. II, Academic Press, Inc., New York, N. Y., 1955, p. 523.

(22) P. A. Trudinger, *Biochem. J.*, **64**, 274 (1956).

(23) A. B. Foster, *Chem. & Ind.*, 1050 (1952).

(24) C. S. Hanes and F. A. Isherwood, *Nature*, **164**, 1107 (1949).

(25) Mylar is du Pont's registered trademark for its polyester film. P. M. Hayes, University of California Radiation Laboratory Report, UCRL 2932, March, 1955, p. 48.

(26) B. L. Horecker, J. Hurwitz and A. Weissbach, *J. Biol. Chem.*, **218**, 785 (1956).

(27) J. R. Quayle, University of California Radiation Laboratory Report, UCRL 3017, January, 1956.

(28) J. B. Neilands and M. D. Cannon, *Anal. Chem.*, **27**, 29 (1955).

(29) The following abbreviations will be used throughout this paper: ATP, adenosine triphosphate; RMP, ribose 5-phosphate; RuDP, ribulose 1,5-diphosphate; PGA, phosphoglyceric acid; HmDP, hamamelonic acid diphosphate; RuMP, ribulose 5-phosphate.




Fig. 2.—Flow diagram for the preparation of phosphoribulose-1,5-bisphosphate.

^a All centrifugations were done in a Spinco Model L, ultracentrifuge. ^b After each addition of ammonium sulfate the solution was adjusted to pH 7.0 with 2 N NH₄OH.

dition of the isomerase-kinase preparation (5 ml.) and stopped, after 37 minutes when the uptake of alkali (1.5 ml.) had slowed down, by the addition of 20 ml. of 50% trichloroacetic acid. The nucleotides²⁸ were removed by the addition of acid-washed Norite A (120 g.). The solution was centrifuged (2000 r.p.m. for 10 minutes) and the residue washed successively with four 200 ml. increments of water. Barium acetate (4 ml. of 1 M) was added to the combined supernatant liquid and washings and the pH adjusted to 6.4 with saturated barium hydroxide. The precipitate was removed by centrifugation and washed with water (50 ml.). The supernatant liquid and washings were combined and an equal volume of ethanol was added. RuDP precipitates in a gelatinous form; it was collected by centrifugation, washed with 80% ethanol and dried *in vacuo* over P₂O₅; yield 1.75 g. (crude).

Analysis of the preparation showed it to contain 0.4 μ mole of inorganic phosphate and 2.31 μ moles of organic phosphate per mg. These analytical figures must be regarded as approximate only. Assuming the only phosphate ester present to be the dibarium salt of RuDP, the product was

69% pure. In the experiments with cyanide, this preparation was used without further purification.

Purity of RuDP as Determined by Column Chromatography.—The column employed was 0.8 cm. (diameter) \times 27 cm. The resin was Dowex 1 formate; it was freed from fines by decantation and converted from the chloride to the formate form with 2 N formic acid.

The elution apparatus consisted of two vessels of equal size and shape. One, the mixing vessel, was provided with a stirrer and had an outlet to the top of the column; water (100 ml.) was placed in this vessel. Ammonium formate (100 ml., 2 M) was placed in the other vessel and the two vessels connected together by means of a siphon tube. This system provides a linear gradient of ammonium formate at the outlet of the mixing vessel.

RuDP (100 mg. Ba salt) was dissolved in water (30 ml.) with the aid of a little Dowex 50 (H⁺ form), and the pH adjusted to 6.0 with KOH. The solution was centrifuged and run through the column. The column was washed with water and subjected to gradient elution; fractions were collected every 5 minutes using an automatic device. The flow rate was 0.44 ml./min. The fractions found to contain phosphorus by qualitative test²⁴ were assayed for total phosphate³⁰ and RuDP. The method used for the determination of RuDP was based on the fixation of C¹⁴O₂ by RuDP in the presence of carboxydismutase. The reagent solution was prepared by mixing NaHC¹⁴O₃ (200 μ l. of 0.036 N, 400 μ c./ml.), MgCl₂ (80 μ l. of 0.01 M in 0.04 N HCl), tris³¹ buffer (20 μ l., pH 7.83, 1 M with respect to tris) and carboxydismutase (50 μ l. of a spinach preparation fractionated between 35 and 39% saturated ammonium sulfate and containing 6 mg. of protein per ml.). An aliquot from each column fraction (5 μ l.) was placed on a planchet and the reagent solution (15 μ l.) added. The planchet was placed on moist filter paper in a covered vessel, to reduce evaporation, and allowed to incubate at room temperature. After 2 hr. the reaction was stopped by the addition of acetic acid (10 μ l. of 6 N) and the planchet prepared for counting in the usual manner. The elution pattern obtained is shown in Fig. 3.

Two well separated phosphorus-containing peaks are observed. The first to emerge contains inorganic phosphate;




Fig. 3.—Elution pattern of RuDP: Klett reading of 230 corresponds to 0.5 μ mole phosphate.

(30) R. J. L. Allen, *Biochem. J.*, **34**, 858 (1940).

(31) Tris-(hydroxymethyl)-amino-methane.

the second peak is in the position expected for a diphosphate ester, and the enzyme assay shows it to contain essentially all the RuDP emerging from the column. It is noteworthy that the RuDP and phosphate peaks do not coincide in the second peak, suggesting the presence of at least one phosphate ester in addition to RuDP; this impurity emerges from the column slightly ahead of RuDP. From the ratio RuDP/organic phosphate, the quantity of this contaminant can be calculated. This constitutes the shaded area on the graph, Fig. 3, and represents about 20% of the material in the second peak. The fractions emerging between A and B (sample a) and B and C (sample b) were bulked. To each of them barium acetate (500 μ l. of 1 *M*) and an equal volume of 95% ethanol were added. The solutions were cooled in ice and the precipitate collected by centrifugation. The precipitates were washed with 50% ethanol (20 ml.), 95% ethanol (10 ml.), absolute ethanol (10 ml.), and dried *in vacuo* over P_2O_5 ; material recovered: sample a, 22 mg.; sample b, 23 mg.

Both samples contained a small quantity of inorganic phosphate, though very much less than the starting material. By treatment with acid phosphatase the only sugar found to be present in sample b was ribulose; sample a appeared to contain a small quantity of ribose in addition to ribulose. Sample b is considered to be a highly purified dibarium salt of ribulose diphosphate.

Reaction of RuDP with $KC^{14}N$.—RuDP (12 mg. Ba salt, crude) was triturated with water (1 ml.) and Na_2SO_4 (40 μ mole). The residue of barium sulfate was removed by centrifugation and washed with water (200 μ l.). The combined supernatant liquid and washings were allowed to react with 1.3 mg. of $KC^{14}N$ (1 μ c./ μ mole) at room temperature for 16 hr. An aliquot was plated in 6 *N* acetic acid and counted; 21,000 d./m./ μ l. were fixed as an acid stable compound.³² This represents approximately 100% reaction, when allowance for the impurities present in RuDP are taken into account.

Two-dimensional chromatography in the usual solvent system showed 87% of the activity was in the diphosphate area, as shown in Fig. 4. One dimensional chromatography of the unhydrolyzed reaction product in butanol-propionic acid-water for 48 hr. showed the presence of one main spot, with the same R_f value as HmDP with a small amount of other impurities as shown in Fig. 5.

An aliquot of the above material (25 μ l.) plus 100 μ g. of carrier authentic hamamelonic acid³³ was hydrolyzed with 1 *N* HCl by heating for 16 hr. in a sealed tube at 120°. The resulting solution was chromatographed in two dimensions (butanol-propionic acid-water followed by phenol-water) and is shown in Fig. 6. It was found that the radioactivity and the silver spray did not coincide exactly. This is almost certainly due to the probability that the radioactive material consists of both epimers at the α -carbon atoms. A similar spot in the lactone position was produced by the action of acid phosphatase on the material; this indicates that the reaction product contains a carboxyl group and is not cyanohydrin, which is presumably formed first, but is subsequently hydrolyzed.

Preparation of Hamamelonic Acid Diphosphate (2-Carboxyribitol 1,5-Diphosphate).—RuDP (117 mg. Ba salt, crude) was triturated with water (10 ml.) containing K_2SO_4 (75 mg.). The insoluble material was removed by centrifugation and KCN (900 μ l. of 0.24 *M*) added. After 18 hr. at room temperature the solution was centrifuged and the small amount of residue discarded. The supernatant liquid was cooled in ice and barium acetate (600 μ l. of 1 *M*) added. The copious precipitate was centrifuged off, washed with 5-ml. portions of 80 and 90% ethanol and dried *in vacuo* over P_2O_5 ; yield 120 mg. A further 15 mg. of hamamelonic acid diphosphate Ba salt could be recovered by the addition of ethanol (2 ml.) to the supernatant liquid. The material (120 mg. plus 15 mg.) was contaminated with $BaSO_4$ which was removed by trituration with HCl (10 ml. of 0.1 *N*) and centrifuged. The residue was washed with

(32) The symbol HmDP is used in this manuscript to designate the product of the addition of KCN to RuDP. It is derived from the name hamamelonic acid diphosphate. As will appear later the compound upon hydrolysis produces a mixture of stereoisomeric hydroxy acids including hamamelonic acid.

(33) We are grateful to Prof. O. Th. Schmidt for a gift of several mg. of that stereoisomer which he has established to be hamamelonic acid; O. Th. Schmidt and K. Heintz, *Ann.*, **515**, 77 (1935).

water (5 ml.); it weighed 42 mg. and consisted mainly of $BaSO_4$.

The solution was adjusted to pH 7.1 and cooled in an ice-bath. The precipitate of hamamelonic acid diphosphate, Ba salt, centrifuged off, washed with 50% ethanol (7.5 ml.), 90% ethanol (5 ml.) and dried *in vacuo* over P_2O_5 , yield 26 mg. (A). A further crop was obtained by adding equal volumes of ethanol to the supernatant liquid. This was centrifuged, washed with 50% ethanol (4 ml.), 90% ethanol (5 ml.) and dried *in vacuo* over P_2O_5 ; yield 22 mg. of hamamelonic acid diphosphate Ba salt (B).

Both A and B were cochromatographed with the products of the reaction of $KC^{14}N$ on RuDP and coincidence of radioactivity and the phosphomolybdate color was obtained.

Chromatography in the solvent system of Cowgill, as modified by Trudinger, showed a sharp separation from RuDP.

Preparation of Carrier Hamamelonic Acid.—Hamamelonic acid diphosphate A (5 mg.) was dissolved in HCl (1 ml. of 1 *N*) and heated for 16 hr. in a sealed tube at 110°. For carrier 100 μ l. of this solution was used. Before use it was treated with Na_2SO_4 solution (50 μ l. of 0.05 *M*) and centrifuged.

Results

A radioautograph of a typical $C^{14}O_2$, CN^- experiment is shown in Fig. 7. The general picture is rather similar to that obtained in the absence of cyanide. From a cursory examination it might be concluded that cyanide is acting only as an enzyme poison. That this is far from the truth is illustrated in Fig. 8, which is the radioautograph of a typical experiment, CO_2 , $C^{14}N^-$. The sole source of radioactivity here is cyanide; hence, all the spots are the products of the reactions of cyanide with materials present in the algae.

Radioautographs of dephosphorylated diphosphate areas of the above experiments are shown in Figs. 9 and 10 (enzymic hydrolysis) and in Figs. 11 and 12 (acid hydrolysis). Both of these radioautographs show large amounts of a material not observed before in experiments without cyanide. Since it has been demonstrated that cyanide participated in the formation of this material, it is probable that this substance is formed by a cyanohydrin reaction. As this substance originates in the diphosphate area, the possibility was investigated that it was produced by a cyanohydrin reaction on ribulose diphosphate. This reaction has been shown to yield hamamelonic acid diphosphate (together possibly with some epimer). A comparison of the electrophoretic behavior of the diphosphate area of the experiment, $C^{14}O_2$, CN^- , with synthetic hamamelonic acid diphosphate (produced by the non-enzymatic action of $C^{14}N^-$ on RuDP) is illustrated in Fig. 13. It can be seen that the two materials are electrophoretically indistinguishable under these conditions.

The diphosphate areas obtained using the material from experiments $C^{14}O_2$, CN^- and CO_2 , $C^{14}N^-$ were chromatographed with carrier RuDP and carrier HmDP using the solvent system of Trudinger. The radioactive areas coincided with the areas reacting to the phosphate spray. In a control where RuDP was added as a carrier, no such coincidence was obtained. This is evidence that the new material produced in experiments with cyanide on the algae is hamamelonic acid diphosphate, produced by the action of the cyanide on RuDP, either inside the algae or during the extraction procedure. A photograph of the sprayed paper was not included

as it was not possible to photograph it before it disintegrated.

Further confirmation was obtained by hydrolyzing the diphosphate area in 1 *N* HCl for 16 hr. The material obtained from experiments using C¹⁴O₂, CN⁻ and CO₂, C¹⁴N⁻ was shown to be identical with synthetic hamamelonic acid (made from HmDP) by cochromatography in one dimension in butanol-propionic acid-water.

Conclusions

It has been shown that the new material which appears in photosynthesis experiments with algae by the use of high concentrations of cyanide is hamamelonic acid diphosphate together with some epimer. This compound has not been detected in normal photosynthesis experiments without cyanide addition. A substance which is chromatographically rather similar but not identical has been detected by Moses³⁴ in normal photosynthesis experiments. This material on dephosphorylation and reduction by KBH₄ yields hamamelonic acid and its isomers.

The source of hamamelonic acid diphosphate is clearly a cyanohydrin reaction on RuDP. In order to form appreciable amounts of HmDP, it is essential to have a light period between the cyanide addition and the injection of boiling alcohol to kill the algae.¹⁵ Furthermore, the net fixation of radiocarbon as PGA diminishes when the algae are illuminated after the addition of cyanide. The amount of radiocarbon fixed as hamamelonic acid

(34) V. Moses, unpublished work from this Laboratory.

diphosphate is consistent with the amount disappearing from the pentose phosphates and the PGA pools. This can be interpreted if it is assumed that the addition of cyanide has little effect on the ability of the algae to reduce PGA at least for an initial short period of time. Then in the light period after the cyanide addition, PGA is reduced to triose phosphate which cycles into RuDP and is trapped by the cyanide. The principal effect of the cyanide would then be to block the uptake of CO₂, probably by inhibition of the carboxydismutase system,¹⁴ and trap the RuDP as it accumulated from other intermediates of the photosynthetic cycle. In addition, it can be deduced that the reaction involving the phosphorylation of RuMP is also occurring during the light period after the addition of cyanide. This is consistent with the finding of Kandler,³⁵ who has taken his results on cyanide inhibition to indicate that a light phosphorylation is not inhibited by high concentrations of cyanide in the first few minutes. However, the validity of this method of determining phosphorylation rates must be re-examined in the light of the broad reactivity of CN⁻ here demonstrated.

It also has been demonstrated that a facile reaction occurs between RuDP and KCN to produce HmDP. It is feasible that this reaction occurs readily because the intermediate cyanohydrin can cyclize to give an imino-lactone which should undergo rapid hydrolysis.

(35) O. Kandler, *Naturwiss.*, **42**, 390 (1955).

BERKELEY 4, CALIFORNIA

[CONTRIBUTION FROM THE DEPARTMENT OF CHEMISTRY, WASHINGTON UNIVERSITY]

The Condensation of 2-Nitroethanol with the D-Aldopentoses

BY JOHN C. SOWDEN AND DONALD R. STROBACH

RECEIVED JANUARY 20, 1958

D-*gluco*-Heptulose, D-*manno*-heptulose, D-*galacto*-heptulose, D-*talo*-heptulose, 2,7-anhydro-β-D-*ido*-heptulopyranose and 2,7-anhydro-β-D-*altro*-heptulopyranose (sedoheptulosan) have been obtained by application of the 2-nitroethanol synthesis to the appropriate D-aldopentoses.

A preliminary communication¹ from this Laboratory reported the condensation of sodium *aci*-2-nitroethanol with D-arabinose to give, after hydrolysis of the intermediate sodium *aci*-2-deoxy-2-nitroheptitols, D-*gluco*-heptulose and D-*manno*-heptulose. The synthesis now has been extended to D-lyxose, D-xylose and D-ribose to give, respectively, D-*galacto*-heptulose plus D-*talo*-heptulose, 2,7-anhydro-β-D-*ido*-heptulopyranose and 2,7-anhydro-β-D-*altro*-heptulopyranose (sedoheptulosan). Although the yields (5–22%) by this method are not high, the relative simplicity of the synthesis rates it favorably with other available methods for obtaining the higher-carbon ketoses.

2,7-Anhydro-β-D-*ido*-heptulopyranose has been synthesized previously² from D-xylose through consecutive condensations with nitromethane and formaldehyde, followed by hydrolysis of the resulting *aci*-2-deoxy-2-nitroheptitols. However, the

present synthesis using 2-nitroethanol provides the anhydroheptulose with greater ease and in considerably higher yield.

The laboratory preparation of 2-nitroethanol is accomplished readily by the condensation of nitromethane with formaldehyde,³ and the product can be isolated safely by co-distillation with diphenyl ether.⁴ Extraction of the separated, crude 2-nitroethanol thus obtained with an equal volume of hexane or low-boiling petroleum ether, to remove diphenyl ether, then gives a product (50% yield based on formaldehyde) of sufficient purity for condensation with the sugars.

Experimental

D-*gluco*-Heptulose and D-*manno*-Heptulose.—To a suspension of 10 g. of D-arabinose in 24 ml. of 2-nitroethanol

(3) I. M. Gorsky and S. P. Makarow, *Ber.*, **67**, 996 (1934); J. Controulis, M. C. Rebstock and H. M. Crooks, Jr., *THIS JOURNAL*, **71**, 2463 (1949).

(4) H. T. Roy, Jr. (to the General Tire and Rubber Co.), U. S. Patent 2,710,830, June 14, 1955; W. E. Noland, H. I. Freeman and M. S. Baker, *THIS JOURNAL*, **78**, 188 (1956).

(1) J. C. Sowden, *THIS JOURNAL*, **72**, 3325 (1950).

(2) J. K. N. Jones, *J. Chem. Soc.*, 3643 (1954).