

(III) which was cyclized by base to dihydroconesine (IV) (80% yield from II), m.p. 101.5–102.5°,  $[\alpha]_D^{26} +53.5^\circ$ , infrared spectrum identical with that of authentic dihydroconesine, mixture m.p. undepressed.

The above synthesis of dihydroconesine confirms the previously assigned structure<sup>6</sup> and stereochemistry.<sup>7</sup>

The introduction of other functional groups at C<sub>18</sub> via intermediates such as III is an obvious possibility which is presently under investigation in these laboratories. In addition, the application of these methods to the functionalization of methyl groups in other systems is under study.<sup>8</sup>

(6) R. D. Haworth, J. McKenna, R. G. Powell and G. H. Whitfield, *Chem. and Ind.*, 215 (1952).

(7) V. Cerny, L. Labler and F. Sorm, *Coll. Czech. Chem. Comm.*, **22**, 76 (1957).

(8) Drs. O. Jeger, D. Arigoni and co-workers have also been concerned with this problem and our results are published simultaneously with theirs by friendly agreement.

(9) Predoctoral Research Fellow (AF-7544, 1957–58) of the National Institute of Arthritis and Metabolic Diseases; Alfred P. Sloan Foundation Fellow (1956, 1957).

DEPARTMENT OF CHEMISTRY AND  
CHEMICAL ENGINEERING  
UNIVERSITY OF ILLINOIS  
URBANA, ILLINOIS

E. J. COREY  
W. R. HERTLER<sup>9</sup>

RECEIVED MAY 1, 1958

**6 $\alpha$ -METHYL-17 $\alpha$ -HYDROXYPROGESTERONE  
17-ACYLATES; A NEW CLASS OF  
POTENT PROGESTINS<sup>1</sup>**

Sir:

Recent communications from These Laboratories<sup>2,3</sup> and others<sup>4</sup> have described the preparation of a number of 6-methyl steroids. We now wish to report the synthesis of 6 $\alpha$ -methyl-17 $\alpha$ -hydroxyprogesterone (Ia), the acetate (II) of which is believed to be the most active progestational agent yet known.

The bisethylene acetal (IV) of 17 $\alpha$ -hydroxyprogesterone (III)<sup>5</sup> was treated with peracetic acid to give a mixture of 5 $\alpha$ ,6 $\alpha$ -epoxy-17 $\alpha$ -hydroxypregnane-3,20-dione bisethylene acetal (Va), m.p. 216–218.5°,  $[\alpha]_D -70^\circ$ , and the corresponding 5 $\beta$ ,6 $\beta$ -epoxide (Vb), m.p. 170–172.5°,  $[\alpha]_D -14^\circ$ , which could be separated by crystallization from acetone. The  $\alpha$ -epoxide (Va) when refluxed with methylmagnesium bromide in tetrahydrofuran, afforded the bisethylene acetal of 5 $\alpha$ ,17 $\alpha$ -dihydroxy-6 $\beta$ -methylpregnane-3,20-dione (VI), m.p. 160–163°,  $[\alpha]_D -38^\circ$ . Upon hydrolysis in acidic acetone, 5 $\alpha$ ,17 $\alpha$ -dihydroxy-6 $\beta$ -methylpregnane-3,20-dione (VII), m.p. 274–279°,  $[\alpha]_D -6^\circ$ , was produced. Dehydration of VI by very dilute sodium hydroxide in pyridine afforded 6 $\beta$ -methyl-17 $\alpha$ -hydroxyproges-


(1) Rotations were in chloroform unless otherwise specified; ultraviolet maxima were determined in 95% alcohol.

(2) G. B. Spero, J. L. Thompson, A. R. Hanze, H. C. Murray, O. K. Sebek and J. A. Hogg, *THIS JOURNAL*, **78**, 6213 (1956).

(3) J. A. Campbell, J. C. Babcock and J. A. Hogg, *ibid.*, in press.

(4) (a) O. S. Madaeva, M. I. Ushakov, N. F. Kosheleva, *J. Gen. Chem. (USSR)*, **10**, 213 (1940); *C. A.*, **34**, 7292 (1944); (b) M. Ehrenstein, *J. Org. Chem.*, **8**, 83 (1943); (c) G. Cooley, B. Ellis, D. N. Kirk, and V. Petrov, *J. Chem. Soc.*, 4412 (1957), and preceding papers; (d) H. J. Ringold, E. Batres and G. Rosenkranz, *J. Org. Chem.*, **22**, 99 (1957).

(5) S. Bernstein, M. Heller and S. M. Stolar, *THIS JOURNAL*, **76**, 5674 (1954).



terone (Ib), m.p. 232–240°,  $[\alpha]_D +34^\circ$ ,  $\lambda_{\max}$  242 m $\mu$  ( $a_M$  16,500), which was epimerized in chloroform<sup>6</sup> saturated with gaseous hydrogen chloride to 6 $\alpha$ -methyl-17 $\alpha$ -hydroxyprogesterone (Ia), m.p. 220–223.5°,  $[\alpha]_D +75^\circ$ ,  $\lambda_{\max}$  241 m $\mu$  (16,150). Alternatively, dehydration and epimerization of VII to Ia could be effected directly with chloroform–hydrogen chloride.

Acylation of Ia with acetic anhydride–acetic acid–*p*-toluenesulfonic acid<sup>7</sup> produced the 17-acetate II, m.p. 205–209°,  $[\alpha]_D +56^\circ$ ,  $\lambda_{\max}$  240 m $\mu$  ( $a_M$  15,950). With cyclopentylpropionic acid and trifluoroacetic anhydride<sup>8</sup> there was obtained the 17-( $\beta$ -cyclopentylpropionate), m.p. 135–137°,  $[\alpha]_D +44^\circ$ ,  $\lambda_{\max}$  240.5 m $\mu$  ( $a_M$  15,775). Similarly, the propionate (m.p. 155–157°,  $[\alpha]_D +45^\circ$  (EtOH),  $\lambda_{\max}$  240 m $\mu$  ( $a_M$  16,075)), caproate (m.p. 105–107°,  $[\alpha]_D +46^\circ$ ,  $\lambda_{\max}$  240 m $\mu$  ( $a_M$  15,300)), phenylacetate (m.p. 164–166°,  $[\alpha]_D +62^\circ$  (EtOH),  $\lambda_{\max}$  240 m $\mu$  ( $a_M$  16,125)) and related esters were prepared.

In the McPhail modification of the Clauberg assay<sup>9</sup> 6 $\alpha$ -methyl-17 $\alpha$ -hydroxyprogesterone 17-acetate (II, Provera<sup>10</sup>) was 50–60 times more active than progesterone on subcutaneous administration and 100–300 times more active than ethisterone

(6) Commercial chloroform (containing 0.7% alcohol) was used.

(7) (a) Huang-Minlon, E. Wilson, N. L. Wendler, and M. Tishler, *THIS JOURNAL*, **74**, 5394 (1952); (b) R. B. Turner, *ibid.*, **76**, 3489 (1953).

(8) E. J. Bourne, M. Stacey, J. C. Tatlow and J. M. Tedder, *J. Chem. Soc.*, 2976 (1949).

(9) M. K. McPhail, *J. Physiol.*, **83**, 145 (1934).

(10) Registered Trademark, The Upjohn Company.

(17 $\alpha$ -ethinyltestosterone) on oral administration.<sup>11</sup> As an ovulation inhibitor in the rabbit, II was 10-20 times more active parenterally than progesterone. When administered parenterally to pregnant rats castrated on the 8th day of pregnancy, II was 25-100 times progesterone in maintaining pregnancy to term. These results, which indicate 6 $\alpha$ -methyl-17 $\alpha$ -hydroxyprogesterone 17-acetate to be a progestational agent of exceptional potency, will be described in greater detail shortly.<sup>12</sup>

(11) In the same assays, the corresponding non-methylated steroid, 17 $\alpha$ -hydroxyprogesterone 17-acetate,<sup>7b</sup> was ca. 6 times progesterone (parenterally) and 2-4 times ethisterone (orally).

(12) The chemical portion of this work will be submitted to THIS JOURNAL (Babcock, Gutsell, Herr and Hogg); the endocrine studies will be reported elsewhere (Stucki, Barnes and Dulin).

RESEARCH LABORATORIES  
THE UPJOHN COMPANY  
KALAMAZOO, MICHIGAN

JOHN C. BABCOCK  
ERWIN S. GUTSELL  
MILTON E. HERR  
JOHN A. HOGG  
JACOB C. STUCKI  
LESTER E. BARNES  
WILLIAM E. DULIN

RECEIVED APRIL 24, 1958

### DIRECT INTRODUCTION OF A NITROGEN FUNCTION AT C-18 IN A STEROID<sup>1</sup>

Sir:

Current interest in aldosterone,<sup>1a</sup> has prompted several investigations on the synthesis of C-18 oxygenated steroids, total<sup>2</sup> as well as partial synthesis using steroids lacking substitution at C-18 as starting material.<sup>3</sup> In the latter case opening of ring D has been used for modification of C-18.

We have developed a third way of approaching the problem, the direct introduction of a substituent into the *intact* tetracyclic steroid molecule by the Loeffler-Freytag reaction.<sup>4,5</sup>

3 $\beta$ -Acetoxy-20-keto-5 $\alpha$ -pregnane (I)<sup>6</sup> was converted via the oxime (II) (m.p. 198°, [ $\alpha$ ]<sub>D</sub><sup>20</sup> +17° (CHCl<sub>3</sub>), *Anal.* found: C, 73.49; H, 9.94) to 3 $\beta$ -acetoxy-20 $\alpha$ -amino-5 $\alpha$ -pregnane (III)<sup>7</sup> (m.p. 171°; *pK*<sup>\*8</sup> 9.30) (Pt-H<sub>2</sub>), thence with ethyl formate to the N-formyl derivative (IV) (m.p. 185-186°, [ $\alpha$ ]<sub>D</sub><sup>19</sup> -3° (CHCl<sub>3</sub>), *Anal.* found: C, 74.01; H, 10.13; N, 3.52) and further with LiAlH<sub>4</sub> to 3 $\beta$ -hydroxy-20 $\alpha$ -N-methylamino-5 $\alpha$ -pregnane

(1) This paper is part of a joint effort of CIBA AG,<sup>2a</sup> Basle, Organische Anstalt der Universitat,<sup>2b</sup> Basle, N.V. Organon,<sup>2c</sup> Oss, and organisch-chemisches Laboratorium der Eidg. Technischen Hochschule, Zurich, on synthesis of C-18 oxygenated steroids.

(1a) S. Á. Simpson, J. F. Tait, A. Wettstein, R. Neher, J. von Euw, O. Schindler and T. Reichstein, *Helv. Chim. Acta*, **37**, 1163, 1200 (1954).

(2) (a) P. Wieland, K. Heusler, H. Ueberwasser and A. Wettstein, *Helv. Chim. Acta*, **41**, 416 (1958) and earlier papers; (b) A. Lardon, O. Schindler and T. Reichstein, *ibid.*, **40**, 666, 1034 (1957); (c) W. J. van der Burg, D. A. van Dorp, O. Schindler, C. M. Siegmann and S. A. Szpilfogel, *Rev. Trav. Chim.*, **77**, 171 (1958), and previous papers.

(3) D. H. R. Barton, A. da S. Campos-Neves and A. I. Scott, *J. Chem. Soc.*, 2698 (1957).

(4) (a) A. W. Hofmann, *Ber.*, **18**, 5, 109 (1885); (b) K. Loeffler and C. Freytag, *Ber.*, **42**, 3427 (1909); K. Loeffler, *ibid.*, **43**, 2035 (1910).

(5) For references cf. R. Lukes and M. Ferles, *Coll. Czech.*, **20**, 1227 (1955).


(6) A. Butenandt, U. Westphal and W. Hohlwer, *Z. physiol. Chem.*, **227**, 84 (1934).

(7) The stereochemistry of C-20 in III and related compounds is discussed below.

(8) Measured in methylcellosolve according to W. Simon, E. Kovats, L. H. Chopard-dit-Jean and E. Heilbronner, *Helv. Chim. Acta*, **37**, 1872 (1954).

(V) (m.p. 211°, [ $\alpha$ ]<sub>D</sub><sup>20</sup> +27° (CHCl<sub>3</sub>), *pK*<sup>\*9</sup> 9.17, *Anal.* found: C, 79.20; H, 11.79; N, 4.16). Oxidation of (V) (CrO<sub>3</sub>-HOAc) gave the 3-keto-derivative (VI) (*pK*<sup>\*9</sup> 9.20; N-acetate: m.p. 216-217°, [ $\alpha$ ]<sub>D</sub><sup>19</sup> +15° (CHCl<sub>3</sub>), *Anal.* found: C, 77.06; H, 10.69) which was reduced by the Huang-Minlon procedure to 20 $\alpha$ -N-methylamino-5 $\alpha$ -pregnane (VII) (m.p. 96°, [ $\alpha$ ]<sub>D</sub><sup>20</sup> +32° (CHCl<sub>3</sub>), *pK*<sup>\*9</sup> 9.32, *Anal.* found: C, 83.22; H, 12.33; N, 4.31). The N-chloro derivative (VIII), made from (VII) according to the procedure of Ruschig, *et al.*,<sup>9</sup> furnished by treatment with a mixture of concd. sulfuric and acetic acid<sup>10</sup> a tertiary base, m.p. 107°, formulated as the hitherto unknown conanine (IX),<sup>11,12</sup> [ $\alpha$ ]<sub>D</sub><sup>19</sup> +61° (CHCl<sub>3</sub>), (*pK*<sup>\*8</sup> 8.28 *Anal.* found: C, 83.66; H, 12.08; N, 4.43).

The constitution of (IX) was proved as follows: dihydro-isoconessimine (X)<sup>13</sup> was converted via the N-chloro derivative (XI) to 3-keto-conanine (XII) (m.p. 146°, [ $\alpha$ ]<sub>D</sub><sup>20</sup> +80° (CHCl<sub>3</sub>), *pK*<sup>\*8</sup> 8.26, *Anal.* found: C, 80.15; H, 10.64; N, 4.13 (NaOCH<sub>3</sub> followed by H<sub>3</sub>O<sup>+</sup>). Removal of the 3-keto group in the latter afforded conanine (IX) (m.p. 107°, [ $\alpha$ ]<sub>D</sub><sup>20</sup> +63° (CHCl<sub>3</sub>), *pK*<sup>\*8</sup> 8.28, *Anal.* found: C, 83.85; H, 11.79, N, 4.39), identical in all respects with the tertiary base from the cyclization reaction. Assuming that C-20 is not involved in the latter reaction the precursors of (IX) must therefore be formulated as 20 $\alpha$ -compounds.<sup>12</sup>



(9) H. Ruschig, W. Fritsch, J. Schmidt-Thome and W. Haede, *Ber.*, **88**, 883 (1955).

(10) According to a private communication of Prof. E. J. Corey the cyclization reaction of aliphatic N-chloro-amines is promoted by traces of ferrous sulfate. We thank Prof. Corey for this information. A similar effect we have now noted in the cyclization of compound (VIII).

(11) Nomenclature as proposed by R. D. Haworth and M. Michael, *J. Chem. Soc.*, 4973 (1957).

(12) For the proof of configuration of C-20 cf. V. Cerny, L. Labler and F. Sorm, *Coll. Czech.*, **22**, 76 (1957).

(13) R. D. Haworth, J. McKenna and G. H. Whitfield, *J. Chem. Soc.*, 1102 (1953).