



Further investigation of allyldiazomethane and of bicyclobutane is in progress.

**Acknowledgment.**—The authors are grateful to Miss Elizabeth Kroes for skillful technical assistance and to the National Science Foundation and the American Chemical Society (PRF) for generous financial support.

DEPARTMENT OF CHEMISTRY  
CHEMISTRY RESEARCH BLDG.  
UNIVERSITY OF WISCONSIN  
MADISON 15, WISCONSIN


DAVID M. LEMAL  
FREDRIC MENGES  
GEORGE W. CLARK

RECEIVED JUNE 3, 1963

## Electronic Interactions in Biradicals. I. Reversible Bond Formation

Sir:

There is considerable interest in the chemistry of biradicals which, in some cases, may interact sufficiently to be termed triplets.<sup>1</sup> The synthesis of a potential triplet containing two phenoxy radicals in a cross-conjugated system has been reported by Yang and Castro.<sup>2</sup> We have studied a system containing two individually stable phenoxy radicals not connected to a single  $\pi$ -electron system but held together as closely as possible—bound to the same saturated carbon atom. The system we chose is derived from 2,2-bis-(3,5 di-*t*-butyl-4-hydroxyphenyl)-propane (I).<sup>3</sup>



Metal oxides ( $\text{PbO}_2$ ,  $\text{Ag}_2\text{O}$ ,  $\text{MnO}_2$ ) oxidized I readily, but treatment of a benzene solution of I with aqueous alkaline  $\text{K}_3\text{Fe}(\text{CN})_6$  was most effective for preparative scale work. Evaporation of the solvent with nitrogen left brick-red needles (m.p. ca.  $120^\circ$  dec.) which could be crystallized from pentane. Iodometric titration indicated the consumption of 1.85 equivalents of HI and regenerated I (55% crystallized). The infrared spectrum of the product was interpretable in terms of structure IV, resulting from internal bond formation in the biradical III. There was no band at  $6.4 \mu$ , the characteristic absorption of phenoxy radicals,<sup>4</sup> but there was a doublet, characteristic of this type of cyclohexadienone, at  $6.08$  and  $6.16 \mu$ .<sup>4</sup> Examination of a scale model of III indicated that the *para* positions of the radicals are held close together (the internuclear distance is  $2.5 \text{ \AA}$ . when the central atom is tetrahedral) and a strained bond between these centers is conceivable. A simple spirodienone of this type has been reported by Baird and Winstein.<sup>5</sup>

The oxidation product, IV, was stable as a solid or in an inert solvent as long as it was protected from oxygen. The solid was decolorized slowly by air and solu-

tions of IV reacted readily with oxygen, forming an unstable addition compound. The latter is formulated as the peroxide V on the basis of its analysis (*Anal.* Calcd. for  $\text{C}_{31}\text{H}_{46}\text{O}_4$ : C, 77.13; H, 9.61. Found: C, 76.70; H, 9.84) and its infrared spectrum ( $\lambda_{\text{max}}^{\text{CH}_2\text{Cl}_2}$   $5.97$ ,  $6.07 \mu$ ).

The oxidation of I (benzene or cyclohexane solution) also was carried out with  $\text{Ag}_2\text{O}$  or  $\text{PbO}_2$  in a sealed, degassed system. The initially formed product was identified as the monoradical II by virtue of its e.p.r. spectrum. The major triplet splitting is attributable to the two equivalent *meta* protons ( $\alpha = 1.67$  gauss) with further smaller coupling to the six equivalent protons of the isopropylidene group ( $\alpha = 0.38$  gauss). There is no significant interaction with the adjacent unoxidized phenolic system although the optical spectrum lacks the long wave length band responsible for the blue color of 2,4,6-tri-*t*-butylphenoxy. Further oxidation of the solution caused a marked decrease in the intensity of the e.p.r. spectrum; the addition of excess phenol I regenerated the original spectrum. These observations are readily interpretable in terms of the equilibrium  $\text{I} + \text{IV} \rightleftharpoons 2\text{II}$ . No e.p.r. signal attributable to biradical III was observed and there was no absorption in rigid media other than at  $g = 2$ .

The ultraviolet spectrum of phenol I is essentially that of a simple phenol and has a doublet maximum at  $275$  and  $283 \text{ m}\mu$ . The optical absorption spectrum of the solution was monitored simultaneously with the e.p.r. spectrum as oxidation of the phenol progressed. Initially, a broad absorption at  $325 \text{ m}\mu$  appears as well as a sharp peak at  $263 \text{ m}\mu$ . As the oxidation proceeds, the latter increases in intensity while the former, which we attribute to the monoradical II, decreases. The peak at  $263 \text{ m}\mu$  corresponds well with that reported for spiro(2,5)octa-(1,4)-diene-3-one by Baird and Winstein.<sup>5</sup> The long wave length peak tails into the visible and is responsible for the red color of the product. The variations in e.p.r. intensity as oxidation proceeded were in accord with the conclusions reached on the basis of the optical spectral data.

The intensity of the e.p.r. spectrum of II decreased as the temperature was lowered, indicating that a thermal equilibrium was present. This is the disproportionation of II into I and IV. The Curie dependence of magnetic susceptibility may be allowed for by taking intensity ( $I$ )  $\times$  temperature ( $T$ ) as a measure of radical concentration. An Arrhenius plot of  $\log IT$  vs.  $1/T$  over the range  $25$  to  $-56^\circ$  ( $\text{CH}_2\text{Cl}_2$  solution) was linear and gave  $\Delta H = -3.7$  kcal. The low value reflects the weakness of the strained cyclopropane bond in IV.

It appears that IV is in equilibrium with biradical III on the basis of both physical and chemical evidence although III itself could not be detected. Dienone IV would not be expected to be destroyed rapidly by oxygen, but the reaction takes only a few minutes in solution at  $25^\circ$ . The n.m.r. spectrum of I was in accord with its assigned structure ( $\tau$  scale: *t*- $\text{C}_4\text{H}_9$ , 1.33;  $\text{CH}_3$ , 1.55; OH, 4.76; ArH, 6.81). In solutions of IV, however, only a single moderately broad line was observed ( $1.33 \tau$ ). The absence of the vinyl proton line can be explained by rapid interconversion of III and IV. The amount of monoradical II in solution was too small to permit an explanation based on hydrogen atom exchange.<sup>6</sup> The sharpness of the butyl line may be explained by the low spin density at this position.<sup>7,8</sup> The *t*-butyl proton splitting was not resolved in the e.p.r. spectrum.

(1) See S. I. Weissman, *Ann. Rev. Phys. Chem.*, **12**, 151 (1961).

(2) N. C. Yang and A. J. Castro, *J. Am. Chem. Soc.*, **82**, 6208 (1960).

(3) We thank the Ethyl Corp., Detroit, Michigan, for a gift of this compound. It was prepared by butylation of the corresponding bisphenol with isobutylene and  $\text{Al}(\text{OC}_2\text{H}_5)_3$ . I has m.p.  $155$ – $156^\circ$ . *Anal.* Found: C, 81.94; H, 10.74.

(4) E. Muller and K. Ley, *Chem. Ber.*, **87**, 922 (1954).

(5) R. Baird and S. Winstein, *J. Am. Chem. Soc.*, **85**, 567 (1963).

(6) R. Kreilick and S. I. Weissman, *ibid.*, **84**, 306 (1962).

(7) This phenomenon has been observed by Weissman and Kreilick in line broadening due to hydrogen atom exchange.

(8) H. McConnell and S. B. Berger, *J. Chem. Phys.*, **27**, 230 (1957).