

The reaction of methyllithium with vinyltriphenylphosphonium bromide in ether-tetrahydrofuran medium can be understood in the same manner, although the products observed account for only *ca.* 30% of the lithium reagent consumed. When the reaction mixture was quenched with cyclohexanone, benzene (17%), *n*-propylidenecyclohexane (13%), and methylcyclohexane (6%) were obtained.

All reactions were carried out under an atmosphere of prepurified nitrogen. All new compounds reported had satisfactory analyses. Our work in this area is continuing.

Acknowledgment.—The authors are grateful to the U. S. Army Research Office (Durham) and to the Alfred P. Sloan Foundation for generous support of this work and to M & T Chemicals for gifts of triphenylphosphine.

(17) (a) Alfred P. Sloan Research Fellow; (b) National Science Foundation Graduate Fellow, 1960–1963; National Institutes of Health Predoctoral Fellow, 1963–1964; (c) Fellow of the M.I.T. School for Advanced Study, 1961–1962.

DEPARTMENT OF CHEMISTRY DIETMAR SEYFERTH^{17a}
 MASSACHUSETTS INSTITUTE OF TECHNOLOGY JOSEPH S. FOGEL^{17b}
 CAMBRIDGE, MASSACHUSETTS JAMES K. HEEREN^{17c}


RECEIVED NOVEMBER 22, 1963

Tetrahydrohomofolate, a Specific Inhibitor of Thymidylate Synthetase

Sir:

This communication describes the synthesis and biological properties of homofolic acid (a trivial name for the compound possessing an additional methylene group between positions 9 and 10 of folic acid) and its reduced forms, dihydrohomofolate (homofolate-H₂) and tetrahydrohomofolate (homofolate-H₄).

The general approach of Boon and Leigh¹ for the synthesis of unambiguously 6-substituted 2-amino-4-hydroxypteridines was used to prepare XII. The *N*-phenyl-β-alanine derivative (I) was converted *via* II, III, IV, and V to the aminoketone (VI), the semicarbazone of which was condensed with 2-amino-4-



IX, R₁ = H; R₂ = Ac; R₃ = OEt

X, R₁ = R₂ = H; R₃ = OH

XI, R₁ = Ac; R₂ = CF₃CO; R₃ = OH

XII, R₁ = R₂ = H; R₃ = NHCH(CO₂H)CH₂CH₂CO₂H

(1) W. R. Boon and T. Leigh, *J. Chem. Soc.*, 1497 (1951).

hydroxy-5-phenylazo-6-chloropyrimidine to give the intermediate (VII), after hydrolysis of the semicarbazone function. Hydrogenation of VII resulted in spontaneous cyclization of the ketone carbonyl with the formed 5-amino group to give the 7,8-dihydropteridine (VIII), $\lambda_{\text{max}}^{\text{pH } 13}$ 234 m μ (ϵ 21,200), 277 (7780), 330 (5180), that was oxidized to the pteridine (IX). Saponification gave chromatographically homogeneous "homopteroic acid" (X), $\lambda_{\text{max}}^{\text{pH } 13}$ 256 m μ (ϵ 26,900), 277 (21,900), 365 (7625). *Anal.*² Found: C, 54.7, H, 4.62; N, 25.6. The blocked derivative (XI), after reaction with diethyl L-glutamate by the mixed anhydride method followed by saponification, gave XII, $\lambda_{\text{max}}^{\text{pH } 13}$ 255 m μ (ϵ 24,600), 281 (19,500), 365 (7880). *Anal.* Found (for the hemihydrate): C, 51.2; H, 5.09; N, 21.2. On paper chromatography in 0.1 M (NH₄)HCO₃, 99% of the material was present in a single spot with *R_f* 0.89. The method of synthesis precludes a 7-substituted compound and, considering the ultraviolet spectra that support the pteridine structure, the analytical data, and the chromatographic behavior, there can be little doubt that homofolic acid has structure XII.

Homofolate was converted to the dihydro derivative by dithionite reduction in 1 M 2-mercaptoethanol³ and then was tested spectrophotometrically⁴ as a substrate of dihydrofolate reductase from amethopterin-resistant mouse leukemia cells. Homofolate-H₂ was as effective a substrate as folate-H₂. The enzymatically formed homofolate-H₄ was not only completely inert as a participant in thymidylate synthesis but was a potent inhibitor of this enzyme. At 2.0×10^{-6} M, homofolate-H₄ caused a 50% inhibition of thymidylate synthetase from *E. coli*⁵ in the presence of 80 times as much folate-H₄. Homofolate-H₄ analogs containing α -, β -, and γ -methyl glutamic acid moieties⁶ were less inhibitory than homofolate-H₄.

Data obtained in a survey of the possible inhibitory action of homofolate-H₄ on a variety of tetrahydrofolate-requiring reactions indicate that thymidylate synthetase from *E. coli* appears to be the most sensitive of the enzymes tested (Table I).

In 0.006 M mercaptoethanol, homofolate-H₄ oxidized within 20 min., giving a compound with a spectrum closely resembling that of dihydrohomofolate. Addition of mercaptoethanol after oxidation did not reverse the reaction. Homofolate-H₄ bound formaldehyde mole for mole as does folate-H₄. In the presence of formaldehyde, homofolate-H₄ was stabilized over a period of 1 hr. at room temperature.

Homofolate is a poor growth inhibitor whereas homofolate-H₄ is a potent inhibitor of *Streptococcus faecalis* (ATCC 8043) and *Lactobacillus casei* (ATCC 7469, Table II). The inhibition is completely reversed for both organisms by thymidine (25 γ /ml.) even at levels of homofolate-H₄ 140 times greater than that required for 50% inhibition. *Pediococcus cerevisiae* (ATCC 8081) is not affected by homofolate-H₄. This pattern of behavior is distinct from other antifolate agents.⁷ Homofolate-H₂ surprisingly showed growth-factor activity for *S. faecalis*. It was about one-tenth as active as folate. This activity was not due to contamination with folate, as growth-promoting activ-

(2) Acceptable analytical data were obtained for all the solid substances encountered in the synthesis.

(3) M. Friedkin, E. J. Crawford, and D. Misra, *Federation Proc.*, **21**, 176 (1962).

(4) M. Friedkin, E. J. Crawford, S. R. Humphreys, and A. Goldin, *Cancer Res.*, **22**, 600 (1962).

(5) A. J. Wahba and M. Friedkin, *J. Biol. Chem.*, **237**, 3794 (1962).

(6) Substituted glutamates used in the synthesis of homofolate analogs were generously supplied by Dr. Alton Meister.

(7) R. L. Kisliuk, *Nature*, **188**, 584 (1960).

TABLE I
INHIBITION OF VARIOUS TETRAHYDROFOLATE-REQUIRING
REACTIONS BY HOMOFOLATE-H₄

Enzyme preparation	Source of enzyme and assay ref.	Folate-H ₄ ^a	
		homo-folate-H ₄	Inhibition, %
Thymidylate synthetase	<i>E. coli</i> ^b	95	47
		38	61
		19	77
		10	81
Methionine synthetase	<i>E. coli</i> ^c	1	25
Formylglutamate formyl transferase	Hog liver ^d	15	10
		1	30
Formyltetrahydrofolate synthetase ^e	<i>Clostridium cylindrosporum</i> ^f	5	0
		1	0
Methylene tetrahydrofolate dehydrogenase	<i>E. coli</i> ^{g,h}	3	37
		2	43
		1	63
		4	0
Purine synthetase	Pigeon liver ⁱ	2	3
		1	39
		2	8
		1	33
L-Serine hydroxymethyl transferase	<i>E. coli</i> ^{g,j}	2	17
		1	33
		2	0
Deoxycytidylate hydroxymethylase	T ₄ am82-phage-infected <i>E. coli</i> ^{k,l}	2	0

^a Ratio of tetrahydrofolate concentration to tetrahydrohomofolate concentration in assay mixture. Homofolate-H₄ was prepared by dithionite reduction of homofolate at 75° exactly as described by Silverman and Noronha for the preparation of tetrahydrofolate⁸ and was purified by gradient elution from DEAE-cellulose. It was stored frozen in a buffer mixture containing 0.005 M Tris (pH 7.0), 0.2 M mercaptoethanol, and 0.4 M NaCl. Homofolate-H₄ was also prepared by catalytic reduction with hydrogen and PtO₂ [R. L. Kisliuk, *J. Biol. Chem.*, **227**, 805 (1957)]. ^b See ref. 5. ^c R. L. Kisliuk, *J. Biol. Chem.*, **236**, 817 (1961). ^d M. Silverman, J. C. Keresztesy, G. J. Koval, and R. C. Gardiner, *ibid.*, **226**, 83 (1957). ^e A generous gift of crystalline enzyme from the laboratory of Dr. J. Rabinowitz. ^f J. C. Rabinowitz and W. E. Pricer, Jr., *ibid.*, **237**, 2898 (1962). ^g K. G. Scrimgeour and F. M. Huennekens, *Biochem. Biophys. Res. Commun.*, **2**, 230 (1960). ^h Method used is described in footnote g. TPN was used as cofactor. ⁱ D. A. Goldthwait and G. R. Greenberg, "Methods in Enzymology," Vol. II, S. P. Colowick and N. O. Kaplan, Ed., Academic Press, New York, N. Y., 1955, p. 504. ^j Same method was used as for methylene tetrahydrofolate dehydrogenase assay except serine was substituted for HCHO. The dehydrogenase activity was not limiting at concentrations of homofolate-H₄ that inhibited the transferase. ^k M. L. Dirksen, J. C. Hutson, and J. M. Buchanan, *Proc. Natl. Acad. Sci. U. S.*, **50**, 507 (1963). ^l Enzyme was generously supplied by Dr. J. M. Buchanan.

TABLE II
INHIBITION OF MICROBIAL GROWTH BY HOMOFOLATE AND
REDUCED DERIVATIVES

Compound ^a	Concn. required for 0.5 maximal inhibition (μg./ml.)		
	<i>S. faecalis</i> ^b	<i>L. casei</i> ^b	<i>P. cerevisiae</i> ^c
Homofolate	1000	100	>10,000
Dihydrohomofolate	Supports growth	50	>500
Tetrahydrohomofolate	0.7	6	>1000

^a All compounds were diluted in potassium ascorbate, 6 mg./ml. (pH 6.0), and added to growth media aseptically after autoclaving. Final assay contains 0.6 mg. of potassium ascorbate/ml. to stabilize reduced derivatives [H. A. Bakerman, *Anal. Biochem.*, **2**, 558 (1961)]. ^b Folate, 1.0 μg./ml. ^c Calcium dl-L-5-formyltetrahydrofolate, 1 μg./ml.

ity corresponded exactly to the eluted homofolate-H₂ obtained by gradient elution from DEAE-cellulose.

An outstanding biochemical alteration associated with the development of resistance to amethopterin in mouse leukemia is the pronounced increase of di-

hydrofolate reductase.^{4,9} To take advantage of this augmented activity, we have been seeking folate analogs that can be readily reduced to tetrahydro products with marked inhibitory effect on thymidylate synthetase. Homofolate-H₂ uniquely fulfills these requirements.

Acknowledgments.—These studies were supported by the Cancer Chemotherapy National Service Center (Contract No. SA-43-ph-1892), the Public Health Service (CA-05997), the National Science Foundation (G-15989), and the American Cancer Society (IN 23E).

(9) D. K. Misra, S. R. Humphreys, M. Friedkin, A. Goldin, and E. J. Crawford, *Nature*, **189**, 39 (1961).

LIFE SCIENCE RESEARCH
STANFORD RESEARCH INSTITUTE
MENLO PARK, CALIFORNIA
LEON GOODMAN
JOSEPH DEGRAW

DEPARTMENT OF PHARMACOLOGY
TUFTS UNIVERSITY SCHOOL OF MEDICINE
BOSTON 11, MASSACHUSETTS
ROY L. KISLIUK
MORRIS FRIEDKIN
EDWARD J. PASTORE
ELIZABETH J. CRAWFORD
LAURENCE T. PLANTE
ALY AL-NAHAS
JOHN F. MORNINGSTAR, JR.
GREGORY KWOK
LESLIE WILSON
ELAINE F. DONOVAN
JUDITH RATZAN

RECEIVED DECEMBER 13, 1963

The Synthesis of Cellulose by an Enzyme System from a Higher Plant

Sir:

This report describes the synthesis of cellulose by a particulate enzyme preparation from mung beans (*Phaseolus aureus*); the glucosyl donor in this reaction is guanosine diphosphate D-glucose, which has been shown to be enzymatically formed from guanosine triphosphate and α-D-glucose 1-phosphate.¹ Although the cell-free synthesis of cellulose has previously been demonstrated with extracts of the bacterium, *Acetobacter xylinum*, via the glucosyl donor, uridine diphosphate D-glucose,² the synthesis of this important polysaccharide has never before been shown by enzyme isolated from higher plants.

Mung bean seedlings were grown in a moist atmosphere in the dark. One hundred grams (wet weight) of hypocotyls and roots were homogenized in 1.0 M tris(hydroxymethyl)aminomethane-HCl buffer (Tris), pH 7.6, and the homogenate was filtered through cheesecloth and centrifuged at 1000g for 5 min. The supernatant fluid then was centrifuged further at 20,000g for 20 min. The precipitated material was suspended in 3 ml. of 0.1 M Tris buffer, pH 7.6, and used as the source of enzyme. This preparation contained about 53% protein and 4% cellulose. Washing of the particulate material resulted in a substantial loss of activity.

All sugar nucleotides were labeled uniformly with C¹⁴ in the glucosyl moiety (106 μc./μmole). Guanosine diphosphate D-glucose was prepared enzymatically.¹ Uridine diphosphate D-glucose and thymidine diphosphate D-glucose were also prepared enzymatically.³ Adenosine diphosphate D-glucose and cytidine diphosphate D-glucose were chemically synthesized.⁴

As shown in Table I, incubation of guanosine diphosphate D-glucose with the particles led to the incorporation of radioactivity into a material which was insoluble in hot water and hot alkali. None of the other C¹⁴-glucose nucleotides was active in this system

(1) G. A. Barber and W. Z. Hassid, unpublished data.

(2) L. Glaser, *J. Biol. Chem.*, **232**, 627 (1957).

(3) G. A. Barber, *Biochemistry*, **1**, 463 (1962).

(4) S. Roseman, J. J. Distler, J. G. Moffatt, and H. G. Khorana, *J. Am. Chem. Soc.*, **83**, 659 (1961).

(8) M. Silverman and J. M. Noronha, *Biochem. Biophys. Res. Commun.*, **4**, 180 (1961).