

but Michael and Noyes⁵ have reaffirmed the original suggestions. If the energy-transfer mechanism from benzophenone outlined in reactions 1 to 3 is correct, then the triplet state of the carbonyl compounds decomposes to free-radical products. These results show that it does not yield type 2 products. It can be inferred, therefore, that the type 2 decomposition takes place through a singlet intermediate as was originally suggested.

The free-radical products in the normal photodecomposition may well arise from a triplet state, but it is clear from the work of Parmenter and Noyes⁶ that the type 1 products (alkane and carbon monoxide) also arise from singlet excited states.

Acknowledgment.—The author wishes to thank the D.S.I.R. and the London Chemical Society for apparatus grants.

(5) J. L. Michael and W. A. Noyes, *J. Am. Chem. Soc.*, **85**, 1027 (1963).

(6) C. S. Parmenter and W. A. Noyes, *ibid.*, **85**, 416 (1963).

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF KEELE
KEELE, STAFFORDSHIRE
GREAT BRITAIN

PETER BORRELL

RECEIVED MAY 18, 1964

An X-Ray Diffraction Study of Nonplanar Carbanion Structures

Sir:

Recently, spectroscopic investigations of the compound $\text{KC}(\text{CN})_3$ have been reported by Long, Carrington, and Cravenor,¹ and Miller and Baer.² An assignment can be made, on the basis of both infrared and Raman spectra, for either a planar trigonal or a pyramidal model for the anion. Both sets of investigators favor the planar structure. A preliminary X-ray diffraction study by Anderson and Klewe³ of $\text{KC}(\text{CN})_3$ also favors a planar anion within rather large experimental limits.

We have determined, by X-ray diffraction techniques, the crystal structure of the compounds ammonium tricyanomethide, $\text{NH}_4\text{C}(\text{CN})_3$, and pyridinium dicyanomethylide, $\text{C}_5\text{H}_5\text{N}^+\text{-C}^-(\text{CN})_2$. Complete three-dimensional refinements of both structures show the trigonal carbon atom environments to be significantly nonplanar.

Both compounds crystallize in the monoclinic system with the unit cell constants shown in Table I. Both structures were solved through Patterson projections down the short axis followed by a three-dimensional analysis based on packing and other considerations.

TABLE I

Space group	$\text{NH}_4\text{C}(\text{CN})_3$ $\text{C}_{2h}^2\text{-P2}_1/\text{c}$	$\text{C}_5\text{H}_5\text{N}_3$ $\text{C}_{2h}^2\text{-P2}_1/\text{m}$
<i>a</i>	9.055 ± 0.007 Å.	7.87 ± 0.02 Å.
<i>b</i>	3.87 ± 0.010 Å.	12.512 ± 0.004 Å.
<i>c</i>	17.325 ± 0.014 Å.	3.86 ± 0.01 Å.
β	104.6 ± 0.2°	114.8 ± 0.1°
<i>Z</i>	4	2

The structure of $\text{NH}_4\text{C}(\text{CN})_3$ was refined by two-dimensional Fourier and least-squares techniques fol-

(1) D. A. Long, R. A. G. Carrington, and R. B. Cravenor, *Nature*, **196**, 371 (1962).

(2) T. A. Miller and W. K. Baer, *Spectrochim. Acta*, **19**, 73 (1963).

(3) P. Anderson and B. Klewe, *Nature*, **200**, 464 (1963).




Fig. 1.—Bond distances and angles in pyridinium dicyanomethylide.

lowed by a complete full matrix isotropic weighted least-squares. The final *R* value for the 112 independent (*hkl*) data was 8%. The *R* value for the 429 independent three-dimensional data was 10%. The observed dimensions of the $\text{C}(\text{CN})_3^-$ ion are: C-C bond length, 1.40, 1.40, and $1.40 \pm .01$ Å.; C-N bond length, 1.15, 1.14, and 1.16 ± 0.1 Å.; C-C-C bond angle $119^\circ 40'$, $119^\circ 31'$, and $119^\circ 32' \pm 1^\circ$; C-C-N bond angle 180, 180, and $180 \pm 1^\circ$. If one passes a plane through the three nitrogen atoms of this ion, the central carbon atom is found to be 0.13 Å. above this plane and the cyanocarbon atoms are all 0.08 Å. above this plane. Each C-C-N unit makes an angle of 3° with respect to its projection in this plane. There are no abnormal interionic distances.

The structure of $\text{C}_5\text{H}_5\text{N}^+\text{-C}^-(\text{CN})_2$ was refined by the same procedure as that employed for $\text{NH}_4\text{C}(\text{CN})_3$. The final value of *R*, based on 325 independent (*hkl*) data was 12%. The observed dimensions of the molecule are shown in Fig. 1. The molecule lies across a crystallographic mirror plane through atoms 1 and 4. The pyridinium ring as well as the trigonal carbon atom are coplanar within ± 0.01 Å. The two cyano groups are inclined with respect to this plane such that the distance from the plane to atoms 6 and 7 are 0.08 Å. and 0.13 Å., respectively. All distances reported in this molecule have an associated estimated standard deviation of ± 0.01 Å.

It thus appears that in both these systems a significant deviation from planarity of the carbanion group exists, even though the possibility of resonance stabilization of a planar configuration is possible. Detailed accounts of these results will be presented elsewhere.

Acknowledgment.—This work was supported by grants from the National Aeronautics and Space Administration and the Robert A. Welch Foundation.

DEPARTMENT OF CHEMISTRY
RICE UNIVERSITY
HOUSTON, TEXAS

CHARLES BUGG
ROBERT DESIDERATO
RONALD L. SASS

RECEIVED JUNE 12, 1964

Cubane

Sir:

We have completed the synthesis and fundamental characterization of the hydrocarbon cubane.

The bromocyclopentadienone dimer I, prepared as described in an earlier communication,¹ is converted

(1) P. E. Eaton and T. W. Cole, Jr., *J. Am. Chem. Soc.*, **86**, 962 (1964).