


Baeyer-Villiger oxidation of **10** utilizing trifluoro-peracetic acid gave the acetate **11** which reacted with methyllithium to give 4-hydroxy-*trans*-bicyclo[5.1.0]octane (**12**) in 64% yield based on **10**. *Anal.* Calcd for  $C_8H_{14}O$ : C, 76.14; H, 11.18. Found: C, 76.53; H, 11.32. The continued presence of the cyclopropyl moiety was demonstrated by near-infrared ( $1.643 \mu$  ( $\epsilon$  0.276)) and nmr spectroscopy (two-proton multiplet at  $\tau$  9.88). A comparison of **12** with **13** and **14** showed that these compounds had different ir and nmr spectra and different vpc retention times, indicating that these compounds were distinct entities. However, the mass spectra of **12**, **13**, and **14** were virtually identical, thus demonstrating the structural similarity between these three isomers. The nonidentity of **12** with both **13**



and **14** required that the cyclopropyl ring of **12** be *trans* fused,<sup>12</sup> and completed the structure proof.

The availability of this new class of strained *trans*-fused bicyclo[*n*.1.0]alkanes will permit a wide variety of theoretically interesting studies. We are currently investigating those aspects of *trans*-bicyclo[5.1.0]octane chemistry related to the unique nature of the strain incorporated into this molecule.

**Acknowledgment.** This research was supported by Grant GP7063 from the National Science Foundation.

(12) Compounds **13** and **14** were prepared by a modification of the published procedure: A. C. Cope, S. Moon, and C. H. Park, *J. Am. Chem. Soc.*, **84**, 4843 (1962).

(13) Alfred P. Sloan Research Fellow, 1967-1969.

(14) On leave from the Israel Ministry of Defense Scientific Department.

Paul G. Gassman,<sup>13</sup> Frank J. Williams, Jacob Seter<sup>1</sup>  
 Department of Chemistry, The Ohio State University  
 Columbus, Ohio 43210  
 Received September 3, 1968

## Additions and Corrections

**Linkage Isomerism in Phenylmercuric Benzenesulfinate** [*J. Am. Chem. Soc.*, **90**, 493 (1968)]. By G. B. DEACON and P. W. FELDER, Chemistry Department, Monash University, Clayton, Victoria, Australia.

The values for the sulfur-oxygen stretching frequencies of isomer A of phenylmercuric benzenesulfinate (the O-sulfinate) in chloroform (p 494) are incorrect. It has now been found that isomer A in chloroform reacts with sodium chloride plates, and the reported sulfur-oxygen stretching frequencies correspond closely to those of sodium benzenesulfinate. When silver chloride plates are used, the values for the chloroform solution (1053 and 850  $cm^{-1}$ ) are close to those (p 494) for solid isomer A (Nujol mull). This observation invalidates the conclusion (p 494) that isomer A is polymeric or dimeric (structure IV) in the solid state. It now appears probable that this isomer is a monomeric O-sulfinate complex (structure II or III) in the solid state, as in chloroform. It follows that the other  $RHg(SO_2R')$  derivatives (Table I, p 495) with sulfur-oxygen stretching frequencies similar to those of isomer A are likely to have a similar structure.

**The Dimerization of Styrene** [*J. Am. Chem. Soc.*, **90**, 1289 (1968)]. By FRANK R. MAYO, Stanford Research

Institute, Menlo Park, California 94025.

Reference 2 was inadvertently omitted. It should read: (2) F. R. Mayo, *J. Am. Chem. Soc.*, **75**, 6133 (1953).

**A Revised Structure for Ceanothine-B** [*J. Am. Chem. Soc.*, **90**, 4179 (1968)]. By ROBERT E. SERVIS and ALVIN I. KOSAK, Department of Chemistry, New York University, New York, New York 10003.

The second to last sentence should read: We have found that ceanothamine-A<sup>4</sup> is identical with frangulanine<sup>8</sup> (obtained from *Rhamnus frangula*) and that ceanothamine-B<sup>4</sup> is identical with adouetine-X<sup>2,8</sup> (isolated from *Waltheria americana*).

**Charge Distribution and Nucleophilic Reactivity in Sulfur Ligand Chelates. Dialkyl Derivatives of Nickel(II), Palladium(II), and Platinum(II) Bis(*cis*)ethylenedithiolates** [*J. Am. Chem. Soc.*, **90**, 4297 (1968)]. By G. N. SCHRAUZER and H. N. RABINOWITZ, Department of Chemistry, University of California at San Diego, Revelle College, La Jolla, California.

There is an error in Table VI. The corrected table is

Table VI. Observed and Calculated Bond Lengths (Å) in the Complexes  $NiS_4C_4R_4^{0,-2-}$  <sup>a</sup>

Ni-S		C-S		C-C		Compound	Ref
Obsd	Calcd	Obsd	Calcd	Obsd	Calcd		
2.101 (0.002)	<i>c</i>	1.71 (0.01)	1.70	1.37 (0.014)	1.39	$NiS_4C_4Ph_4$	25
2.146 (0.001)	<i>c</i>	1.71 (0.004)	1.72	1.35 (0.007)	1.37	$Ni(MNT)_2^-$	<i>b</i>
2.165 (0.005)	<i>c</i>	1.75 (0.01)	1.73	1.33 (0.02)	1.37	$Ni(MNT)_2^{2-}$	24

<sup>a</sup> Standard deviations in parentheses. <sup>b</sup> C. J. Fritchie, *Acta Cryst.*, **20**, 107 (1966). <sup>c</sup> Ni-S distances were not varied in this calculation.