

Table I. Relative Chemical Shifts and Dipolar Coupling Constants of **1** and **2** (All Values in Hertz)

Compd	$\nu_2 - \nu_3^a$	$\nu_2 - \nu_7$	$\nu_3 - \nu_7$	D_{23}^a	D_{24}	D_{25}	D_{34}	D_{27}	D_{37}	$D_{77'}$	Δf^b	$c, \text{mol} \%$
1	0.7	346.5	347.2	-450.4	-51.6	-20.4	-183.6	-50.1	-79.5	1795.0	0.65	16
2	-17.2 ^d			-393.5	14.1	36.6	330.1	-48.6	-48.2	455.1	0.63	13
	-21.7 ^e			-393.8	14.2	37.2	329.7					
	-17.3 ^e			-393.4	14.1	36.5	330.5					
	-11.9 ^e			-393.5	14.1	36.5	330.0					

^a Probable error of parameters <0.8 Hz. ^b Root-mean-square error for calculated and observed line positions. ^c Concentration. ^d Complete analysis. ^e Subspectral analysis.

Table II. r_{ij} Ratios and Orientation Parameters for **1** and **2**

Compd	r_{23}/r_{34}	r_{24}/r_{34}	r_{25}/r_{34}	r_{27}/r_{34}	r_{37}/r_{34}	$r_{77'}/r_{34}$	S_{xx}^a	S_{yy}^a	S_{zz}^a
1	1.028	1.771	2.081	1.669	2.214	0.760	0.0233	-0.1003	0.0770
	± 0.020	± 0.018	± 0.028	± 0.038	± 0.051	± 0.013	± 0.0001	± 0.0061	± 0.0061
2	1.024	1.769	2.081	1.741	2.293	0.879	-0.0419	-0.0487	0.0906
	± 0.006	± 0.006	± 0.015	± 0.028	± 0.030	± 0.055	± 0.0001	± 0.0021	± 0.0021
Benzene	1.000	1.732	2.000						

^a Based on an assumed H₃-H₄ distance of 2.48 Å.


been analyzed separately. The results for both compounds are collected in Table I.

Orientation parameters S_{kl} and the ratios of the various H,H, H,F, and F,F distances r_{ij} were then calculated from the experimental data of Table I using an iterative procedure.⁹ Since both compounds possess C_{2v} symmetry,¹⁰ two independent parameters are sufficient to characterize their orientation. For these calculations, the distance H₃-H₄ was held constant. The r_{ij} ratios are insensitive to this limitation.¹ To check the results obtained in this way, the ratios of the H,H distances in the six-membered ring of **1** and **2** have also been derived from explicit expressions given for the oriented AA'BB' system.¹¹ They were identical with those found by the iterative method. The results, together with the orientation parameters and the relevant r_{ij} ratios of benzene, are presented in Table II. From these values those based on H,F- and F,F-coupling constants in **2**, i.e., r_{27}/r_{34} , r_{37}/r_{34} , and $r_{77'}/r_{34}$, are less reliable, since the experimental D_{HF} and D_{FF} values contain unknown contributions from the anisotropy of the indirect spin-spin coupling.¹²

From the negative signs of the dipolar coupling constants it follows that **1**, as other aromatics,¹ orients with its x,z plane preferentially parallel to the external magnetic field. For **2**, however, a different orientation is indicated by the positive sign found for D_{24} , D_{34} , and D_{25} and, more directly, by the different S_{xx} value. In contrast to **1**, the y as well as the x axis of the molecular coordinate system orient preferentially perpendicular to the magnetic field direction.

Within the limits of error, our measurements yield identical proton geometry for the six-membered ring of **1** and **2**. Compared to benzene, all r_{ij}/r_{34} ratios have increased. This seems to indicate that the benzocyclopropene system suffers a distortion of type **3** with positive changes δ for the CCC bond angles at C₃ and C₄ and negative changes for these parameters

at C₂ and C₅. Similar angle distortions have been measured for 7,8-dichlorobenzocyclobutene¹³ and ben-



zocyclobutene-7,8-dione,¹⁴ where the strain effects should be comparable to those operating in **1** and **2**. Attempts to obtain the "best" carbon skeleton which fits into the geometry determined by the proton positions in both compounds using "molecular mechanics"¹⁵ are at present under way.

Acknowledgments. We wish to thank Dr. D. Schäpel and Professor Dr. E. Vogel, Cologne, for samples of **1** and **2**, Professor Dr. P. Diehl, Basel, for a copy of his program LAOCOONOR, and one of the referees for a helpful comment. This work was supported by the Deutsche Forschungsgemeinschaft and by the Fonds der Chemischen Industrie.

(13) G. L. Hardgrove, L. K. Templeton, and D. H. Templeton, *J. Phys. Chem.*, **72**, 668 (1968).

(14) F. H. Allen and J. Trotter, *J. Chem. Soc. B*, 916 (1970).

(15) F. H. Westheimer in "Steric Effects in Organic Chemistry," M. S. Newman, Ed., Wiley, New York, N. Y., 1956.

Josef Bernd Pawliczek, Harald Günther*

Institut für Organische Chemie, Universität Köln
Cologne, Germany

Pyrolysis of Cyclopropyl Azides. A Route to 1-Azetines¹

Sir:

Photolytic or pyrolytic decomposition of cyclopropyl azides with ring enlargement or substituent migration presents a potential route to 1-azetines or N-substituted cyclopropylimines. Neither the latter nor 2-alkyl- (or aryl-) 1-azetines have heretofore been reported.² In

(1) Stereochemistry. LXIII. For the previous paper in this series, see G. L'abbé and A. Hassner, *Angew. Chem.*, **83**, 103 (1971).

(2) The only 1-azetines in the literature have ether or thioether linkages at the 2 position and are obtained from β -lactams. See, for

(9) Details of the method will be given in the full paper.

(10) The magnetic equivalence of X₇ and X_{7'} in **1** and **2** demonstrates C_{2v} symmetry only on the nmr time scale, but it seems reasonable to assume that both compounds are planar.

(11) P. Diehl, C. L. Khatripal, and U. Lienhard, *Can. J. Chem.*, **46**, 2645 (1968).

(12) H. Spiesecke and A. Saupe, *Mol. Cryst. Liq. Cryst.*, **6**, 287 (1970).


Table I. Yields and Physical Data in the Formation and Pyrolysis of Cyclopropyl Azides 2

Azide	% yield (isolated)	Nmr spectra, τ^a			Bp, °C (mm)	Pyrolysis rate, $k \times 10^{-4}$ sec ⁻¹
		R ₁	R ₂	R ₃		
2a	42	2.60 (s)	7.85, (d, $J = 8.0$)	8.09 (d, $J = 8.0$)	83-85 (0.1)	4.6
<i>cis</i> - 2b	53	2.57 (s)	8.80 (d, $J = 6.5$)	7.99 (q, $J = 6.5$)	90 (0.2)	14.7
<i>cis</i> - 2c	25	8.50 (s)	8.81 (d, $J = 6.5$)	8.30 (q, $J = 6.5$)	64 (7)	8.2
<i>trans</i> - 2d	66	2.57 (s)	7.89 (q, $J = 6.5$)	8.60 (d, $J = 6.5$)	82 (0.1)	2.9
<i>trans</i> - 2e	52	8.27 (s)	8.61 (q, $J = 4.5$)	8.80 (d, $J = 4.5$)	65 (7)	1.8

^a s = singlet, d = doublet, q = quartet; J values in hertz.

fact, since cyclopropyl halides are neither readily susceptible to displacement reactions nor to solvolysis without rearrangement, cyclopropyl azides have remained a virtually unknown class of compounds.³

Our recent discovery of a general synthesis of vinyl azides from olefins⁴ suggested their reaction with dichlorocarbene as a route to **2**. *N*-Octyl azide has been reported to react with dichlorocarbene to give the corresponding isocyanide dichloride.⁵ In sharp contrast, the vinyl azides **1** gave solely the corresponding 1-azido-2,2-dichlorocyclopropanes **2** in moderate yields (see Table I). This is in analogy with recent studies which showed that vinyl azides react at carbon with other electrophiles such as acid⁶ and bromine.⁷ In a typical experiment 21.9 g (0.115 mol) of ethyl trichloroacetate was added dropwise over a period of 30 min to a solution of 0.1 mol of the vinyl azide and 12.8 g (0.2 mol) of sodium methoxide in 100 ml of pentane at 0° under nitrogen. After stirring for 20 hr at room temperature the mixture was filtered through Celite. Removal of the pentane *in vacuo* left an oil containing starting material and azide. Purification of the latter was accomplished by careful chromatography followed by distillation below 100° at reduced pressure. (*Caution:* these compounds explode on heating at *ca.* 110°.)



Azides **2a-c** decompose smoothly between 105 and 125° in a highly regiospecific⁸ manner to give the corresponding 3,3-dichloroazetines **3** in 50-75% isolated yields (see Table II). No spectral evidence for formation of **4** or **5** was observed. The primary side

example, G. Pifferi, P. Consonni, G. Pelizza, and E. Testa, *J. Heterocycl. Chem.*, **4**, 619 (1967); D. Bormann, *Justus Liebig's Ann. Chem.*, **752**, 124 (1969). For another recent example, see A. Hassner, J. O. Currie, A. S. Steinfeld, and R. F. Atkinson, *Angew. Chem., Int. Ed. Engl.*, **9**, 731 (1970).

(3) Cyclopropyl azide itself has been reported by W. Kirmse and H. Schutte, *J. Amer. Chem. Soc.*, **89**, 1284 (1967), and W. Kirmse and H. Schutte, *Chem. Ber.*, **101**, 1674 (1968).

(4) F. W. Fowler, A. Hassner, and L. A. Levy, *J. Amer. Chem. Soc.*, **89**, 2077 (1967); A. Hassner, *Accounts Chem. Res.*, **4**, 9 (1971).

(5) J. E. Baldwin, *Chem. Commun.*, 968 (1968).

(6) H. W. Moore, H. R. Shelden, D. W. Deters, and R. J. Wikholm, *J. Amer. Chem. Soc.*, **92**, 1675 (1970); A. Hassner, E. S. Ferdinandi, and R. J. Isbister, *ibid.*, **92**, 1672 (1970).

(7) A. Hassner and A. B. Levy, *ibid.*, in press.


(8) Regio is used to describe a directional preference in bond making or breaking: A. Hassner, *J. Org. Chem.*, **33**, 2684 (1968).

Table II. Yields and Nmr Data of Azetines 3


Azetine	% yield	Nmr spectra, τ^a
3a	55	5.49 (s, 2 H) 2.51, 2.05 (2 m, 5 H)
3b	72	8.49 (d, 3 H, $J = 7$) 5.40 (q, 1 H, $J = 71$) 2.60, 2.03 (2 m, 5 H)
3c	75	8.59 (d, 3 H, $J = 6.5$) 7.91 (d, 3 H, $J = 1.6$) 5.56 (q of q, 1 H, $J = 1.6$ and 6.5)

^a s = singlet, d = doublet, q = quartet, m = multiplet; J values in hertz.

products appear to be the nitrile **6** and the olefin **7** (*ca.* 5%), possibly resulting from the fragmentation of the azetine ring. Typically, 5 g of the azide was




refluxed in 200 ml of toluene for 9 hr, the toluene removed *in vacuo*, and the resultant azetine chromatographed on silica gel.



The structures of **3a-c** were indicated by ir, nmr, and mass spectra. The mass spectra showed strong peaks corresponding to the molecular ion, the nitriles **6**, and the olefins **7**. Conclusive evidence for the structures came from acid hydrolysis and LiAlH₄ reduction to give the analogous β -amino- α,α -dichloro ketones **8** and azetidines **9**, respectively.

The structures of the amino ketones **8a-c** were indicated by spectral data, the mass spectra being especially helpful since they showed the expected amine α cleavage corresponding to R₂R₃C=NH₂⁺ and no α cleavage at *m/e* 98, 100, and 102, corresponding to CCl₂=NH₂⁺. The crude nmr spectra of **8a-c** indicated



no other products present. The structures of azetidines **9a-c** were obvious from their ir, nmr, and mass spectra. That the geminal hydrogens were in the 3 position in **9a** and **b** was indicated by the low-field (τ 5.4) broad triplet for H_a . In **9c** H_a could not be resolved, but the mass spectra showed the expected α cleavage at m/e 43 ($CH_3CH=NH\cdot^+$) and no peak at m/e 30 ($CH_2=NH\cdot^+$).

The regioselectivity in the formation of **3** over **4** and **5** cannot be equated with conformational effects⁹ governing a concerted loss of N_2 with ring enlargement, since the conformers of **2** are expected to be of nearly equal stability and should produce a mixture of **3**, **4**, and **5**.

The driving force for ring expansion in preference to formation of a cyclopropylimine **5** (Ph migration) must be relief of ring strain in going from a three- to a four-membered ring. Apparently, the reaction involves selective migration of the electron richer CH_2 (or CR_2) rather than the CCl_2 ring residue to the electron-deficient nitrogen atom.

Rate data on the pyrolysis of cyclopropyl azides **2** (Table I) indicate that in general cis substituents (see **2b** and **2c**) accelerate the reaction. This implies that the rate-determining step cannot be loss of N_2 unless the reaction is concerted. A concerted ring expansion with loss of nitrogen is in fact suggested by the negative entropy of activation ($\Delta S^\ddagger = -18$ eu for **2c** \rightarrow **3c**) and is in accord with data on pyrolysis of simple alkyl azides.¹⁰

Acknowledgment. Support of this work by a grant (CA 04474) from the National Cancer Institute (Public Health Service) is gratefully acknowledged.

(9) Different migratory aptitudes in the photolysis of organic azides have been ascribed to ground-state conformational preferences; see R. M. Moriarty and R. C. Reardon, *Tetrahedron*, **26**, 1379 (1970).

(10) W. Lwowski, "Nitrenes," Interscience, New York, N. Y., 1970, Chapters 3 and 4.

(11) NIH Predoctoral Fellow, 1968-1971.

A. B. Levy,¹¹ A. Hassner*

Department of Chemistry, University of Colorado
Boulder, Colorado 80302

Received December 5, 1970

Effect of CH Scalar Coupling on ^{13}C Transverse Relaxation Times

Sir:

In view of the increasing interest in ^{13}C nmr spectroscopy, we wish to report some preliminary relaxation studies of ^{13}C -enriched compounds. We find that the

transverse relaxation times, T_{2C} ($\equiv R_{2C}^{-1}$), are usually shorter than the longitudinal relaxation times, T_{1C} ($\equiv R_{1C}^{-1}$). This result influences the sensitivity and resolution in both conventional and Fourier transform (FT) ^{13}C spectra and, in particular, affects the signal enhancement obtainable by the use of refocusing schemes such as driven equilibrium and spin-echo Fourier transform (DEFT and SEFT) techniques.¹⁻³

The T_{2C} measurements were performed at 15.08 MHz by the usual Meiboom-Gill⁴-modified Carr-Purcell⁵ technique on a high-resolution magnet with external fluorine lock. The π pulse width was 6 μ sec and the inhomogeneity over the sample gave a free-induction decay constant $T_2^* \sim 100$ msec. Degassed and sealed sample tubes were constricted below the liquid surface to minimize interchange between liquid and vapor. T_{1C} and T_{1H} values were measured by π - τ - $\pi/2$ pulse sequences, T_{1C} 's at 15.08 MHz, T_{1H} 's at 23.2 MHz. The T_1 and T_2 values were obtained by the usual least-squares fitting procedures.

Table I shows that T_{2C} is substantially shorter than

Table I. Relaxation Times^a for 60% ^{13}C -Enriched Compounds at 25°

	T_{1C} , sec	T_{2C} , sec	T_{1H} , ^b sec
$^{13}CH_3I$	13.4	3.9	10.9 ^c
$^{13}CH_3COOCD_3$ ^d	19.2	6.1	12.5
$^{13}CS_2$	48 ^e	38	

^a The experimental error is estimated to be $\sim 5\%$. ^b T_{1H} for 100% enriched compounds will be less than these values. ^c T_{1H} for $^{12}CH_3I$ at 25° is reported [M. D. Zeidler, *Ber. Bunsenges. Phys. Chem.*, **69**, 659 (1965)] to be 12.5 sec. ^d Deuterium enrichment is 99%. ^e $T_{1C} = 30$ sec (15 MHz, 26-28°) was measured by rapid adiabatic passage; A. Olivson, E. Lippmaa, and J. Past, *Eesti NSV Tead. Akad. Toim.*, **16** 390 (1967).

T_{1C} ⁶ for methyl iodide and methyl- d_3 acetate- ^{13}C . We would like to point out that short T_{2C} 's occur when relaxing spin- $1/2$ nuclei, e.g., protons, are coupled to ^{13}C such that

$$A \equiv 2\pi J \gg R_{1H} \quad (1)$$

where $R_{1H} \equiv 1/T_{1H}$. Gutowsky, Vold, and Wells⁸ have pointed out the analogy between chemically exchanging, equally populated, uncoupled two-site systems and relaxing, scalar-coupled AX systems. They suggest that the two cases are related by the simple substitution of $\Omega \rightarrow A/2$ and $k \rightarrow R_{1X}$, where Ω is half the chemical shift between the two sites, k the exchange rate, and R_{1X} the spin-lattice relaxation rate of X.

Scalar contributions to R_{2A} for a general AX_n system originate from relaxation processes within the X manifold which preserve the spin state of A while changing the scalar coupling energy by a nonzero multiple of J . These processes randomly alter the precession frequency

(1) E. D. Becker, J. A. Ferretti, and T. C. Farrar, *J. Amer. Chem. Soc.*, **91**, 7784 (1969).

(2) A. Allerhand, *ibid.*, **92**, 4482 (1970).

(3) J. S. Waugh, *J. Mol. Spectrosc.*, **35**, 298 (1970).

(4) S. Meiboom and D. Gill, *Rev. Sci. Instrum.*, **29**, 688 (1958).

(5) H. Y. Carr and E. M. Purcell, *Phys. Rev.*, **94**, 630 (1954).

(6) We realize that T_1 is not rigorously defined for strongly coupled AX_n systems (see, e.g., ref 7, p 295) but we could detect no deviation from exponential behavior.

(7) A. Abragam, "The Principles of Nuclear Magnetism," Oxford University Press, London, 1961, p 311.

(8) H. S. Gutowsky, R. L. Vold, and E. J. Wells, *J. Chem. Phys.*, **43**, 4107 (1965).