

Table I. Aluminum Amalgam Reductions^a

Reaction	Conditions	Yield, %	Starting material [α] ^{25D} (acetone); mp, °C	Product [α] ^{25D} (acetone); mp, °C
1 → 4 ^b	15°, 3 hr	74	+183°, liq	+173°, 49–52
2 → 5 ^b	20°, 2 hr	54	+36.5°, 31–33	+79.9°, 95–101 ^c
6 → 7	25°, 2 hr	41	+43.7°, 99–100	+46.4°, ^d liq
8 → 9 ^e	25°, 24 hr	53	+12.6°, 128–130	-142.6°, ^f liq
10 → 4 ^b	20°, 4 hr	76	Rac, 87–89	Rac, liq
11 → 4 ^b	25°, 4 hr	46	Rac, liq	Rac, liq

^a The amalgam was prepared from commercial grade aluminum foil by immersing it into 2% aqueous mercuric chloride for 15–20 sec, followed by ethanol and ether rinses, and it was used immediately (ref 4). Generally, 10 g-atom of aluminum/mol of compound was used. The reactions were followed by tlc. All compounds have been identified and characterized by standard analytical procedures. ^b A small amount of benzenethiol was produced in these reactions by further reduction of the sulfinamide. The products were purified by chromatography on silica gel. ^c After recrystallization from ether, [α]^{25D} +82.9°, mp 102–103°. ^d Lit. +43.7° (neat) [R. L. Burwell, Jr., A. D. Shields, and H. Hart, *J. Amer. Chem. Soc.*, **76**, 908 (1954)]; authentic sample, +44.0° (neat), +47.1° (acetone). ^e In this reaction, 5 g-atom of Al/mol of **10** was used. ^f Lit. for enantiomer +149° [J. Jacobus and K. Mislow, *J. Amer. Chem. Soc.*, **89**, 5228 (1967)].

The aluminum amalgam reduction of (+)-(*S*)-*N,N*-dimethyl-*S*-phenylsulfoximine (**1**)^{6a} resulted in the cleavage of the sulfur-alkyl bond to give (+)-(*S*)-*N*-methylbenzenesulfinamide (**4**). The absolute configurations of **1** and **4** have been established;⁷ the reduction proceeds with retention of configuration at the chiral sulfur. Sulfur-alkyl bond cleavage was also observed with (+)-(*S*)-*S*-methyl-*S*-phenylsulfoximine (**2**)^{6b} which yielded (+)-(*S*)-benzenesulfinamide (**5**).⁸ At first glance it may appear that a method for the transformation of a sulfoximine to a sulfinamide would offer little other than a new and useful method to correlate configurations. However, it should be noted that optically active primary sulfinamides have not been previously reported, few optically active secondary sulfinamides are known, and the synthesis of these materials in high optical purity is difficult to achieve due to racemization under the more usual reaction conditions.⁹ The mild media of the aluminum amalgam reductions of the easily resolved sulfoximines allow high retention of optical activity in the production of sulfinamides.


The Al(Hg) reduction of sulfoximines is a key step in a general reaction sequence under development in our laboratory to produce optically pure alcohols. For example, reduction of sulfoximine **6** produced optically pure (+)-(*R*)-1-phenylethanol (**7**). Under these mild conditions no hydrogenolysis or racemization occurred at the sensitive benzylic carbon. Analogous β -hydroxysulfones are inert to these reduction conditions. Employing slightly higher temperatures

(6) (a) Prepared by methylation of resolved **2**, [α]^{25D} +36.5° (c 1.2, acetone). (b) For the preparation of this sulfoximine see R. Fusco and F. Tericoni, *Chem. Ind. (Milan)*, **47**, 61 (1965), and C. R. Johnson, M. Haake, and C. W. Schroeck, *J. Amer. Chem. Soc.*, **92**, 6594 (1970). A forthcoming paper from our laboratory will describe an improved procedure for the resolution of this compound.

(7) E. U. Jonsson and C. R. Johnson, *ibid.*, **93**, 5308 (1971).


(8) Absolute configuration based on the correspondence of the sign of rotation with that of **4** and the assumption that the reduction proceeded with retention of configuration at sulfur.

(9) A. Nudelman and D. J. Cram, *J. Amer. Chem. Soc.*, **90**, 3869 (1968).




and a greater excess of the aluminum, β -hydroxy sulfoxides are quantitatively reduced to the corresponding sulfides. *N-p*-Toluenesulfonylsulfoximine (**3**) proved resistant to reduction under these mild conditions.

An interestingly different mode of cleavage occurs in the reduction of salts prepared by *N,N*-dimethylation of sulfoximines. When (+)-(*S*)-(dimethylamino)-methylphenyloxosulfonium fluoroborate (**8**) was treated with the metal amalgam, the dimethylamino group was lost with retention of configuration at sulfur to give (-)-(*S*)-methyl phenyl sulfoxide (**9**); some methyl phenyl sulfide was also isolated. This represents an alternative to the use of nitrosyl hexafluorophosphate for the removal of the nitrogen from a sulfoximine.¹⁰



Other sulfonimidoyl compounds which were subjected to Al(Hg) reduction included phenyl *N*-methylbenzenesulfonyl imidate (**10**)¹¹ and *N,N,N'*-trimethylbenzenesulfonyl imidamide (**11**);¹¹ in each case sulfinamide **4** was produced.



We are continuing to investigate the scope and mechanistic details of these reductions.

(10) D. J. Cram, J. Day, D. R. Rayner, D. M. von Schrittz, D. J. Duchamp, and D. C. Garwood, *ibid.*, **92**, 7369 (1970).

(11) Prepared by the methods described for the optically active compounds in ref 7.

Calvin W. Schroeck, Carl R. Johnson*

Department of Chemistry, Wayne State University
Detroit, Michigan 48202

Received July 6, 1971

Preparation of Sulfonimidoyl Chlorides by Chlorination of Sulfinamides¹

Sir:

Derivatives of "sulfonimidic acids" (**1**) are a relatively new class of compounds² which, in comparison

(1) Part XXXII in the series "Chemistry of Sulfoxides and Related Compounds." We gratefully acknowledge support by the National Science Foundation (GP 19623).