

brated under the reaction conditions with partial consumption of **5c** (see last line of Table I). This is the likely reason that reaction of $\text{Me}_2\text{N}\cdot$ with 6/94 (thermodynamically equilibrated) *cis/trans* **5a** gives a poor correspondence between isomeric ratios of reacted **5a** and product **5c** as well as a poor yield of **5c**. The dominant *trans*-**5a** is less reactive than is *cis*-**5a**, hence dimethylamino radicals react rapidly with *cis*-**5c** leading finally to its equilibration with *trans*-**5c**. Note, however, that the *trans/cis* ratio for product **5c** from reaction of 6/94 **5a** is reduced as expected for an inversion pathway.

When **5b** is the reactant, the minor *trans*-isomer is the thermodynamically less stable form⁸ and the more reactive isomer toward $\text{Me}_2\text{N}\cdot$. The data again argue for *inversion about phosphorus as the predominant pathway*. Yields of $\text{PhCH}_2\text{CH}_2\text{Ph}$ as high as 80% attest to the free-radical nature of the displacement on **5b**.

We interpret these results (inversion) to mean that if a phosphoranyl radical is involved in these processes, its geometry must be such that X and Y are either coaxial or coequatorial. The attacking radical $\text{Me}_2\text{N}\cdot$ clearly is not introduced into the phosphoranyl intermediate in a configurationally random manner, a result consistent with earlier conclusions drawn from investigations of reactions of alkoxy radicals with trialkyl phosphites.² If as assumed above, $\text{X}\cdot$ enters the intermediate axially, and the odd electron is equatorial, then structure **4** will give the observed stereochemistry.

Since the alkyl substituents on phosphorus are relatively electropositive and according to esr results³ appear to exhibit an equatorial preference in the most thermodynamically stable form of intermediates such as $\text{Me}_3\text{P}\cdot\text{-}t\text{-Bu}$ and $\text{Me}_2\text{P}\cdot(\text{OR})_2$, kinetic factors may favor formation of **4**. Alternatively, the stereochemistry of these systems may be controlled by the presence of the six-membered ring which in pentacoordinate systems has been suggested¹¹ to have a thermodynamic preference for diequatorial attachment to phosphorus. Finally, the particular systems investigated may react *via* a concerted (synchronous SH2) mechanism. These possibilities are being investigated.

Acknowledgment. This research was generously supported by grants from the National Cancer Institute of Public Health Service (CA-11045) and by the National Science Foundation (GP-36637X1).

(11) P. Gillespie, P. Hoffman, H. Klusacek, D. Marquarding, S. Pfohl, F. Ramirez, E. A. Tsolis, and I. Ugi, *Angew. Chem., Int. Ed. Engl.*, **10**, 687 (1971).

Wesley G. Bentrude,* Wajid A. Khan
Masashi Murakami, Han-Wan Tan
Department of Chemistry, University of Utah
Salt Lake City, Utah 84112


Received April 20, 1974

Direct Determination of the Enthalpy Change for the Migration of Para-Substituted Benzyl and Trifluoromethyl Groups in Coordinatively Unsaturated Iridium(III) Acyl Complexes


Sir:

The insertion of carbon monoxide into a metal-carbon bond or its reverse, the migration of an alkyl or aryl group from an acyl carbon to a metal site in an acyl complex, are of widespread occurrence in organo-

metallic syntheses and homogeneous catalytic processes.¹ A reaction sequence which accounts for what is known of the mechanism for these transformations is shown in




eq 2.¹ In general the intermediate of reduced coordi-



nation number, $\text{L}_{n-1}\text{MC(O)R}$, is not detected and the actual migration (or insertion) step is not observed independent of the loss (or addition) of a ligand from the metal coordination sphere. It is not surprising, then, that no thermochemical data have yet been reported for the reaction step in which the alkyl or aryl group migrates to or from the coordinated CO group.

In 1971 Kubota, Blake, and Smith reported the synthesis of a series of five-coordinated phenylacetyl complexes of iridium(III).^{2a} These compounds were used as models for the coordinatively unsaturated intermediate in eq 2, and the rate of the migration step was studied.² We now wish to report that these five-coordinated phenylacetyl complexes, as well as the analogous trifluoroacetyl complex,^{2b} undergo the migration reaction in the solid state. This has made it possible to determine the enthalpy change, ΔH_T , at the reaction temperature, T , and at standard conditions, ΔH_0 .

The reaction studied is shown in eq 3. When the



orange solid, Ia-d, is heated to a temperature, T , which depends on the R group, the migration reaction occurs to give white compounds IIa-d. These were shown to be identical with the products formed in homogeneous solution by ir spectroscopy. The reaction was not complicated by weight loss and the exothermic peak for the migration reaction was well separated from the melting transition of the product. The enthalpy change for the reaction was determined at temperature T by standard differential calorimetric methods.^{3,4} These

(1) A. Wojcicki, *Advan. Organometal. Chem.*, **11**, 87 (1973).

(2) (a) M. Kubota, D. M. Blake, and S. A. Smith, *Inorg. Chem.*, **10**, 1430 (1971); (b) D. M. Blake, S. Shields, and L. Wyman, *ibid.*, **13**, 1595 (1974).

(3) A Perkin-Elmer Model DSC-1B differential scanning calorimeter was used in the study. Peak areas for the reaction were calibrated with respect to the heat of fusion of standard indium samples. A sapphire sample was used as a standard for the heat capacity determinations.

(4) (a) W. J. Smothers and Y. Chiang, "Handbook of Differential Thermal Analysis," Chemical Publishing Co., New York, 1966; (b) Instructions for Differential Scanning Calorimetry, Perkin-Elmer Corporation Publication No. 990-9556 (1966); (c) M. J. O'Neill, *Anal. Chem.*, **38**, 1331 (1966).

Table I. Thermodynamic and Kinetic Parameters for an Alkyl Group Migration in Iridium(III) Complexes
$$\{\text{IrCl}_2[\text{C}(\text{O})\text{R}][\text{P}(\text{C}_6\text{H}_5)_3]_2\}_{\text{solid}} \longrightarrow \{\text{IrCl}_2(\text{CO})\text{R}[\text{P}(\text{C}_6\text{H}_5)_3]_2\}_{\text{solid}}$$

R	T^a °K	ΔH_T^b kJ mol ⁻¹	Standard deviation of ΔH_T (kJ mol ⁻¹)	$\int_{298}^T \Delta C_p dT$ (kJ mol ⁻¹)	ΔH_0 kJ mol ⁻¹	$t_{1/2}$ (in solution) (min)	ΔH^\ddagger (in solution) (kJ mol ⁻¹)
Ia, CF ₃	445	-83.88	3.8	0.08	-84 ± 4	44.59 ^d (at 343°K)	+86.5
Ib, <i>p</i> -NO ₂ C ₆ H ₄ CH ₂ -	444	-16.40	1.30	0.01	-16 ± 1	29.67 ^{d,e} (at 303°K)	+84.5 ^d
Ic, <i>p</i> -CH ₃ C ₆ H ₄ CH ₂ -	428	-14.39	1.51		-14 ± 2 ^c	13.83 ^{d,e} (at 303°K)	+68.2 ^d
Id, <i>p</i> -CH ₃ OC ₆ H ₄ CH ₂ -	419	-13.34	3.81		-13 ± 4 ^c	12.57 ^{d,e} (at 303°K)	+62.8 ^d

^a Temperature of peak maximum. ^b Each value is the average of 12 determinations. ^c $\Delta H_0 = \Delta H_T$ by analogy with compounds Ia and Ib. ^d Ia in toluene, Ib, c, and d, in benzene. ^e M. Kubota, D. Blake, and S. Smith, *Inorg. Chem.*, **10**, 1430 (1971).

Table II. Heat Capacity Data, $C_p = (a + bT + cT^2 + dT^3)$ (4.184 J °K⁻¹ mol⁻¹)

Compound ^a	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>	$\int_{298}^T C_p dT$ (kJ mol ⁻¹)	<i>T</i> ^b
IrCl ₂ [C(O)CF ₃]L ₂	179.49	1.8279 × 10 ⁻¹	-2.0533 × 10 ⁻³	4.83 × 10 ⁻⁶	131.67	445°K
IrCl ₂ (CO)CF ₃ L ₂	128.26	-1.2987	7.5246 × 10 ⁻³	9.02 × 10 ⁻⁶	131.75	
IrCl ₂ [C(O)CH ₂ C ₆ H ₄ NO ₂]L ₂	-39.686	1.3955	-2.8362 × 10 ⁻³	3.01 × 10 ⁻⁶	145.81	444°K
IrCl ₂ (CO)(CH ₂ C ₆ H ₄ NO ₂)L ₂	-71.013	2.4474	-779.67 × 10 ⁻³	9.29 × 10 ⁻⁶	145.82	


^a L = P(C₆H₅)₃. ^b T = temperature at which the migration reaction occurs.

data are shown in Table I. In order to determine the enthalpy change at standard conditions, ΔH_0 , the following relation was used

$$\Delta H_0 = \Delta H_T - \int_{298}^T \Delta C_p dT$$

Molar heat capacities, C_p , were determined for the pairs of compounds Ia and IIa and Ib and IIB using differential scanning calorimetry.^{4c} The data for C_p as a function of temperature were fitted to a third degree polynomial equation in T from 298° to the reaction temperature for each compound. The integrals were then evaluated using Simpson's rule.⁵ These data are shown in Table II. The magnitude of the integral term for ΔC_p was much smaller than the standard deviation in ΔH_T so that ΔH_0 may be assumed to be equal to ΔH_T . These data are shown in Table I, along with kinetic data for the reaction in homogeneous solution.

The magnitude of the enthalpy change is smaller than metal-carbon monoxide bond energies (*ca.* 120–170 kJ mol⁻¹)⁶ or metal-carbon bond energies (*ca.* 150–300 kJ mol⁻¹).⁷ The enthalpy change is opposite in sign from that found for the decarbonylation of benzoyl platinum complexes (eq 4) for which $\Delta H = +18.6$ to



From the data it can be noted that there is an inverse correlation between the migration rate and the enthalpy change. The enthalpy change in solution is anticipated to be close in value to that in the solid state since heats of solution of the compounds I and II are likely to be small and of similar magnitude in nonpolar solvents. The large difference in the magnitude of ΔH_0 for the tri-

fluoromethyl, -83 kJ mol⁻¹, and benzyl groups, *ca.* -15 kJ mol⁻¹, is noteworthy. The enthalpy change for the migration reaction includes four bond energy terms (Ir-C≡O, Ir-C(O)R, Ir-R, and C(O)-R). Unfortunately there is not at present sufficient information available about the individual contributions to make a meaningful discussion of the difference in the overall values of ΔH possible. A kinetic and calorimetric study is now in progress for the series of five-coordinated complexes analogous to I where R = CF_{*n*}H_{3-*n*} (*n* = 3, 2, 1, or 0). Preliminary results show that the enthalpy change for reaction 3 decreases in the series CF₃ > CF₂H > CFH₂ > CH₃ ≈ *p*-XC₆H₄CH₂. Other work is in progress to extend the data to other migrating groups and to seek correlations of the enthalpy changes with kinetic and spectral parameters of the complexes.

Acknowledgment. The authors thank the Robert A. Welch Foundation and University Organized Research Fund for support of this work.

(9) (a) Robert A. Welch Foundation Predoctoral Fellow. (b) Supported in part as a Robert A. Welch Foundation Undergraduate Fellow and by funds from the University Organized Research Fund.

Daniel M. Blake,* Joseph de Faller
Yen Lung Chung,^{9b} Alan Winkelman^{9b}

Department of Chemistry, The University of Texas at Arlington
Arlington, Texas 76019

Received May 15, 1974

MINDO/3 Study of the Bisdehydrobenzenes

Sir:

While the three bisdehydrobenzenes **1**, **2**, and **3** were characterized some years ago by Berry, *et al.*,¹⁻³ while the ortho isomer **1** (benzyne) has been proposed as an

(1) R. S. Berry, G. N. Spokes, and M. Stiles, *J. Amer. Chem. Soc.*, **84**, 3570 (1962).

(2) R. S. Berry, J. Clardy, and M. E. Shafer, *Tetrahedron Lett.*, 1003 (1965).

(3) R. S. Berry, T. Clardy, and M. E. Shafer, *Tetrahedron Lett.*, 1101 (1965).

(5) D. D. McCracken and W. S. Dorn, "Numerical Methods and Fortran Programming," Wiley, New York, N. Y., 1968.

(6) H. A. Skinner, *Advan. Organometal. Chem.*, **2**, 49 (1964).

(7) G. Wilkinson, *Pure Appl. Chem.*, **30**, 627 (1972).

(8) S. J. Ashcroft, A. Maddock, and G. Beach, *J. Chem. Soc., Dalton Trans.*, 462 (1974).