

material from this paper only or microfiche (105 × 148 mm, 24× reduction, negatives) containing all of the supplementary material for the papers in this issue may be obtained from the Journals Department, American Chemical Society, 1155 16th St., N.W., Washington, D. C. 20036. Remit check or money order for \$3.00 for photocopy or \$2.00 for microfiche, referring to code number JACS-74-6785.

(13) Camille and Henry Dreyfus Foundation Teacher-Scholar Grant Awardee 1972-1977 and Fellow of the Alfred P. Sloan Foundation 1973-1975.

Jose Fayos, David Lokensgard, Jon Clardy*¹⁸
Ames Laboratory-USAEC and Department of Chemistry
Iowa State University
Ames, Iowa 50010

Richard J. Cole, Jerry W. Kirksey
National Peanut Research Laboratory
U. S. Department of Agriculture
Dawson, Georgia 31742
Received June 26, 1974

[CF₂Cl⁻], Chlorodifluoromethide Ion. The Capture of an Elusive Species

Sir:


The concerted nature of difluorocarbene formation *via* the decomposition of chlorodifluoroacetic acid and its derivatives is presently accepted as there is no concrete evidence for the existence of the chlorodifluoromethide ion in the literature to date. Herein is reported evidence supporting the existence of the chlorodifluoromethide ion as a reactive intermediate in the decarboxylation of methyl chlorodifluoroacetate. Independent experiments show that the observed products do not result from the initial formation and subsequent reactions of difluorocarbene.

The usefulness of trichloroacetic acid and its derivatives as dichlorocarbene precursors has been known for years.¹⁻⁴ More recently, this interest has been extended to the potential use of chlorodifluoroacetic acid and its derivatives as synthetically useful sources of difluorocarbene. Alkali metal chlorodifluoroacetates have been used in the synthesis of steroid derivatives⁵ and *gem*-difluorocyclopropanes,⁶ in homologation reactions,⁷ and in the preparation of 1,1-difluoroolefins.⁸ Sodium ethoxide induced decomposition of ethyl chlorodifluoroacetate resulted in difluoromethylation of 2,3-dimethylindole.⁹ Decomposition of lithium chlorodi-


fluoroacetate in the presence of 9,10-phenanthrene quinone gave a hydroxy ketone which might possibly be a chlorodifluoromethide ion trapping product.¹⁰ However, since initial studies by Hine and coworkers, it has been accepted that halodifluoromethide ions have no finite existence. In studies of both the hydrolysis of chlorodifluoromethane¹¹ and the decarboxylation of chlorodifluoroacetic acid,¹² it was concluded that difluorocarbene formation is a concerted process and that there is no intermediate formation of halodifluoromethide ion. Attempts to trap these carbanions even by protonation have met with little success.¹²

We have observed in this laboratory that lithium chloride initiates the facile decarboxylation of methyl chlorodifluoroacetate (I) in hexamethylphosphoramide (HMPA). A study of this decarboxylation in the presence of polyfluorinated ketones has been done, and the results are reported below.

Decarboxylation of I by lithium chloride in the presence of trifluoroacetophenone (II) in HMPA proceeded smoothly at 65 to 70°C.¹³ After 20 hr, steam distillation resulted in isolation of 1-chloro-2-phenylpentafluoro-2-propanol (III) as the only product in 62% yield. Similarly, decarboxylation of I in the presence of 1,1,1-trifluoro-2-hexanone (IV) afforded 1-chloro-1,1-difluoro-2-trifluoromethyl-2-hexanol (V) as the only product in 39% yield. Decarboxylation of I in the presence of



chlorodifluoroacetophenone (VI), however, yielded an olefin, 1-chloro-1-phenyldifluoroethylene (VII), as the major product (50%) as well as 1,3-dichloro-2-phenyltetrafluoro-2-propanol (VIII) (18%) after steam distillation.



Two mechanistic interpretations have been considered to explain the observed results. The first interpretation is that the products in each case result from initial formation of the chlorodifluoromethide ion upon decarboxylation of I with subsequent attack of the carbanion at the carbonyl carbon atoms of ketones II, IV, and VI to form alcoholates IX, X, and XI, respectively (Scheme I). For reasons as yet unexplained, alcoholate XI can undergo an intramolecular S_N2 displacement of chloride ion by oxygen to form oxirane

(10) M. Derenberg and P. Hodge, *J. Chem. Soc., Perkin Trans.*, 1056 (1972).

(11) J. Hine and P. B. Langford, *J. Amer. Chem. Soc.*, **79**, 5497 (1957).

(12) J. Hine and D. C. Duffey, *J. Amer. Chem. Soc.*, **81**, 1131 (1959).

(13) In one experiment methyl chloride was detected and identified by ¹H nmr and infrared spectroscopy. Quantitative analysis by ¹H nmr showed a 92% yield of methyl chloride. A yield of 88% CO₂ was determined by precipitation as barium carbonate.

(1) F. H. Verhoek, *J. Amer. Chem. Soc.*, **56**, 571 (1934).

(2) W. E. Parham and F. C. Loew, *J. Org. Chem.*, **23**, 1705 (1958).

(3) W. M. Wagner, *Proc. Chem. Soc., London*, 229 (1959).

(4) W. M. Wagner, *et al.*, *Recl. Trav. Chim. Pays-Bas*, **80**, 740 (1961).

(5) See, for example, L. H. Knox, E. V. Velarde, S. M. Berger, and D. H. Caudriello, *Chem. Ind. (London)*, 860 (1962); L. H. Knox, E. V. Velarde, S. M. Berger, D. H. Caudriello, P. W. Landis, and A. D. Cross, *J. Amer. Chem. Soc.*, **85**, 1851 (1963); C. Beard, N. H. Dyson, and J. H. Fried, *Tetrahedron*, **28**, 3281 (1966); C. Beard, I. T. Harrison, L. Kirkham, and J. H. Fried, *ibid.*, **28**, 3287 (1966); T. L. Popper, F. E. Carlon, H. M. Marigliano, and M. D. Yudis, *Chem. Commun.*, 277 (1968); P. Hodge, J. A. Edwards, and J. H. Fried, *Tetrahedron Lett.*, 5175 (1966); P. Crabbé, H. Carpio, E. V. Velarde, and J. H. Fried, *J. Org. Chem.*, **38**, 1478 (1973).

(6) J. M. Birchall, G. W. Cross, and R. N. Haszeldine, *Proc. Chem. Soc., London*, 81 (1960).

(7) P. Crabbé, H. Carpio, A. Cervantes, J. Iriarte, and L. Tökes, *Chem. Commun.*, 79 (1968); P. Crabbé, A. Cervantes, A. Cruz, E. Galeazzi, J. Iriarte, and E. V. Velarde, *J. Amer. Chem. Soc.*, **95**, 6655 (1973).

(8) See, for example, S. A. Fuqua, W. G. Duncan, and R. M. Silverstein, *J. Org. Chem.*, **30**, 2543 (1965); D. J. Burton and F. E. Herkes, *Tetrahedron Lett.*, 1883 (1965); F. E. Herkes, Ph.D. Thesis, University of Iowa, 1966; D. J. Burton and F. E. Herkes, *J. Org. Chem.*, **32**, 1311 (1967); **33**, 1854 (1968).

(9) C. W. Rees and C. E. Smithen, *J. Chem. Soc.*, 938 (1964).

